



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

### About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>





600047143P





PLANE TRIGONOMETRY.



AN ELEMENTARY TREATISE  
ON  
PLANE TRIGONOMETRY;

WITH  
A Numerous Collection of Examples,  
CHIEFLY DESIGNED FOR THE USE OF SCHOOLS AND BEGINNERS.

BY  
R. D. BEASLEY, M.A.  
HEAD MASTER OF GRANTHAM GRAMMAR SCHOOL,  
FORMERLY FELLOW OF ST JOHN'S COLLEGE, CAMBRIDGE,  
AUTHOR OF 'ARITHMETIC FOR SCHOOLS.'


*EIGHTH EDITION.*

London:  
MACMILLAN AND CO.  
1883

[All Rights reserved.]

18381 . e . 1 .





**Cambridge :**

**PRINTED BY C. J. CLAY, M.A. & SON,  
AT THE UNIVERSITY PRESS.**

## PREFACE.

THE following pages are specially intended for use in Schools. In the choice of matter I have been chiefly guided by the requirements of the three days' Examination at Cambridge, with the exception of proportional parts in logarithms which I have omitted. In one point, I have ventured to deviate from the usual custom. I have denoted angles throughout by the Greek characters, even before the explanation of circular measure. My reasons for doing so were chiefly two; first, that they are much more distinctly and easily written, and it is advisable that a boy should be accustomed in the text to the notation he uses on his paper; secondly, I am thus enabled to insert the algebraical symbols for the angles in the figures, a very great advantage in such propositions as that proved in Art. 27. I have added about 400 Examples mainly collected from the Examination Papers of the last ten years, and I have taken great pains to exclude from the body of the work any which might dishearten a beginner by their difficulty.

In conclusion, I must express my great obligations to my friend the Rev. R. B. Mayor, of Rugby. Without his kind encouragement I should not have ventured to have offered these pages to the public; and I am indebted to him, not only for many valuable suggestions, but for a careful revision of the whole work.

R. D. BEASLEY.

ST JOHN'S COLLEGE, CAMBRIDGE,  
*September, 1858.*

## ADVERTISEMENT TO THE SECOND EDITION.

THE chief alterations in this Edition are the addition of Articles 63, bis, and 73—75, and of the easy series of Examples marked A.

GRAMMAR SCHOOL, GRANTHAM, *January, 1865.*

## ADVERTISEMENT TO THE THIRD EDITION.

In this Edition the Examples have been largely increased, Articles 76—78 have been added, and the series of Examination Papers.

GRAMMAR SCHOOL, GRANTHAM, *February, 1872.*

## ADVERTISEMENT TO THE EIGHTH EDITION.

In this Edition Section IX. on proportional parts has been added, and the number of Examination Papers considerably increased.

BOURNEMOUTH, *March, 1883.*

# CONTENTS.

SECTION	PAGE
I. Units of Measurement. Use of Signs + and -. Meaning of the term "Angle" in Trigonometry. The Trigonometrical Ratios. Practical Application. Instruments for Surveying . . . .	1
Examples. A. . . . .	12
" B. . . . .	15
" C. . . . .	18
II. Trigonometrical Ratios of Angles greater than 90°. Relations amongst the Ratios. Change in Sign and Magnitude of the Ratios through the Four Quadrants. Change from Positive to Negative Infinity. Other Trigonometrical Ratios . . . .	22
Examples. A. . . . .	31
" B. . . . .	33
" C. . . . .	35
III. Trigonometrical Ratios of the Sums and Differences of Angles, and of Multiple Angles. Explanation of the Double Sign in Trigonometrical Formulæ . . . . .	38
Examples. A. . . . .	47
" B. . . . .	51
" C. . . . .	56
IV. Relations between the Sides and Angles of a Triangle. The Radii of the Circumscribed and Inscribed Circles . . . . .	65
Examples. A. . . . .	70
" B. . . . .	73
" C. . . . .	76
V. Solution of Triangles. Four Cases. Their Treatment in Euclid. Application . . . . .	82
Examples. A. . . . .	87
" B. . . . .	89
" C. . . . .	91

SECTION	PAGE
VI. Logarithms. Their use. Advantage of Logarithms to Base 10.	
Arrangement of the Tables. Logarithms of the Trigonometrical Ratios . . . . .	94
Examples. A. . . . .	100
„ B. . . . .	101
„ C. . . . .	102
VII. Application of Logarithms to the Solution of Triangles; Illustration . . . . .	104
Examples. A. . . . .	108
„ B. . . . .	109
„ C. . . . .	110
VIII. Circular Measure. Polygons inscribed, and circumscribed about a Circle. Area of a Circle. General values of $\sin^{-1} a$ , &c. . . . .	111
Examples. A. . . . .	120
„ B. . . . .	122
„ C. . . . .	124
IX. On the use of proportional parts in Logarithmic and Trigonometrical Tables . . . . .	127
Examples A. . . . .	132
„ B. . . . .	133
„ C. . . . .	135
Miscellaneous Examples . . . . .	138
Examination Papers . . . . .	162

# PLANE TRIGONOMETRY.

## SECTION I.

UNITS OF MEASUREMENT. USE OF SIGNS + AND -. MEANING OF THE TERM "ANGLE" IN TRIGONOMETRY. THE TRIGONOMETRICAL RATIOS. PRACTICAL APPLICATION. INSTRUMENTS FOR SURVEYING.

1. *Object of Trigonometry.* In Trigonometry we apply Algebraical symbols to establish certain relations between the magnitudes of the sides and angles of plane rectilineal figures. These relations are useful for all the higher branches of Mathematics, and are specially applicable to surveying, and to determining the heights and distances of inaccessible objects. In the present treatise we shall confine ourselves to the simpler relations, and some practical applications of them.

We must first consider the mode of estimating algebraically the magnitudes of lines, areas, and angles.

2. *Measurement of lines.* As lines have neither breadth nor thickness we have only to measure their length. To do this we take some standard length, as one foot, one inch, five inches, or any other definite length as our unit of measurement; and the length of any line is then measured and represented by the number, whether whole or fractional, of these units which it contains. Thus if 5 inches is our unit, a line of 20 inches is measured by the number 4, and is called the line 4, that is, the line whose length is 4 times the unit of length. So also the line  $a$  would be the line whose length is  $a$  times the unit of length.

In investigations involving only algebraical symbols it is indifferent what unit is employed: but we must be careful to remember that throughout the same investigation we are using the same unit.

Thus if two lines  $a, b$  enter into our calculations we must consider the length of the line  $a$  to be  $a$  times *some* unit of length, and of the line  $b$  to be  $b$  times the *same* unit. When we apply our general results to numerical examples we must be careful to consider the special unit employed.

3. In the measurement of superficies we take the square of which one side is the unit of length, as the unit of superficies; and then any area is measured by the number of these units it contains.

N. B. The area of a rectangle whose sides contain  $a, b$  linear units respectively, contains  $ab$  superficial units. Also the areas of a parallelogram and a triangle whose bases are  $b$  and altitudes  $a$  are  $ab$  and  $\frac{ab}{2}$  respectively.

4. *Measurement of angles.* Since the idea of a right angle is simple, and all right angles are equal, it is conveniently taken as a standard by which to measure the magnitudes of other angles. But since most of the angles we have to deal with are less than right angles, and would therefore be represented by fractions or decimals, it is found convenient to divide it.

In England we divide the right angle into 90 equal parts called degrees; each degree is divided into 60 minutes; each minute into 60 seconds. Any angle is then measured by the number of degrees, minutes, seconds, and decimal parts of a second it contains; an angle containing 27 degrees, 13 minutes, 24.53 seconds is written thus,  $27^{\circ}, 13', 24''.53$ .

In France the decimal system is adopted. A right angle contains 100 grades, a grade 100 minutes, a minute 100 seconds; an angle containing 33 grades, 27 minutes, 45.5 seconds is written thus,  $33^g, 27', 45''.5$ . The advantage of this system is, that any angle can be reduced to the decimal of a grade at once, and *vice versa*. Thus  $33^g, 27', 45''.5 = 33^g.27455$ , and  $27^g.35679 = 27^g, 35', 67''.9$ .

5. We will now shew how to change our unit from degrees to grades, and *vice versa*.

Let  $D$  be the number of degrees in any angle  $ACB$ ,  $G$  the number of grades in the same angle; then since the number of

degrees in two angles must bear to one another the same ratio which the angles themselves do, we have

$$\frac{\text{angle } ACB}{\text{a right angle}} = \frac{D}{90};$$

for a like reason


$$\frac{\text{angle } ACB}{\text{a right angle}} = \frac{G}{100}.$$

Hence  $\frac{D}{90} = \frac{G}{100}$ : from which equation, if we know the number of degrees, we can find the number of grades, and *vice versa*.

There is also another mode of measuring angles, called *Circular measure*. For an account of this, the student may, if he pleases, turn at once to Section VIII, Articles 65—69, and to the Examples A, appended to that section.

6. *Use of signs + and - to represent contrariety of position.* In applying Algebra to the solution of Arithmetical Problems, we frequently meet with negative results, which require interpretation in each particular case; we shall meet with similar results in Trigonometry; and the question arises, how are we to interpret a negative quantity when it represents a line or angle?

Suppose  $BAB'$ ,  $DAD'$  to be two lines at right angles to one another and fixed in position.




Let  $AB = a$ ,  $BC = b$ ,  $AC = x$ ; then, representing the relation between these lines by Algebraical symbols, we have

$$x = a - b.$$



Now so long as  $b < a$ ,  $x$  is positive, and  $C$  lies to the right of  $A$ : but if  $b > a$ ,  $x$  is negative, and  $C$  lies to the left of  $A$ : that is, in using Algebraical symbols, if we arrive at a negative quantity representing a line, the natural interpretation appears to be, that the line must be considered as measured in an opposite direction to what it would be if the quantity representing it were positive. Hence we are led to the following convention: any line measured along  $AB$  or parallel to it is said to be a positive line, and any line measured along  $AB'$  or parallel to it is said to be a negative line, and the symbol representing it must have the sign  $-$  placed before it. So also lines measured along  $AD$  are positive, and along  $AD'$  negative. Lines measured in any other direction, not parallel to either of these two, will be considered positive. The advantage of these conventions will appear to the student as he proceeds.

7. *Magnitudes of angles unlimited in Trigonometry.*



In Euclid the magnitude of an angle is the absolute amount of the inclination of the two lines which contain it, to each other; this amount can never exceed two right angles. In Trigonometry the idea of an angle is extended. Suppose  $AOA'$  to be a line fixed

in space, and  $OP$  to be a line revolving from its original position  $OA$  towards  $OB$ , and tracing out the angle  $AOP$ : till  $OP$  reaches  $OA'$ , this angle does not differ from Euclid's idea of an angle, because it is less than two right angles; but when  $OP$  has passed  $OA'$ , and arrived at such a position as  $OP_3$ , the Trigonometrical idea of the angle differs from Euclid's: Euclid would now measure the angle  $AOP_3$  by the amount of inclination of  $OP_3$  to  $OA$ , that is, he would consider it as less than two right angles; but in Trigonometry we consider it as the angle actually described by the revolving line  $OP$ , that is, as  $180^\circ + \angle A'OP_3$ . So also when the revolving line reaches  $OP_4$ , the Trigonometrical angle  $AOP_4$  will be  $270^\circ + \angle B'OP_4$ ; and when  $OP$  has passed its initial position  $OA$  and reached  $OP_1$  a second time, the Trigonometrical angle  $AOP_1$  will be  $360^\circ + \angle AOP_1$ . Thus if  $AOP_1$  is  $30^\circ$ , then  $AOP_2$  is  $150^\circ$ ,  $AOP_3$  is  $210^\circ$ ,  $AOP_4$  is  $330^\circ$ ; and in the second revolution, when  $OP$  has reached  $OP_1$ , the  $\angle AOP_1$  is  $390^\circ$ . The same plan for estimating the value of the angle extends to any number of revolutions of  $OP$ , and in this sense the magnitude of an angle in Trigonometry is unlimited.

If the line  $OP$  on starting from  $OA$  had revolved in the opposite direction, towards  $OB'$ , then in accordance with the principle laid down in Art. 6, we should have considered the angle described as a negative angle; and we have the same convention for angles that we have for lines, viz. that angles measured upwards from  $OA$  towards  $OB$  are positive angles, and angles measured downwards from  $OA$  to  $OB'$  are negative angles, and have the sign  $-$  placed before the symbol representing them.

The same position of the line  $OP$  may be considered as making either a positive or negative angle with  $OA$ . Thus when  $OP$  is at  $OP_3$ , it may be supposed to have revolved either in a positive direction through  $OB$ ,  $OA_1$ , or in a negative direction through  $OB_1$ , i. e.,  $AOP_3$  may be either  $210^\circ$  or  $-150^\circ$ .

*Obs.* The space between  $OA$  and  $OB$  is called the first quadrant; when  $OP$  lies between  $OA$  and  $OB$ , the angle  $AOP$  is said to be an angle in the first quadrant. So also angles  $AOP_2$ ,  $AOP_3$ ,  $AOP_4$  are angles in the second, third, and fourth quadrants respectively.

8. *Complement; Supplement.* The complement of an angle is that angle which must be added to it to make it equal to a right angle. Thus the complement of

$30^\circ$  is  $60^\circ$ , of  $120^\circ$  is  $-30^\circ$ , of  $-75^\circ$  is  $165^\circ$ .

The supplement of an angle is that angle which must be added to it to make it equal to two right angles. Thus the supplement of

$30^\circ$  is  $150^\circ$ , of  $220^\circ$  is  $-40^\circ$ , of  $-75^\circ$  is  $255^\circ$ .

It is evident that in a right-angled triangle one of the acute angles is the complement of the other; and in any triangle any one of the angles is supplementary to the sum of the other two.

9. *Trigonometrical Ratios.* Having now shewn how lines and angles are to be measured in Trigonometry, and the interpretation which is to be given to the signs + and - when applied to quantities that represent them, we will proceed to shew how lines are connected with angles: this is done by means of what are called Trigonometrical Ratios.

Let the revolving line  $OP$  (fig. Art. 7) have come into any position  $OP_1, OP_2, \dots$ , and from  $P_1, P_2, \dots$ , drop the perpendiculars  $P_1N, P_2M, \dots$  upon the initial line  $OA$ , or  $OA$  produced backwards to  $A'$ ; the triangles  $P_1NO, P_2MO, \dots$  will be right-angled triangles. We then have the following definitions.

The ratio of the perpendicular to the hypotenuse is called the *sine* of the angle described: that is,

$$\frac{P_1N}{OP_1}, \frac{P_2M}{OP_2}, \dots$$

are the sines of the angles  $AOP_1, AOP_2, \dots$  respectively.

The ratio of the base to the hypotenuse is called the *cosine* of the angle, that is,

$$\frac{ON}{OP_1}, \frac{OM}{OP_2}, \dots$$

are the cosines of  $AOP_1, AOP_2, \dots$  respectively.

The ratio of the perpendicular to the base is called the *tangent* of the angle, that is,

$$\frac{P_1N}{ON}, \frac{P_2M}{OM}, \dots$$

are the tangents of  $\angle AOP_1, \angle AOP_2, \dots$  respectively.

The ratio of the base to the perpendicular is called the *cotangent* of the angle, that is,

$$\frac{ON}{P_1N}, \frac{OM}{P_2M}, \dots$$

are the cotangents of  $\angle AOP_1, \angle AOP_2, \dots$  respectively.

The ratio of the hypotenuse to the base is called the *secant* of the angle, that is,

$$\frac{OP_1}{ON}, \frac{OP_2}{OM}, \dots$$

are the secants of  $\angle AOP_1, \angle AOP_2, \dots$  respectively.

The ratio of the hypotenuse to the perpendicular is called the *cosecant* of the angle, that is,

$$\frac{OP_1}{P_1N}, \frac{OP_2}{P_2M}, \dots$$

are the cosecants of  $\angle AOP_1, \angle AOP_2, \dots$  respectively.

These definitions are quite independent of the direction and number of the revolutions of  $OP$  about  $O$ ; that is, they apply to negative angles, and to angles greater than four right angles.

The Trigonometrical Ratios of any angle  $\alpha$  (that is, of any angle which contains  $\alpha$  degrees, grades, or any other unit we may employ) are usually written

$$\sin \alpha, \cos \alpha, \tan \alpha, \cot \alpha, \sec \alpha, \operatorname{cosec} \alpha.$$


10. The question here naturally arises—are these Trigonometrical ratios definite quantities for any given value of the angle, or do they depend upon the magnitude of the revolving line  $OP$ ?

Let  $\angle AOP$  be any angle, i.e. let  $OP$  be any position whatever of the revolving line. Take  $Q$  any point in  $OP$ , or  $OP$  produced; draw  $PM$ ,  $QN$  perpendicular to  $OA$ .

Then by similar triangles  $PMO$ ,  $QNO$ ,

$$QN : ON : OQ = PM : OM : OP.$$

Hence the Trigonometrical Ratios of the angle  $\angle AOP$  are the same whether we consider  $OQ$  or  $OP$  the revolving line.



So also if  $RS$  is perpendicular to  $OP$ , we have

$$RS : OS : OR = PM : OM : OP.$$

Hence we conclude, that, *disregarding signs*, the values of the Trigonometrical Ratios are quite independent of the magnitude of the revolving line, and depend only on the absolute inclination of the two lines containing the angle to each other; and so for a given value of the angle are determinate.

The converse of this proposition (viz. that for a given value of the Trigonometrical Ratio, the angle is determinate) is only true with certain limitations, as will appear hereafter.

*Obs.* . The necessity of the proviso *disregarding signs* will appear on reading the next section.

11. Let the angle  $\angle AOP$  (fig. Art. 7) contain  $\alpha$  degrees, then  $\angle P_1OB$ , and  $\angle OP_1N$  (since they either of them with  $\angle AOP_1$  make a right angle) contain  $90 - \alpha$  degrees each. Drawing  $P_1N'$  perpendicular to  $OB$ , we have by our definitions

$$\sin P_1OB = \frac{P_1N'}{OP_1} = \frac{ON}{OP_1} = \cos \angle AOP_1,$$

$$\text{or } \sin (90^\circ - \alpha) = \cos \alpha.$$

Again, 
$$\tan P_1OB = \frac{P_1N'}{ON'} = \frac{ON}{P_1N} = \cot \angle AOP_1,$$

$$\text{or } \tan (90^\circ - \alpha) = \cot \alpha,$$

$$\text{so also } \sec (90^\circ - \alpha) = \operatorname{cosec} \alpha,$$

and these equations may be proved to exist for any position of the revolving line  $OP$ .

It is from these relations, viz. that the cosine, cotangent, cosecant of an angle, are respectively the same as the sine, tangent, and secant of the complement, that they derive their names.

From the same relations we also have

$$\sin a = \cos (90^\circ - a),$$

$$\tan a = \cot (90^\circ - a),$$

$$\sec a = \operatorname{cosec} (90^\circ - a).$$

Hence we can always express a Trigonometrical Ratio of the complement of an angle in terms of some Trigonometrical Ratio of the simple angle.

12. *The Trigonometrical Ratios of  $30^\circ$ ,  $45^\circ$ ,  $60^\circ$ .* The calculation of the values of the Trigonometrical Ratios of all angles is a matter of considerable labour, involving processes not described in this treatise, but they may be easily found in some particular cases.

Let  $ABC$  be an isosceles triangle having a right angle at  $C$ ; then  $CAB =$  half a right angle  $= 45^\circ$ .

Let  $AC = x = BC$ ,

$$\text{then } AB^2 = AC^2 + BC^2 = 2x^2$$

$$\text{or } AB = x\sqrt{2},$$

and we have

$$\sin 45^\circ = \frac{BC}{AB} = \frac{x}{x\sqrt{2}} = \frac{1}{\sqrt{2}},$$


$$\cos 45^\circ = \frac{AC}{AB} = \frac{x}{x\sqrt{2}} = \frac{1}{\sqrt{2}},$$

$$\tan 45^\circ = \frac{BC}{AC} = \frac{x}{x} = 1,$$

$$\cot 45^\circ = \frac{AC}{BC} = \frac{x}{x} = 1,$$

$$\sec 45^\circ = \frac{AB}{AC} = \frac{x\sqrt{2}}{x} = \sqrt{2},$$

$$\operatorname{cosec} 45^\circ = \frac{AB}{BC} = \frac{x\sqrt{2}}{x} = \sqrt{2}.$$



Again, let  $ABC$  be an equilateral triangle, then angle  $ABC = \frac{180^\circ}{3} = 60^\circ$ . Let  $AD$  be perpendicular to  $BC$ , then angle  $BAD = 30^\circ$ . Let  $BD = x$ ,  $AB = 2x$ ; then

$$AB^2 = AD^2 + BD^2,$$

$$\text{or } 4x^2 = AD^2 + x^2,$$

and therefore  $AD = x\sqrt{3}$ ; and we have,

$$\sin 60^\circ = \cos 30^\circ = \frac{AD}{AB} = \frac{x\sqrt{3}}{2x} = \frac{\sqrt{3}}{2},$$

$$\cos 60^\circ = \sin 30^\circ = \frac{BD}{AB} = \frac{x}{2x} = \frac{1}{2},$$

$$\tan 60^\circ = \cot 30^\circ = \frac{AD}{BD} = \frac{x\sqrt{3}}{x} = \sqrt{3},$$

$$\cot 60^\circ = \tan 30^\circ = \frac{BD}{AD} = \frac{x}{x\sqrt{3}} = \frac{1}{\sqrt{3}},$$

$$\sec 60^\circ = \operatorname{cosec} 30^\circ = \frac{AB}{BD} = \frac{2x}{x} = 2,$$



$$\operatorname{cosec} 60^\circ = \sec 30^\circ = \frac{AB}{AD} = \frac{2x}{x\sqrt{3}} = \frac{2}{\sqrt{3}}.$$

Thus we have found the values of all the Trigonometrical Ratios of  $30^\circ$ ,  $45^\circ$ , and  $60^\circ$ .

13. We will now assume that we know, and have arranged in Tables the values of the Trigonometrical Ratios of all angles, as well as of  $30^\circ$ ,  $45^\circ$ , and  $60^\circ$ , and shew how they may be used to solve many interesting problems.


Let  $ABC$  be a right-angled triangle,  $C$  the right angle, and let angles  $CAB$ ,  $CBA$  contain  $\alpha^\circ$ ,  $\beta^\circ$  respectively. Let the lengths of the sides opposite to  $A$ ,  $B$ ,  $C$  be  $a$ ,  $b$ ,  $c$  respectively. Then  $\sin \alpha = \frac{a}{c}$ , or

$$a = c \sin \alpha; \tan \alpha = \frac{a}{b}, \text{ or } a = b \tan \alpha;$$



$\cos a = \frac{b}{c}$ , or  $b = c \cos a$ ; also if  $p$  be the perpendicular from the angle  $C$  on the hypotenuse  $AB$ , we have  $\frac{p}{b} = \sin a = \frac{a}{c}$ , or  $cp = ab$ ; and similarly other relations may be found to exist between the sides and angles of a right-angled triangle. The student will best learn by practice to apply these relations to solve problems in each particular case; the following may be taken as an example.

A person on one side of a river at  $C$  wishes to determine the height of a tower  $AB$  on the other side. He observes, by an instrument made for the purpose, the angle which  $CA$  makes with the horizontal line  $CB$ . (Let this be  $a$ .) He then places a mark at  $D$ , and measures the distance  $CD$  ( $a$  feet) at right angles to  $CB$ , and when at  $D$  he observes the angle  $ADC$  ( $\beta$  suppose). He then calculates the height of the tower thus;  $ACD$  is a right-angled triangle, and therefore



$$AC = CD \tan ADC \quad \text{or} \quad AC = a \tan \beta.$$

So also  $ABC$  is a right-angled triangle, and therefore

$$AB = AC \sin ACB = AC \sin a.$$

But

$$AC = a \tan \beta,$$

$$\therefore AB = a \tan \beta \sin a.$$

Hence since we know the values of  $a$ ,  $\tan \beta$ ,  $\sin a$ , we can calculate the height of the tower.

$$\text{Ex. } a = 50 \text{ yards, } \beta = 45^\circ, a = 30^\circ,$$

$$\text{height of tower} = 50 \times 1 \times \frac{1}{2} = 25 \text{ yards.}$$

Instead of observing the  $\angle ADC$ , we might have observed the  $\angle BDC$  ( $\gamma$  suppose), and proceeded as follows,

$$BC = DC \tan BDC = a \tan \gamma,$$

$$AB = BC \tan ACB = a \tan \gamma \tan a.$$

This method would also give the breadth of the river,  $BC = a \tan \gamma$ .



14. The two principal instruments by which observations are taken in surveying are the sextant and the theodolite. The construction of these instruments depends on certain properties of light, the discussion of which falls under the science of optics. By the sextant we are enabled to observe the angle which any two objects subtend at the eye of the observer, as for instance the angle  $ADC$  in the case above.

By the theodolite we can observe the elevation of any object, that is to say, the angle which the line joining the object and the eye of the observer makes with the horizontal plane, as the angle  $ACB$  in the case above. We can also observe the angle which two objects in the same horizontal plane subtend at the eye of the observer; the sextant might do this, if we knew that the objects were in the same horizontal plane; the theodolite tells us that they are, as well as the amount of the angle they subtend. When an object is lower than the observer, the angle which the line joining it and the eye of the observer makes with the horizontal plane is called its depression.

### EXAMPLES. (A).

1. Find the number of degrees in the angle  $17^\circ, 34', 27''$ .

We have  $17^\circ, 34', 27'' = 17.3427,$

and  $\frac{D}{90} = \frac{G}{100},$  or  $D = \frac{9}{10} \times 17.3427 = 15.60843$  degrees

$$= 15^\circ, 36', 30''.348.$$

*Ans.*  $15^\circ, 36', 30''.348.$

2. If 7 feet is the unit of length, by what number is 1421 yards represented? *Ans.* 609.

3. If 3 inches is the unit of length, what is the unit of superficies? *Ans.* 9 square inches.

4. If 1728 square inches is represented by the number 108, what is the unit of length? *Ans.* 4 inches.

5. Reduce to grades, &c. the following angles,  $30^\circ$ ;  $22^\circ, 30'$ ;  $18^\circ$ ;  $29^\circ, 30'$ ;  $37^\circ, 22', 19''$ .

*Answers.*  $33^\circ, 33', 33''.3$ ;  $25^\circ$ ;  $20^\circ$ ;  $32^\circ, 77', 77''.7$ ;  $41^\circ, 52', 43''.44$ .

6. Reduce to degrees, &c. the angles,  $27^\circ, 5', 22''$ ;  $46^\circ, 17', 19''$ .

*Answers.*  $24^\circ, 20', 49''.128$ ;  $41^\circ, 33', 16''.956$ .

7. What is the unit of measurement when  $66\frac{2}{3}$  grades is represented by  $20$ ?

*Ans.* 3 degrees.

8. If  $\frac{1}{8}$ th of a right angle be the unit, what is the value of an angle  $5.09296$  in degrees?

*Ans.*  $57.2958$ .

9. Find all the angles less than a right angle, which can be expressed by an integral number both of degrees and grades.

10. Find the complements and supplements of the following angles,  $27^\circ$ ;  $29^\circ, 13', 15''$ ;  $47^\circ, 29', 47''$ ;  $107^\circ, 22'$ ;  $84^\circ, 10'$ .

11. One angle of a right-angled triangle is  $10^\circ$  greater than the other; find them.

*Ans.*  $40^\circ, 50^\circ$ .

12. One angle of a right-angled triangle is three times as great as the other; find them.

*Ans.*  $67^\circ, 30'$ ;  $22^\circ, 30'$ .

13. The three angles of a triangle are in A.P. and the common difference is  $12^\circ$ ; find them.

*Ans.*  $48^\circ, 60^\circ, 72^\circ$ .

14. Of the three angles of a triangle, one is double either of the other two; find them.

*Ans.*  $90^\circ, 45^\circ, 45^\circ$ .

15. The three angles of a triangle are in the ratio  $2 : 3 : 5$ ; find them.

*Ans.*  $36^\circ, 54^\circ, 90^\circ$ .

16. In a triangle, twice the difference between the first and third angle is equal to the sum of the second and third; and the third is equal to three times the difference between the first and second; find them.

*Ans.*  $80^\circ, 70^\circ, 30^\circ$ .

17. In the isosceles triangle of Euclid iv. 10, find the angles.

*Ans.*  $36^\circ, 72^\circ, 72^\circ$ .

18. The vertical angle of an isosceles triangle being  $80^\circ$ , find the other angles.

*Ans.*  $50^\circ$ .

19. The sides of a right-angled triangle are in A.P., shew that they are in the ratio 3 : 4 : 5.

20. If one angle of a right-angled triangle is  $39^\circ$ , and the hypotenuse is 7 feet, find the other sides.

*Ans.* 4.4052428, 5.440022 feet.

21. If one angle of a right-angled triangle is  $26^\circ$ , and the side adjacent to it 5 feet, find the other sides.

*Ans.* 5.5630095, 2.438663 feet.

22. If one angle of a right-angled triangle is  $37^\circ, 10'$ , and the side opposite to it 124 feet, find the other sides.

*Ans.* 205.25193, 163.5614684 feet.

23. If the two sides of a right-angled triangle are 30 feet and 40 feet, find the hypotenuse and the angles.

*Ans.* 50 feet ;  $36^\circ, 53'$  ;  $53^\circ, 7'$ .

24. If the hypotenuse and one side of a right-angled triangle are 13 feet and 5 feet, find the other side and the angles.

*Ans.* 12 feet ;  $67^\circ, 23'$  ;  $22^\circ, 37'$ .

25. If two sides of a right-angled triangle are 6 feet and 8 feet, find the perpendicular on the hypotenuse from the right angle.

*Ans.* 4.8 feet.

26. If one angle of a right-angled triangle is  $25^\circ$ , and the hypotenuse 7 feet, find the perpendicular on the hypotenuse.

*Ans.* 2.6811554.

27. A ladder is set 4 feet from a wall, and just reaches a window 20 feet from the ground, find its length.

*Ans.* 20.39... feet.

28. The length of an upright stick is 6 feet, and the length of its shadow 12 feet, find the altitude, or angle of elevation, of the sun.

*Ans.*  $26^\circ, 34'$ .

29. The length of a kite string is 300 yards, and the elevation of the kite is  $32^\circ$ , find its height.

*Ans.* 158.97579 yards.

30. A person in a balloon whose elevation is  $37^\circ$  drops a stone, which falls 560 yards from the observer, find the height of the balloon.

*Ans.* 421.990296 yards.

31. Construct the angles whose tangents are 3, and  $\frac{2}{3}$  respectively.

32. Construct the angles whose sines are  $\frac{1}{3}$  and  $\frac{3}{5}$  respectively.

33. A church tower on the side of a river is 200 feet high, and its elevation from the other side is  $29^\circ$ , find the breadth of the river.  
*Ans.* 360.80956 feet.

34. A hill whose slope makes an angle of  $12^\circ$  with the horizon is one mile long, find the height of the hill.

*Ans.* 1097.773776 feet.

35. If the road up the hill in the last question were made obliquely so as to be two miles long instead of one, find the inclination of the road to the horizon.  
*Ans.*  $5^\circ, 58'$ .

36. The elevation of a tower is  $30^\circ$  to a man 6 feet high, 140 feet from the foot of the tower, find its height.

*Ans.* 86.8 feet nearly.

EXAMPLES. (B).

1. Construct the angle whose sine is  $\frac{1}{\sqrt{3}}$ .

Take  $BD$  (fig. Art. 12)  $= x$ , then  $AD = x\sqrt{3}$ ; with centre  $B$ , distance equal to  $AD$ , describe a circle meeting  $AD$  in  $E$ , join  $BE$ , then

$$\sin BED = \frac{BD}{BE} = \frac{x}{x\sqrt{3}} = \frac{1}{\sqrt{3}},$$

and the angle  $BED$  has been found as required.

2. Find the linear unit, when an acre is 100,000 square units.  
*Ans.* 1 link.

3. If a rectangle 3 yards long by 2 broad be taken as the unit of superficies, what quantity will represent 2 square feet?

*Ans.*  $\frac{1}{27}$

4. How is 37 yards, 2 feet, 8 inches represented, when the unit of length is 4 inches? *Ans.* By 341.

5. If the angle of an equilateral triangle be taken as the unit of measurement, how many degrees will be contained in the angle  $\frac{1}{3}$ ?

6. Find the number of grades in  $27^\circ, 35', 24''$ .

*Ans.*  $30^\circ.65$ .

7. What is the supplement of  $227^\circ, 35', 46''$ ?

8. The complement of  $45^\circ \pm a$  is  $45^\circ \mp a$ .

9. The difference of two acute angles of a right-angled triangle is  $8^\circ$ , find the angles.

10. The vertical angle of an isosceles triangle is  $a^\circ$ , find the angles at the base.

11. The length of a person's shadow is twice his height, find the altitude of the sun, having given  $\tan 26^\circ, 34' = .5$ .

12. Shew that  $(\sin 60^\circ - \sin 45^\circ)(\cos 30^\circ + \cos 45^\circ) = \sin^2 30^\circ$ .

13. Shew that  $(\sin 30^\circ + \cos 30^\circ)(\sin 60^\circ - \cos 60^\circ) = \cos 60^\circ$ .

14. Shew that  $\frac{\sin 45^\circ - \sin 30^\circ}{\sin 45^\circ + \sin 30^\circ} = (\sec 45^\circ - \tan 45^\circ)^2$ .

15. Determine the height of an object whose elevation is  $40^\circ$ , the observer being 140 feet distant, and his eye 5 feet from the ground, having given that  $\tan 40^\circ = .85$ .

*Ans.* 124 feet.

16. Construct the angle whose tangent is  $\sqrt{2}$ .

17. Construct the angle whose sine is  $\frac{\sqrt{2}-1}{2}$ .

18. A ladder of length  $l$  is placed against a wall so that the angle it makes with the ground is double that which it makes with the wall: find how far from the wall the foot of the ladder is.

*Ans.*  $\frac{l}{2}$ .

19. The town *C* is halfway between the towns *D* and *E*; and the towns *C*, *E*, *F* are equidistant from each other. If distance from *D* to *E* is 12 miles, find the distance from *D* to *F*.

*Ans.*  $6\sqrt{3}$  miles.

20. A ladder 20 feet long just reaches the top of a wall when its foot is 13 feet from the foot of the wall; shew that when its foot is 5 feet from the wall, the ladder projects 4 feet beyond the top of the wall.

21. In a right-angled triangle the lengths of lines drawn from the acute angles to the middle points of the opposite sides are  $d$ ,  $d'$  respectively. Find the sides of the triangle.

$$\text{Ans. } \frac{2}{\sqrt{15}} \sqrt{(4d^2 - d'^2)}, \frac{2}{\sqrt{15}} \sqrt{(4d'^2 - d^2)}.$$

22. A person travelling southwards observes two objects towards the S.E. After 8 miles travelling, one of them is N.E., and the other E. Their distances from him are then  $4\sqrt{2}$  miles, and 8 miles respectively.

23. Whilst sailing due West, I observe two ships at anchor directly North of me: after sailing 6 miles the directions of the ships make angles  $60^\circ$ ,  $30^\circ$  with my course respectively. The distance between them is  $4\sqrt{3}$  miles.

24. A staff 1 foot long stands on the top of a tower 200 feet high. Shew that the angle it subtends at a place 100 feet from the foot of the tower is  $6'$ .

$$\text{Given } \tan 63^\circ.27' = 2, \tan 63^\circ.33' = 2.01.$$

25. The angles of depression of the top and bottom of a column observed from a tower 108 feet high are  $30^\circ$ ,  $60^\circ$  respectively. Shew that the height of the column is 72 feet.

26. From the top of a column 100 feet high, the angles of depression of two objects in a line with the column, and in the horizontal plane on which the column stands, are observed to be  $30^\circ$  and  $60^\circ$ . The distance between them is  $\frac{200}{\sqrt{3}}$  feet.

27. The length of a road, in which the ascent is 1 foot in 5, from the foot of a hill to the top is a mile and two-thirds : what will be the length of a zigzag road in which the ascent is 1 foot in 12? *Ans.* 4 miles.

### EXAMPLES. (C).

1. The same line is represented by  $m$  and  $n$  in two systems of measurement, compare the units of length in the two systems.

2. What is the greatest unit of length with which .625 feet may be represented by an integer?

3. Find the number of French minutes in one English minute, and the number of French seconds in one English second.

*Ans.*  $1^{\circ}.851$ ,  $3^{\circ}.086419753$ .

4. The angles of a triangle are in A. P.; shew that one of them must equal  $60^{\circ}$ .

5. The supplement of one angle of a triangle is double the complement of another, and triple that of the third: find the angles.

*Ans.*  $81\frac{2}{11}$ ,  $40\frac{1}{11}$ ,  $57\frac{3}{11}$  degrees.

6. If with two units of angular measurement differing by  $10^{\circ}$  the measures of an angle are as 3 : 2, what are those units?

*Ans.*  $20^{\circ}$ ,  $30^{\circ}$ .

7. If the measure of an angle equal the sum of the number of degrees and half the grades in it, what is the unit of angular measure?

*Ans.*  $\left(\frac{9}{14}\right)^{\circ}$ .

8. Three angles are in A. P. The number of grades in the greatest is equal to the number of degrees in the sum of the other two: shew that the angles are in the ratio 11 : 19 : 27.

9. In an isosceles triangle whose sides are  $a$ ,  $a$ ,  $c$ , if  $p$  be the perpendicular from one of the angles at the base on the opposite side, then  $p = \frac{c\sqrt{4a^2 - c^2}}{2a}$ .

10. A ladder (whose inclination to the ground is  $\alpha$ ) just reaches the top of a wall; at a distance  $a$  feet up the ladder the wall subtends an angle  $2\alpha$ ; shew that the height of the wall is  $2a \sin \alpha$ .

11. A person observes the angular elevation of a column; after approaching  $a$  feet, he finds its elevation doubled; again approaching  $b$  feet it is again doubled. Shew that the second point of observation is  $\frac{a^2}{2b}$  feet from the foot of the tower.

If  $\alpha = b \sqrt{3}$ , shew that the first elevation was  $15^\circ$ .

12. Construct the angle whose tangent is  $\sqrt{5} - 1$ .

13. At the foot of a mountain the elevation of its summit is  $45^\circ$ . After ascending 1 mile up a slope of  $30^\circ$  its elevation is found to be  $60^\circ$ . The height of the mountain is  $\frac{\sqrt{3}+1}{2}$  miles.

14. A person at the edge of a river observes the elevation  $\alpha$  of a tower on the other side; on retreating  $a$  feet he finds its elevation to be  $\beta$ : the height of the tower is  $\frac{a \tan \alpha \tan \beta}{\tan \alpha - \tan \beta}$ , the breadth of the river is  $\frac{a \tan \beta}{\tan \alpha - \tan \beta}$ .

15. A rock is observed from a ship to bear N. N. W.; after sailing 10 miles in direction E. N. E. the rock is due West: its distance from the ship at the first observation was  $10(\sqrt{2}-1)$  miles. Given  $\tan 22\frac{1}{2}^\circ = \sqrt{2}-1$ .

16. A ship sailing North sees two lighthouses due East. After sailing an hour they are S. E., and S. S. E.; the distance between them is 8 miles: find the rate of the ship.

*Ans.* 13.6 miles nearly.

17. Two persons  $A$  and  $B$  start from two points distant 400 yards.  $B$  starts at right angles to the line joining the two points at the rate of 90 yards a minute.  $A$  starts in a direction to catch  $B$  as soon as possible at the rate of 150 yards a minute; find how



long he will be before he catches him, and the direction in which he will walk, having given  $\sin 36^\circ.53' = \frac{3}{5}$ .

*Ans.* 3 minutes, 20 seconds.

18. A tower is situated at the top of a hill whose inclination is  $30^\circ$ . The angle subtended by the tower at the foot of the hill is  $15^\circ$ ; and on walking  $a$  yards up the hill it is found to be  $30^\circ$ . The height of the tower is  $\frac{a}{\sqrt{3}}$  feet.

19. From two points in the diameter of a circle produced tangents are drawn to the circle. Given the distance  $c$  between the points, and the inclinations  $\alpha, \beta$  of the tangents to the diameter, shew that the radius of the circle is  $\frac{c}{\operatorname{cosec} \beta - \operatorname{cosec} \alpha}$ .

20. A person travelling along a road observes the elevation  $\alpha$  of a tower the nearest distance ( $a$ ) of which from the road is known; at the same time he observes the angular distance  $\beta$  of the top of the tower from an object in the road. The height of the tower is  $\frac{a \sin \alpha}{\sqrt{(\cos^2 \alpha - \cos^2 \beta)}}$ .

21. At three positions in the same horizontal plane distant from each other 60, 80, 100 feet respectively, the elevation of a tower is observed to be  $45^\circ$ ; find its height. *Ans.* 50 feet.

22. Two spectators at two stations, distant  $2a$  from each other, observe the elevation of a kite to be  $\alpha$  at each station, and the angle subtended by the kite and the other station to be  $\beta$ ; shew that the height of the kite is  $a \sec \beta \sin \alpha$ .

23. A person standing on the top  $A$  of a light-house  $AB$ , of known height 300 feet, observes a ship sailing from  $C$  to  $D$  in a straight line: he knows that  $CD$  is perpendicular to the plane  $ACB$ , and he observes the angles  $CAD = 30^\circ$ ,  $BAD = 60^\circ$ . Shew that  $CD = 300$  feet;  $BD = 300\sqrt{3}$  feet.

24. The altitude of the sun is  $45^\circ$ , and it is at the point of the compass  $60^\circ$  from the south. The breadth of the shadow of a south wall is one-half its height.

25. The angular altitude and breadth of a cylindrical tower are observed to be  $\alpha$  and  $\beta$  respectively; but at a point  $a$  feet nearer the foot of the tower they are  $\alpha'$ ,  $\beta'$  respectively: find the height and radius of the tower.

$$\text{Ans. Height} = \frac{a}{\cot \alpha - \cot \alpha'}, \text{ radius} = \frac{a}{\operatorname{cosec} \frac{\beta}{2} - \operatorname{cosec} \frac{\beta'}{2}}.$$

26. A person  $a$  feet from the foot of a tower (height  $h$ ) sees a mountain summit in a line with the top of the tower. At the foot of the tower the elevation of the mountain is  $\alpha$ : shew that the height of the mountain is

$$\frac{ah}{a - h \cot \alpha}.$$

27. An embankment (the height of whose summit is  $c$ ) stands on a plain, and a man looking directly up the slope of the embankment just sees the top of a tower on the plain: when at distance  $a$  from the summit the man's eye is level with the summit, and when at the distance  $b$  from the summit he can just see the foot of the tower. Shew that the height of the tower is  $\frac{ac}{a - b}$ .

## SECTION II.

TRIGONOMETRICAL RATIOS OF ANGLES GREATER THAN  $90^\circ$ .  
RELATIONS AMONGST THE RATIOS. CHANGE IN SIGN AND  
MAGNITUDE OF THE RATIOS THROUGH THE FOUR QUAD-  
RANTS. CHANGE FROM POSITIVE TO NEGATIVE IN-  
FINITY. OTHER TRIGONOMETRICAL RATIOS.

15. We now proceed to express the Trigonometrical Ratios of all angles in terms of Trigonometrical Ratios of positive angles less than a right angle.

We will first find the Trigonometrical Ratios of the supplement of an angle.

Let  $A'OP_2 = AOP_1$  absolutely (fig. Art. 7): then  $AOP_2$  is the supplement of  $AOP_1$  or  $\alpha$ , that is,  $AOP_2 = 180^\circ - \alpha$ .

Let  $ON = x$ ,  $P_1N = y$ ,  $OP_1 = r$ ,  
then  $OM = -x$ ,  $P_2M = y$ ,  $OP_2 = r$ .

Hence  $\sin(180^\circ - \alpha) = \frac{y}{r} = \sin \alpha$ ,

$$\cos(180^\circ - \alpha) = \frac{-x}{r} = -\cos \alpha,$$

$$\tan(180^\circ - \alpha) = \frac{y}{-x} = -\tan \alpha,$$

$$\cot(180^\circ - \alpha) = \frac{-x}{y} = -\cot \alpha,$$

$$\sec(180^\circ - \alpha) = \frac{r}{-x} = -\sec \alpha,$$

$$\operatorname{cosec}(180^\circ - \alpha) = \frac{r}{y} = \operatorname{cosec} \alpha;$$

which equations give the Trigonometrical Ratios of the supplement of an angle, in terms of the Trigonometrical Ratios of the simple angle.

We may observe that they all change in sign except the sine and cosecant.

16. We will next find the Trigonometrical Ratios of a negative angle.

Let  $\angle OP_4 = \angle OP_1$ , absolutely: then  $\angle OP_4 = -\alpha$ ;  
also  $ON = x$ ,  $P_4N = -y$ ,  $OP_4 = r$ .

$$\text{Hence} \quad \sin(-\alpha) = \frac{-y}{r} = -\sin \alpha,$$

$$\cos(-\alpha) = \frac{x}{r} = \cos \alpha,$$

$$\tan(-\alpha) = \frac{-y}{x} = -\tan \alpha,$$

$$\cot(-\alpha) = \frac{x}{-y} = -\cot \alpha,$$

$$\sec(-\alpha) = \frac{r}{x} = \sec \alpha,$$

$$\operatorname{cosec}(-\alpha) = \frac{r}{-y} = -\operatorname{cosec} \alpha;$$

which equations give the Trigonometrical Ratios of a negative angle in terms of those of the positive angle of the same absolute magnitude.

We may observe that they all change in sign except the cosine and secant.

17. Let  $\angle A'OP_2 = \angle OP_1 = \alpha$ , then  $\angle AOP_2 = 180^\circ + \alpha$ ,  
also  $OM = -x$ ,  $P_2M = -y$ ,  $OP = r$ .

$$\text{Hence} \quad \sin(180^\circ + \alpha) = \frac{-y}{r} = -\sin \alpha,$$

$$\cos(180^\circ + \alpha) = \frac{-x}{r} = -\cos \alpha,$$

$$\tan(180^\circ + \alpha) = \frac{-y}{-x} = \tan \alpha,$$

$$\cot(180^\circ + \alpha) = \frac{-x}{-y} = \cot \alpha,$$

$$\sec(180^\circ + \alpha) = \frac{r}{-x} = -\sec \alpha,$$

$$\operatorname{cosec}(180^\circ + \alpha) = \frac{r}{-y} = -\operatorname{cosec} \alpha.$$

Here all the ratios change in sign except the tangent and cotangent.

18. It is evident that, when the revolving line  $OP$  has reached  $OP$ , the second time, that is, has described the angle  $(360^\circ + a)$ , all the Trigonometrical Ratios are the same as those of the angle  $a$ , for the triangle which determines their values is identical in the two cases. The same remark applies to any number of revolutions, so that we may say generally that the Trigonometrical Ratios of  $(n \cdot 360^\circ + a)$  are equal severally to those of  $a$ .

19. The formulæ in Articles 12, 15, 16, 17, 18, have been proved for the case in which  $a$  is less than  $90^\circ$ ; that they are true for all values whatever of  $a$  the student may satisfy himself by drawing figures to represent the different cases: for a more logical proof that these forms are true for all angles whatever he must be content to wait till he is further advanced, as the reasoning is of too subtle a nature for the present.

These formulæ may also be classed as follows :

$$\begin{aligned}\sin a &= \sin(180^\circ - a) = -\sin(180^\circ + a) = -\sin(-a) \\ &= \sin(n \cdot 360^\circ + a), \\ \cos a &= -\cos(180^\circ - a) = -\cos(180^\circ + a) = \cos(-a) \\ &= \cos(n \cdot 360^\circ + a), \\ \tan a &= -\tan(180^\circ - a) = \tan(180^\circ + a) = -\tan(-a) \\ &= \tan(n \cdot 360^\circ + a), \\ \cot a &= -\cot(180^\circ - a) = \cot(180^\circ + a) = -\cot(-a) \\ &= \cot(n \cdot 360^\circ + a), \\ \sec a &= -\sec(180^\circ - a) = -\sec(180^\circ + a) = \sec(-a) \\ &= \sec(n \cdot 360^\circ + a), \\ \operatorname{cosec} a &= \operatorname{cosec}(180^\circ - a) = -\operatorname{cosec}(180^\circ + a) = -\operatorname{cosec}(-a) \\ &= \operatorname{cosec}(n \cdot 360^\circ + a).\end{aligned}$$

We obtain also from the above formulæ the signs which the Trigonometrical Ratios of any angles must have according to the quadrant in which their bounding line is situated.

20. We are now in a position to express the Trigonometrical Ratios of any angle whatever in terms of the Ratios of a positive angle less than  $90^\circ$ .

If the angle is negative make it positive by formulæ in Art. 16.

If the angle is greater than  $360^\circ$ , suppress  $360^\circ$  as often as necessary to make it less than  $360^\circ$  by Art. 18.

If the angle is now greater than  $180^\circ$ , suppress  $180^\circ$  by Art. 17.

If the angle is now greater than  $90^\circ$  take its supplement by Art. 15.

When the angle is negative labour is sometimes saved by adding  $360^\circ$  once or oftener, till the angle is positive, instead of using Art. 16.

Ex.  $\tan 995^\circ = \tan 275^\circ = \tan 95^\circ = -\tan 85^\circ$ ,  
 $\operatorname{cosec}(-995^\circ) = -\operatorname{cosec} 995^\circ = -\operatorname{cosec} 275^\circ = \operatorname{cosec} 95^\circ = \operatorname{cosec} 85^\circ$ ,  
 or thus,  $\operatorname{cosec}(-995^\circ) = \operatorname{cosec}(1080^\circ - 995^\circ) = \operatorname{cosec} 85^\circ$ .

21. From the formulæ in Arts. 15, 16, 17, 18 we deduce the following, which are true for all values of the angle,

$$\sin a \operatorname{cosec} a = 1,$$

$$\cos a \sec a = 1,$$

$$\tan a \cot a = 1.$$

$$\text{Also in the first quadrant } \tan a = \frac{y}{x} = \frac{\frac{y}{r}}{\frac{x}{r}} = \frac{\sin a}{\cos a},$$

$$\text{in the second quadrant } \tan a = \frac{y}{-x} = \frac{\frac{y}{r}}{-\frac{x}{r}} = \frac{\sin a}{\cos a},$$

$$\text{in the third quadrant } \tan a = \frac{-y}{-x} = \frac{-\frac{y}{r}}{-\frac{x}{r}} = \frac{\sin a}{\cos a},$$

$$\text{in the fourth quadrant } \tan a = \frac{-y}{x} = \frac{-\frac{y}{r}}{\frac{x}{r}} = \frac{\sin a}{\cos a}.$$

nititude of the Trigonometrical Ratios as the angle (fig. Art. 7)  $AOP$  or  $\alpha$  passes through all values from  $0^\circ$  to  $360^\circ$ . Let

$$ON = x, \quad PN = y, \quad OP = r,$$

for all positions of  $OP$ : then  $r$  is always positive and never less than  $x$  and  $y$ , and therefore the sine and cosine are never greater than 1, and the secant and cosecant never less than 1: and since  $x$  and  $y$  may have any ratio to each other, the tangent and cotangent may have any values whatever.

When  $OP$  coincides with  $OA$ ,  $x = r$ ,  $y = 0$ .

$$\text{Hence, } \sin 0^\circ = \frac{0}{r} = 0; \quad \tan 0^\circ = \frac{0}{r} = 0; \quad \sec 0^\circ = \frac{r}{r} = 1;$$

$$\cos 0^\circ = \frac{r}{r} = 1; \quad \cot 0^\circ = \frac{r}{0} = \infty; \quad \operatorname{cosec} 0^\circ = \frac{r}{0} = \infty.$$

As  $OP$  revolves from  $OA$  to  $OB$ ,  $x$  diminishes,  $y$  increases, and both are positive; hence the sine, tangent, secant increase, the cosine, cotangent, cosecant diminish, and they are all positive.

When  $OP$  coincides with  $OB$ ,  $x = 0$ ,  $y = r$ .

$$\text{Hence, } \sin 90^\circ = \frac{r}{r} = 1; \quad \tan 90^\circ = \frac{r}{0} = \infty; \quad \sec 90^\circ = \frac{r}{0} = \infty;$$

$$\cos 90^\circ = \frac{0}{r} = 0; \quad \cot 90^\circ = \frac{0}{r} = 0; \quad \operatorname{cosec} 90^\circ = \frac{r}{r} = 1.$$

As  $OP$  revolves from  $OB$  to  $OA'$ ,  $x$  increases and is negative,  $y$  diminishes and is positive; hence the sine, tangent, secant diminish, the cosine, cotangent, cosecant increase; also the sine and cosecant are positive, the rest negative.

When  $OP$  coincides with  $OA'$ ,  $x = -r$ ,  $y = 0$ .

Hence,

$$\sin 180^\circ = \frac{0}{r} = 0; \quad \tan 180^\circ = \frac{0}{-r} = 0; \quad \sec 180^\circ = \frac{r}{-r} = -1;$$

$$\cos 180^\circ = \frac{-r}{r} = -1; \quad \cot 180^\circ = \frac{-r}{0} = -\infty; \quad \operatorname{cosec} 180^\circ = \frac{r}{0} = \infty.$$

As  $OP$  revolves from  $OA'$  to  $OB$ ,  $x$  diminishes and is negative,  $y$  increases and is negative; hence the sine, tangent, secant increase,

and the cosine, cotangent, cosecant diminish, also the tangent and cotangent are positive, the rest negative.

When  $OP$  coincides with  $OB'$ ,  $x = -0$ ,  $y = -r$ .

Hence,

$$\sin 270^\circ = \frac{-r}{r} = -1; \tan 270^\circ = \frac{-r}{-0} = \infty; \sec 270^\circ = \frac{r}{-0} = -\infty;$$

$$\cos 270^\circ = \frac{-0}{r} = 0; \cot 270^\circ = \frac{-0}{-r} = 0; \operatorname{cosec} 270^\circ = \frac{r}{-r} = -1.$$

As  $OP$  revolves from  $OB'$  to  $OA$ ,  $x$  increases and is positive, and  $y$  diminishes and is negative. Hence the sine, tangent, secant diminish, and the cosine, cotangent, cosecant increase; also the cosine and secant are positive, the rest negative.

When  $OP$  coincides with  $OA$  the second time, the values of the Trigonometrical Ratios become equal to those of  $0^\circ$ .

24. We have used above the relation  $x = -0$ , and so made

$$\sec 270^\circ = \frac{r}{x} = \frac{r}{-0} = -\infty,$$


instead of  $+\infty$  which it would have been if we had put  $x = 0$ . The reason of this is, that  $x$  has been continually diminishing and negative, and therefore the secant has been a continually increasing negative quantity; and we have thought it better to say that as the angle reaches  $270^\circ$ , the secant becomes an infinitely great negative quantity. If however the line  $OP$  had approached the position  $OB'$  from  $OA$ , by revolving in the negative direction, the secant of the angle  $-90^\circ$  would be  $+\infty$ .

Thus for the same position of  $OP$ , supposed to revolve in opposite directions to reach this position, we have two values of the secant of the angle of opposite signs. In fact, positive and negative infinity seem to have a near approach to each other, for we find the Ratios which become infinite always pass from one to the other. The case above and similar ones are only in accordance with our preceding formulæ; for since  $270^\circ$  and  $-90^\circ$  are supplementary angles, it follows that their secants should be equal and of opposite signs.



The following illustration may help to remove the difficulty from the Student's mind.

Let  $CA$  be drawn perpendicular to  $AB$  a line of unlimited length,  $CD$  drawn through  $C$  to meet  $AB$  in  $D$ . Let  $AD = x$



Now if angle  $ACD$  is made gradually to increase, the value of  $x$  increases without limit and is positive till  $ACD$  becomes a right angle. As soon as  $ACD$  becomes greater than a right angle, the value of  $x$  becomes suddenly an infinitely large negative quantity, and then diminishes to zero. The change in position of the point  $D$  for a small change in the revolving line, becomes infinitely great at this position of the line, though in general it is but gradual.

25. Besides the Trigonometrical Ratios above enumerated, the following are sometimes used.

The versine of the angle, which equals  $1 - \cos$ , that is,

$$\text{vers } \alpha = 1 - \cos \alpha.$$

The coversine, which is the versine of the complement, that is,

$$\text{covers } \alpha = 1 - \cos (90 - \alpha) = 1 - \sin \alpha.$$

The suversine, which is the versine of the supplement, that is,

$$\text{suvers } \alpha = 1 - \cos (180 - \alpha) = 1 + \cos \alpha.$$

Also the chord of the angle, which is twice the sine of half the angle, that is,

$$\text{chd } \alpha = 2 \sin \frac{\alpha}{2}.$$

If we turn to fig. Art. 7, we see that

$$\text{vers } \alpha = \frac{AN}{AO}, \quad \text{covers } \alpha = \frac{BN'}{AO}, \quad \text{suvers } \alpha = \frac{A'N}{AO};$$

and if  $AP$  be joined, chord  $\alpha = \frac{AP}{AO}$ .

26. The following notation is also used: 'if the sine of the angle  $a$  is  $a$ , that is, if  $\sin a = a$ , then the angle whose sine is  $a$  is written  $\sin^{-1}a$ , that is,  $a = \sin^{-1}a$ ; and so for the other Trigonometrical Ratios.

Also, since  $a = \sin^{-1}a$ ;  $\therefore \sin a = a$ ,  $\cos a = \sqrt{1-a^2}$ ,

$$\tan a = \frac{a}{\sqrt{1-a^2}}. \quad \text{Hence } a = \sin^{-1}a = \cos^{-1} \sqrt{1-a^2} = \tan^{-1} \frac{a}{\sqrt{1-a^2}}.$$

EXAMPLES. (A).

1. Express in the corresponding Trigonometrical Ratios of angles less than  $90^\circ$ ,  $\sin 732^\circ$ ,  $\sin 2451^\circ$ ,  $\cos 674^\circ$ ,  $\cos (-424^\circ)$ ,  $\tan 582^\circ$ ,  $\tan (-1975^\circ)$ ,  $\cot 873^\circ$ ,  $\operatorname{cosec} 565^\circ$ ,  $\sec (-1892^\circ)$ .

*Ans.*  $\sin 12^\circ$ ,  $-\sin 69^\circ$ ,  $\cos 46^\circ$ ,  $\cos 64^\circ$ ,  $\tan 42^\circ$ ,  $\tan 5^\circ$ ,  $-\cot 27^\circ$ ,  $-\operatorname{cosec} 25^\circ$ ,  $-\sec 88^\circ$ .

2. If  $\sin a = \frac{3}{5}$ , find  $\cos a$ ,  $\sec a$ ,  $\tan a$ .

$$\text{Ans. } \cos a = \frac{4}{5}, \sec a = 1\frac{1}{4}, \tan a = \frac{3}{4}.$$

3. If  $\cos a = \frac{5}{13}$ , find  $\sin a$ ,  $\operatorname{cosec} a$ ,  $\cot a$ .

$$\text{Ans. } \sin a = \frac{12}{13}, \operatorname{cosec} a = 1\frac{1}{12}, \cot a = \frac{5}{12}.$$

4. If  $\sec a = 3\frac{4}{7}$ , find  $\tan a$ ,  $\sin a$ .

$$\text{Ans. } \tan a = 3\frac{3}{7}, \sin a = \frac{24}{25}.$$

5. If  $\tan a = 2\frac{2}{5}$ , find  $\sec a$ ,  $\sin a$ .

$$\text{Ans. } \sec a = 2\frac{3}{5}, \sin a = \frac{12}{13}.$$

6. If  $\sin a = \frac{1}{3}$ , find the other Ratios.

$$\text{Ans. } \cos a = \frac{2}{3}\sqrt{2}, \tan a = \frac{1}{4}\sqrt{2}, \cot a = 2\sqrt{2}, \sec a = \frac{3}{4}\sqrt{2}, \operatorname{cosec} a = 3.$$

7. If  $\tan a = \frac{2}{3}$ , find  $\sec a$ ,  $\sin a$ ,  $\text{vers } a$ .

$$\text{Ans. } \sec a = \frac{\sqrt{13}}{3}, \sin a = \frac{2}{\sqrt{13}}, \text{vers } a = \frac{\sqrt{13}-3}{\sqrt{13}}.$$

8. In questions 4, 5, 6, 7 find the values of  $a$  from the Tables.

$$\text{Ans. } 73^\circ, 45'; 67^\circ, 23'; 19^\circ, 28'; 33^\circ, 41'.$$

Prove the truth of the following equations.

9.  $\sin a \text{ covers } a (1 + \text{cosec } a) = \cos^2 a.$

Here

$$\sin a \text{ covers } a (1 + \text{cosec } a)$$

$$= \sin a (1 - \sin a) \left( \frac{1 + \sin a}{\sin a} \right) = 1 - \sin^2 a$$

$$= \cos^2 a. \quad \text{Q.E.D.}$$

10.  $\cos a \tan a + \sin a \cot a = \sin a + \cos a.$

11.  $\cos a \text{ cosec } a = \cot a.$

12.  $(1 + \tan^2 a) \cos^2 a = 1.$

13.  $\frac{\sin a \sec a}{\tan a} = 1.$

14.  $(\cos a \tan a)^2 + (\sin a \cot a)^2 = 1.$

15.  $\frac{\cos a}{\sin a \cot^2 a} = \tan a.$

16.  $\frac{\sec^2 a - 1}{\sec^2 a} = \sin^2 a.$

17.  $\frac{\text{cosec}^2 a - 1}{\text{cosec}^2 a} = \cos^2 a.$

18.  $(\sec^2 a - 1)(\text{cosec}^2 a - 1) = 1.$

19.  $\sec a \text{ cosec } a (\cos^2 a - \sin^2 a) = \cot a - \tan a.$

20.  $\frac{\sin a \sec a}{\cos a \text{ cosec } a} = \frac{\tan a}{\cot a}.$

21.  $\frac{\sin a \cos a}{\sec a \text{ cosec } a} = \sin^2 a - \sin^4 a.$

22.  $\sec^2 a + \text{cosec}^2 a = \sec^2 a \text{ cosec}^2 a.$

23.  $\cot^2 a - \cos^2 a = \cos^2 a \cot^2 a.$
24.  $\cos^4 a - \sin^4 a = \cos^2 a - \sin^2 a.$
25.  $\tan a + \cot a = \sec a \operatorname{cosec} a.$
26.  $\cos a \operatorname{vers} a (1 + \sec a) = \sin^2 a.$
27.  $\sin^2 a (1 + n \cot^2 a) = \cos^2 a (n + \tan^2 a).$
28.  $(\sin a + \cos a)(\tan a + \cot a) = \sec a + \operatorname{cosec} a.$
29.  $\tan(-a) \sec(-a) = \frac{\tan a}{\operatorname{vers} a - 1}.$
30. If  $\tan(a + \beta) = \tan(\theta + \phi)$ , then  $\tan(a - \theta) = \tan(\phi - \beta).$

EXAMPLES. (B).

1. If  $\tan a = \frac{b}{a}$ , then  $\sin a = \frac{b}{\sqrt{a^2 + b^2}}$ ,  $\cos a = \frac{a}{\sqrt{a^2 + b^2}}$ ,  
 for  $\sin a = \frac{\tan a}{\sqrt{1 + \tan^2 a}} = \frac{\frac{b}{a}}{\sqrt{1 + \frac{b^2}{a^2}}} = \frac{b}{\sqrt{a^2 + b^2}},$   
 $\cos a = \frac{1}{\sqrt{1 + \tan^2 a}} = \frac{1}{\sqrt{1 + \frac{b^2}{a^2}}} = \frac{a}{\sqrt{a^2 + b^2}}.$
2. If  $\tan \theta = \frac{1 + \tan a}{1 - \tan a}$ , find  $\sin \theta$ ,  $\cos \theta$ ,  
 $\sin \theta = \frac{1 + \tan a}{\sqrt{\{2(1 + \tan^2 a)\}}} = \frac{1 + \tan a}{\sec a \cdot \sqrt{2}} = \frac{1}{\sqrt{2}} (\cos a + \sin a),$   
 $\cos \theta = \frac{1 - \tan a}{\sqrt{\{2(1 + \tan^2 a)\}}} = \frac{1}{\sqrt{2}} (\cos a - \sin a).$
3. If  $\sin a = \frac{3}{4}$ , shew that  $\tan a = \frac{1}{4} \sqrt{2}$ ,  $\cot a = 2 \sqrt{2}$ ,  
 $\sec a = \frac{5}{4} \sqrt{2}.$

4. If  $\tan a = .3$ , find  $\sin a$ ,  $\cos a$ ,  $\operatorname{cosec} a$ ,  $\operatorname{vers} a$

$$\text{Ans. } \frac{1}{10}\sqrt{10}, \quad \frac{3}{10}\sqrt{10}, \quad \sqrt{10}, \quad \frac{\sqrt{10}-3}{\sqrt{10}}$$

Prove the truth of the following equations :

5.  $\frac{\sin a}{\tan a} \cdot \frac{\cot a}{\cos a} = \frac{\operatorname{cosec} a}{\sec a}$ .
6.  $(\operatorname{cosec} a - \cot a)^2 = \frac{1 - \cos a}{1 + \cos a}$ .
7.  $\cot a + \frac{\sin a}{1 + \cos a} = \operatorname{cosec} a$ .
8.  $\sec a \{1 + \operatorname{cosec} a (\cos^2 a - \sin a)\} = \cot a$ .
9.  $\sin^2 a \cos^2 \beta - \cos^2 a \sin^2 \beta = \sin^2 a - \sin^2 \beta$ .
10.  $\cos^2 a \cos^2 \beta - \sin^2 a \sin^2 \beta = \cos^2 a - \sin^2 \beta$ .
11.  $\cot a - \sec a \operatorname{cosec} a (1 - 2 \sin^2 a) = \tan a$ .
12.  $\cot^2 a - \tan^2 a = (\cos^2 a - \sin^2 a) \sec^2 a \operatorname{cosec}^2 a$ .
13.  $\frac{\tan a + \tan \beta}{\cot a + \cot \beta} = \tan a \tan \beta$ .
14.  $\tan^2 a \tan^2 \beta - 1 = \frac{\sin^2 a - \cos^2 \beta}{\cos^2 a \cos^2 \beta}$ .
15.  $\frac{1 - \tan^2 a \tan^2 \beta}{\tan^2 a \tan^2 \beta} = \frac{\cos^2 a - \sin^2 \beta}{\sin^2 a \sin^2 \beta}$ .
16.  $\sin^2 a \tan^2 a + \cos^2 a \cot^2 a = \tan^2 a + \cot^2 a - 1$ .
17.  $\sin^2 a \tan a + \cos^2 a \cot a + 2 \sin a \cos a = \sec a \operatorname{cosec} a$ .
18.  $\sec^4 a + \tan^4 a = 1 + 2 \sec^2 a \tan^2 a$ .
19.  $\operatorname{cosec} a (\sec a - 1) - \cot a (1 - \cos a) = \tan a - \sin a$ .
20. Solve the equation  $\sin \theta \cos \theta = \frac{2}{5}$ .

$$\sin^2 \theta \cos^2 \theta = \frac{4}{25};$$

$$\therefore \sin^2 \theta - \sin^4 \theta = \frac{4}{25};$$

$$\therefore \sin^4 \theta - \sin^2 \theta + \frac{1}{4} = \frac{1}{4} - \frac{4}{25} = \frac{9}{100};$$

$$\therefore \sin^2 \theta = \frac{1}{2} \pm \frac{3}{10} = \frac{5 \pm 3}{10} = \frac{1}{5} \text{ or } \frac{4}{5};$$

$$\therefore \sin \theta = \pm \frac{1}{\sqrt{5}} \text{ or } \pm \frac{2}{\sqrt{5}};$$

$$\therefore \theta = \sin^{-1} \left( \pm \frac{1}{\sqrt{5}} \right) \text{ or } \sin^{-1} \left( \pm \frac{2}{\sqrt{5}} \right).$$

Solve the following equations :

21.  $2 \sin \theta = \tan \theta.$  *Ans.*  $\theta = 0$ , or  $60^\circ$ .
22.  $2 \sin^2 \theta + 4 \cos^2 \theta = 3.$  *Ans.*  $\theta = \pm 45^\circ$ .
23.  $\tan \theta + 3 \cot \theta = 4.$  *Ans.*  $\theta = 45^\circ$ , or  $\tan^{-1} 3$ .
24.  $8 \sin \theta = 3 \cos^2 \theta.$  *Ans.*  $\theta = \sin^{-1} \frac{1}{2}$ .
25.  $3 \sin \theta = 2 \cos^2 \theta.$  *Ans.*  $\theta = 30^\circ$ .
26.  $6 \cot^2 \theta - 4 \cos^2 \theta = 1.$  *Ans.*  $\theta = \pm 60^\circ$ .
27.  $\sin \theta + \operatorname{cosec} \theta = 2.$  *Ans.*  $\theta = 90^\circ$ .
28.  $\sec^2 \theta - \frac{5}{2} \sec \theta + 1 = 0.$  *Ans.*  $\theta = 60^\circ$ .
29.  $\sec \theta = 2 \tan \theta.$  *Ans.*  $\theta = 90^\circ$  or  $30^\circ$ .
30. Find the values of  $\cos 585^\circ$ ,  $\tan 600^\circ$ ,  $\operatorname{cosec} 690^\circ$ .
31. Shew that  $\sin^{-1} \frac{4}{5} = \cos^{-1} \frac{3}{5} = \tan^{-1} \frac{4}{3} = \sec^{-1} \frac{5}{3}.$
32. If  $\sin (x + a) = \cos (x - a)$ , find  $x.$  *Ans.*  $x = 45^\circ$ .
33. If  $\sin (a - x) = \cos (a + x)$ , find  $x.$  *Ans.*  $x = -45^\circ$ .
34. If  $\tan \theta + \cot \theta = 2$ , then  $\sin \theta + \cos \theta = \sqrt{2}.$

EXAMPLES. (C).

1. If  $\operatorname{cosec} a = 1.3$ , find  $\sin a$ ,  $\cos a$ ,  $\tan a$ ,  $\cot a$ ,  $\sec a$ .

Prove the truth of the following equations :

2.  $(\sin^2 a - \sin^2 a \sin^2 \beta) (\cos^2 a - \cos^2 a \cos^2 \beta)$   
 $= (\sin^2 \beta - \sin^2 a \sin^2 \beta) (\cos^2 \beta - \cos^2 a \cos^2 \beta).$

$$3. \sec^2 \alpha \tan^2 \alpha (\sec^2 \beta - 1) (\tan^2 \beta + 1) = \sec^2 \beta \tan^2 \beta (\sec^2 \alpha - 1) (\tan^2 \alpha + 1)$$

$$4. \sin \alpha (1 + \tan \alpha) + \cos \alpha (1 + \cot \alpha) = \sec \alpha + \operatorname{cosec} \alpha.$$

$$5. \sin^{-1} \frac{p}{q} = \cot^{-1} \frac{\sqrt{(q^2 - p^2)}}{p}.$$

$$6. \sin^{-1} \frac{p - q}{p + q} = \tan^{-1} \frac{p - q}{2\sqrt{(pq)}}.$$

$$7. \sec^2 \alpha \operatorname{cosec}^2 \alpha - 2 = \tan^2 \alpha + \cot^2 \alpha.$$

$$8. \sin \left( \frac{4n+3}{2} 180^\circ + \alpha \right) = -\cos \alpha.$$

$$9. \cos \{(4n+2) 90^\circ \pm \alpha\} = -\cos \alpha.$$

$$10. \text{ If } \sec \theta \tan \theta = 2\sqrt{3}, \text{ then } \theta = 60^\circ.$$

$$11. \text{ If } \tan \theta \tan \phi = 1, \text{ then } \sec \theta = \operatorname{cosec} \phi.$$

$$12. \text{ If } \frac{\sin \alpha}{\sin \beta} = \sqrt{2}, \frac{\tan \alpha}{\tan \beta} = \sqrt{3}, \text{ then } \alpha = 45^\circ, \beta = 30^\circ.$$

$$13. \text{ If } \frac{\cos \theta}{1 + \sin \theta} + \tan \theta = 2, \theta = 60^\circ.$$

$$14. \text{ If } \cos \phi = n \sin \alpha, \cot \phi = \frac{\sin \alpha}{\tan \beta},$$

$$\text{then } \cos \beta = \frac{n}{\sqrt{(1 + n^2 \cos^2 \alpha)}}.$$

$$15. \text{ If } \cos \alpha + \cos (\beta - \gamma) = 0, \text{ then } \cos \beta + \cos (\alpha - \gamma) = 0, \\ \text{and } \cos \gamma + \cos (\alpha - \beta) = 0.$$

$$16. \text{ If } \frac{\tan \alpha}{\sin \theta} - \frac{\tan \beta}{\tan \theta} = \sqrt{\tan^2 \alpha - \tan^2 \beta}, \text{ then } \cos \theta = \frac{\tan \beta}{\tan \alpha}.$$

$$17. \text{ If } \cos \alpha = \cos \beta \cos \gamma \pm \sin \beta \sin \gamma \cos \alpha, \\ \text{then } \cos \beta = \cos \alpha \cos \gamma \pm \sin \alpha \sin \gamma \cos \beta.$$

$$18. \text{ If } \tan \beta = \sec \alpha \sec \gamma + \tan \alpha \tan \gamma, \\ \text{then } \frac{\tan \alpha + \tan \beta \tan \gamma}{\tan \gamma + \tan \beta \tan \alpha} = \frac{\cos \alpha}{\cos \gamma}.$$

19. Prove that

$$\begin{aligned} & \{(x-y)^2 \sin^2 \theta - y^2\}^2 \cos^2 \theta + \{(x-y)^2 \cos^2 \theta - x^2\}^2 \sin^2 \theta \\ & = \{x^2 \sin^2 \theta + y^2 \cos^2 \theta\}^2. \end{aligned}$$

20. Construct the angle whose secant is  $(\sqrt{5} - 1)$ .

21. Given  $\alpha$  the difference of the lengths of the shadows of a vertical stick when the sun's altitude is  $\alpha, \beta$ , respectively; find the length of the stick.

If the length of the stick is a mean proportional between the lengths of the shadows, shew that  $\alpha, \beta$  are complementary.

22. The elevation of Cader Idris at a point in the valley near Dolgelly is  $\cot^{-1} 6$ ; at Ty Gwyn,  $3\frac{1}{2}$  miles down the valley, it has the same elevation; at a point half-way between its elevation is  $\cot^{-1} 5$ . Its height above the valley is  $\frac{7}{4\sqrt{11}}$  miles.




## SECTION III.

TRIGONOMETRICAL RATIOS OF THE SUMS AND DIFFERENCES OF ANGLES, AND OF MULTIPLE ANGLES. EXPLANATION OF THE DOUBLE SIGN IN TRIGONOMETRICAL FORMULÆ.

27. We now proceed to determine the Trigonometrical Ratios of the sum and difference of any number of angles in terms of the ratios of the simple angles.

To find the Ratios of  $(\alpha + \beta)$  in terms of those of  $\alpha$  and  $\beta$ .



Let  $AOB$  be any angle  $\alpha$ ,  $BOC$  any angle  $\beta$ , then  $AOC$  is  $(\alpha + \beta)$ . In  $OC$  take any point  $P$ , and draw  $PN$ ,  $PQ$  perpendicular to  $OA$ ,  $OB$  respectively. From  $Q$  draw  $QM$ ,  $QR$  perpendicular to  $OA$ ,  $PN$ : then, since  $RPQ$ ,  $RQO$  are each complementary to  $PQR$ , therefore  $RPQ = RQO = QOM = \alpha$ .

And we have,

$$\begin{aligned} \sin(\alpha + \beta) &= \frac{PN}{OP} = \frac{QM}{OP} + \frac{PR}{OP} \\ &= \frac{QM}{OQ} \cdot \frac{OQ}{OP} + \frac{PR}{PQ} \cdot \frac{PQ}{OP}. \end{aligned}$$

Now 
$$\frac{QM}{OQ} = \sin \alpha, \quad \frac{OQ}{OP} = \cos \beta,$$

$$\frac{PR}{PQ} = \cos \alpha, \quad \frac{PQ}{OP} = \sin \beta,$$

therefore  $\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$ ;

also 
$$\cos(\alpha + \beta) = \frac{ON}{OP} = \frac{OM}{OP} - \frac{QR}{OP}$$

$$= \frac{OM}{OQ} \cdot \frac{OQ}{OP} - \frac{QR}{PQ} \cdot \frac{PQ}{OP}$$

$$= \cos \alpha \cos \beta - \sin \alpha \sin \beta;$$

also 
$$\tan(\alpha + \beta) = \frac{\sin(\alpha + \beta)}{\cos(\alpha + \beta)} = \frac{\sin \alpha \cos \beta + \cos \alpha \sin \beta}{\cos \alpha \cos \beta - \sin \alpha \sin \beta};$$

divide numerator and denominator of this fraction by  $\cos \alpha \cos \beta$ , and we have,

$$\tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta};$$

this might have been found directly by geometry as in the case of the sine and cosine.

The sine, cosine, and tangent being found as above, we may find the secant, cosecant, and cotangent.

Put  $\beta + \gamma$  for  $\beta$  in the above form, and we have


$$\begin{aligned} \sin(\alpha + \beta + \gamma) &= \sin \alpha \cos(\beta + \gamma) + \cos \alpha \sin(\beta + \gamma) \\ &= \sin \alpha (\cos \beta \cos \gamma - \sin \beta \sin \gamma) + \cos \alpha (\sin \beta \cos \gamma + \cos \beta \sin \gamma) \\ &= \sin \alpha \cos \beta \cos \gamma + \sin \beta \cos \alpha \cos \gamma + \sin \gamma \cos \alpha \cos \beta - \sin \alpha \sin \beta \sin \gamma. \end{aligned}$$

Similarly the forms for  $\cos(\alpha + \beta + \gamma)$ ,  $\tan(\alpha + \beta + \gamma)$  may be found; and the method may be extended to the sum of any number of angles.

28. To find the Trigonometrical Ratios of  $(\alpha - \beta)$ , in terms of those of  $\alpha$  and  $\beta$ .

Let  $AOB$  be any angle  $\alpha$ ,  $BOC$  any angle  $\beta$ , then  $AOC$  is  $(\alpha - \beta)$ . In  $OC$  take any point  $P$ , and draw  $PN$ ,  $PQ$  perpendicular to  $OA$ ,  $OB$  respectively; from  $Q$  draw  $QM$  perpendicular

to  $OA$ , and from  $P$ ,  $PR$  perpendicular to  $QM$ : then  $PQR = QOA = \alpha$ , for  $PQR$ ,  $QOA$  are each complementary to  $OQM$ .



And we have,

$$\begin{aligned}\sin(\alpha - \beta) &= \frac{PN}{OP} = \frac{QM}{OP} - \frac{QR}{OP} \\ &= \frac{QM}{OQ} \cdot \frac{OQ}{OP} - \frac{QR}{PQ} \cdot \frac{PQ}{OP} \\ &= \sin \alpha \cos \beta - \cos \alpha \sin \beta ;\end{aligned}$$

also,

$$\begin{aligned}\cos(\alpha - \beta) &= \frac{ON}{OP} = \frac{OM}{OP} + \frac{PR}{OP} \\ &= \frac{OM}{OQ} \cdot \frac{OQ}{OP} + \frac{PR}{PQ} \cdot \frac{PQ}{OP} \\ &= \cos \alpha \cos \beta + \sin \alpha \sin \beta ;\end{aligned}$$

also

$$\tan(\alpha - \beta) = \frac{\tan \alpha - \tan \beta}{1 + \tan \alpha \tan \beta} ;$$

and by means of these formulæ the Trigonometrical Ratios of the sums and differences of any number of angles may be calculated.

29. The figures in the above proofs have been drawn so that all the angles are less than  $90^\circ$ . It is left as practice to the student to adapt the proof to any case whatever, and to satisfy himself that the forms proved are true for any angles, a logical

proof of which he will find in more advanced treatises on the subject.

Assuming the form for  $\sin(a + \beta)$  to be true for any values of  $a$  and  $\beta$ , the forms for the other ratios might be deduced as follows :

$$\begin{aligned}\cos(a + \beta) &= \sin\{90 - (a + \beta)\} = \sin(90 + a + \beta) & (\text{Art. 15}) \\ &= \sin(90 + a) \cos \beta + \cos(90 + a) \sin \beta \\ &= \cos a \cos \beta - \sin a \sin \beta, \text{ as before :}\end{aligned}$$

again, putting  $-\beta$  for  $\beta$ ,

$$\begin{aligned}\text{we have} \quad \sin(a - \beta) &= \sin a \cos(-\beta) + \cos a \sin(-\beta) \\ &= \sin a \cos \beta - \cos a \sin \beta ;\end{aligned}$$

$$\begin{aligned}\text{also} \quad \cos(a - \beta) &= \cos a \cos(-\beta) - \sin a \sin(-\beta) \\ &= \cos a \cos \beta + \sin a \sin \beta ;\end{aligned}$$

$$\text{since } \sin(-\beta) = -\sin \beta, \text{ and } \cos(-\beta) = \cos \beta. \quad (\text{Art. 16})$$

30. In the forms for  $a + \beta$ , let  $a = \beta$ ,  
 then we have,

$$\begin{aligned}\sin 2a &= 2 \sin a \cos a, \\ \cos 2a &= \cos^2 a - \sin^2 a, \\ \tan 2a &= \frac{2 \tan a}{1 - \tan^2 a} ;\end{aligned}$$

which give the Ratios of the double angles in terms of those of the single angles.


If we wish to determine  $\sin 2a$ ,  $\cos 2a$  in terms of  $\sin a$ , or of  $\cos a$  only, we have

$$\begin{aligned}\sin 2a &= \pm 2 \sin a \sqrt{1 - \sin^2 a}, \\ &\text{or } = \pm 2 \cos a \sqrt{1 - \cos^2 a}, \\ \cos 2a &= 1 - 2 \sin^2 a, \\ &\text{or } = 2 \cos^2 a - 1.\end{aligned}$$

These forms may be found directly thus :

Let  $AOC = BOC = a$  make  $BCA$  perpendicular to  $OC$ , and  $BN$ ,  $CT$ , to  $OA$ . Then

$$ABN = a, \quad BC = CA, \quad NT = TA, \quad BN = 2CT,$$



and we have

$$\sin 2\alpha = \frac{BN}{OB} = 2 \cdot \frac{BN}{AB} \cdot \frac{BC}{OB} = 2 \cos \alpha \sin \alpha;$$

$$\begin{aligned} \text{also, } \cos 2\alpha &= \sqrt{1 - \sin^2 2\alpha} = \sqrt{1 - 4 \sin^2 \alpha \cos^2 \alpha} \\ &= \sqrt{1 - 4 \sin^2 \alpha + 4 \sin^4 \alpha} = 1 - 2 \sin^2 \alpha, \end{aligned}$$

or directly from the figure, thus,

$$\begin{aligned} \cos 2\alpha &= \frac{ON}{OB} = 1 - \frac{AN}{OB} = 1 - 2 \cdot \frac{AN}{AB} \cdot \frac{BC}{OB} \\ &= 1 - 2 \sin^2 \alpha, \end{aligned}$$

or thus,


$$\begin{aligned} \cos 2\alpha &= \frac{ON}{OB} = \frac{OT}{OB} - \frac{NT}{OB} = \frac{OT}{OC} \cdot \frac{OC}{OB} - \frac{AT}{AC} \cdot \frac{AC}{OA} \\ &= \cos^2 \alpha - \sin^2 \alpha; \end{aligned}$$

$$\begin{aligned} \text{also } \tan 2\alpha &= \frac{BN}{ON} = \frac{2CT}{OT - NT} = \frac{2 \cdot \frac{CT}{OT}}{1 - \frac{AT}{CT} \cdot \frac{CT}{OT}} \\ &= \frac{2 \tan \alpha}{1 - \tan^2 \alpha}. \end{aligned}$$

31. The double sign in the form for  $\sin 2\alpha$  may be explained as follows:

Let  $\angle AOP_1$  be the smallest angle whose sine is equal to  $\sin \alpha$ : then we cannot tell whether the  $\sin \alpha$ , from which we are to determine  $\sin 2\alpha$ , corresponds to the bounding line  $OP_1$  or  $OP_2$ , and therefore we ought not, if our formulæ have their proper gene-

rality, to be able to tell whether  $\sin 2a$  corresponds to  $\angle OQ_1$  or  $\angle OQ_2$  (i.e.  $180^\circ + \angle OQ_2$ ).



Now since  $\angle OQ_2 = 2(180^\circ - \angle OP_1) = 360^\circ - \angle OQ_1$ , therefore the sine of  $\angle OQ_2$  must be equal to the sine of  $\angle OQ_1$ , and be of opposite sign, in accordance with the formulæ proved above. In the same way, if we consider the negative angles bounded by  $OP_1$  and  $OP_2$ , we shall find that our formulæ must correspond to the bounding lines  $OQ_1$ ,  $OQ_2$ , and be correct. So also if we suppose the revolving line to have completed any number of revolutions before reaching the positions  $OP_1$ ,  $OP_2$ .

Precisely similar remarks will apply to  $\sin 2a$  when determined in terms of  $\cos a$ .

The student is also recommended to examine why  $\cos 2a$ ,  $\tan 2a$  have not the double sign.

32. Putting  $\beta = 2a$  in the forms for  $(a + \beta)$ , we have

$$\begin{aligned}\sin 3a &= \sin a \cos 2a + \cos a \sin 2a \\ &= \sin a (1 - 2 \sin^2 a) + 2 \cos^2 a \sin a \\ &= \sin a - 2 \sin^3 a + 2 (1 - \sin^2 a) \sin a \\ &= 3 \sin a - 4 \sin^3 a ;\end{aligned}$$

so also  $\cos 3a = 4 \cos^3 a - 3 \cos a.$

And similarly we may find the Ratios of any multiple angle.

33. If in the forms for  $\sin 2a$  we write  $\frac{a}{2}$  for  $a$ ,

we have, 
$$\sin a = 2 \sin \frac{a}{2} \cos \frac{a}{2},$$

also 
$$1 = \cos^2 \frac{a}{2} + \sin^2 \frac{a}{2}.$$

Hence, 
$$\pm \sqrt{1 + \sin a} = \cos \frac{a}{2} + \sin \frac{a}{2} \dots\dots\dots (A).$$

$$\pm \sqrt{1 - \sin a} = \cos \frac{a}{2} - \sin \frac{a}{2} \dots\dots\dots (B).$$

These formulæ are very useful; the following, which are deduced from them, not so much so.

Adding and subtracting (A) and (B), we have

$$2 \cos \frac{a}{2} = \pm \sqrt{1 + \sin a} \pm \sqrt{1 - \sin a},$$

$$2 \sin \frac{a}{2} = \pm \sqrt{1 + \sin a} \mp \sqrt{1 - \sin a}.$$

Each of these expressions is capable of four values, two and two of opposite signs, according to the signs of the radicals: the reason of which may be explained in a manner similar to that in Art. 31. In choosing practically which of the signs must be used in any particular case, we must return to the formulæ (A) and (B), and inquire, what the signs of  $\cos \frac{a}{2}$ ,  $\sin \frac{a}{2}$  must be, and which is the greatest: we can then determine whether  $\cos \frac{a}{2} + \sin \frac{a}{2}$  is positive or negative, and so also, whether  $\cos \frac{a}{2} - \sin \frac{a}{2}$  is positive or negative. Having thus given their proper signs to the radicals, we can add and subtract the equations (A) and (B), and obtain the proper values of  $\sin \frac{a}{2}$ ,  $\cos \frac{a}{2}$ .

**Ex.** Let  $a = 296^\circ$ . Then  $\cos 148^\circ$  is negative, and numerically greater than  $\sin 148^\circ$ , which is positive.

Hence 
$$\cos 148^\circ + \sin 148^\circ = -\sqrt{1 + \sin 296^\circ},$$

$$\cos 148^\circ - \sin 148^\circ = -\sqrt{1 - \sin 296^\circ},$$

whence the proper signs may be given to  $\cos 148^\circ$ ,  $\sin 148^\circ$ .

34. Writing  $\frac{a}{2}$  for  $a$  in the forms for  $\cos 2a$ , we obtain

$$\cos a = \cos^2 \frac{a}{2} - \sin^2 \frac{a}{2};$$

also 
$$\cos a = 2 \cos^2 \frac{a}{2} - 1,$$

and therefore 
$$\cos \frac{a}{2} = \pm \sqrt{\left(\frac{1 + \cos a}{2}\right)},$$

and 
$$\cos a = 1 - 2 \sin^2 \frac{a}{2},$$

and therefore 
$$\sin \frac{a}{2} = \pm \sqrt{\left(\frac{1 - \cos a}{2}\right)},$$

which give  $\cos \frac{a}{2}$ ,  $\sin \frac{a}{2}$  in terms of  $\cos a$ .

We may also find  $\tan \frac{a}{2}$  in terms of  $\tan a$ , but the formula which results is of very little use.

35. Returning to the formulæ in Arts. 27, 28,

$$\sin (a + \beta) = \sin a \cos \beta + \cos a \sin \beta,$$

$$\sin (a - \beta) = \sin a \cos \beta - \cos a \sin \beta,$$

$$\cos (a + \beta) = \cos a \cos \beta - \sin a \sin \beta,$$

$$\cos (a - \beta) = \cos a \cos \beta + \sin a \sin \beta,$$

we get by addition and subtraction

$$2 \sin a \cos \beta = \sin (a + \beta) + \sin (a - \beta),$$

$$2 \cos a \sin \beta = \sin (a + \beta) - \sin (a - \beta),$$

$$2 \cos a \cos \beta = \cos (a + \beta) + \cos (a - \beta),$$

$$2 \sin a \sin \beta = \cos (a - \beta) - \cos (a + \beta),$$

which forms enable us when we have the product of two sines or cosines to replace them by the sum or difference of two other sines and cosines.



Again, in these forms let  $\alpha = \frac{A+B}{2}$ , and  $\beta = \frac{A-B}{2}$ , then  $\alpha + \beta = A$ ,  $\alpha - \beta = B$ : substituting and transposing, we have

$$\sin A + \sin B = 2 \sin \frac{A+B}{2} \cos \frac{A-B}{2},$$

$$\sin A - \sin B = 2 \cos \frac{A+B}{2} \sin \frac{A-B}{2},$$

$$\cos A + \cos B = 2 \cos \frac{A+B}{2} \cos \frac{A-B}{2},$$

$$\cos B - \cos A = 2 \sin \frac{A+B}{2} \sin \frac{A-B}{2};$$

these forms enable us to pass from the sum or difference of two sines or cosines, to the product of two other sines or cosines.

35. *bis.* The following applications of the formulæ proved in this section are very important.

$$\begin{aligned} (1) \quad & \sin(\alpha + \beta) \sin(\alpha - \beta) \\ &= (\sin \alpha \cos \beta + \cos \alpha \sin \beta) (\sin \alpha \cos \beta - \cos \alpha \sin \beta) \\ &= \sin^2 \alpha \cos^2 \beta - \cos^2 \alpha \sin^2 \beta \\ &= \sin^2 \alpha (1 - \sin^2 \beta) - (1 - \sin^2 \alpha) \sin^2 \beta \\ &= \sin^2 \alpha - \sin^2 \beta. \end{aligned}$$

$$(2) \quad \text{Similarly } \cos(\alpha + \beta) \cos(\alpha - \beta) = \cos^2 \alpha - \sin^2 \beta.$$

$$(3) \quad \tan(45^\circ \pm \theta) = \frac{\tan 45^\circ \pm \tan \theta}{1 \mp \tan 45^\circ \tan \theta} = \frac{1 \pm \tan \theta}{1 \mp \tan \theta}.$$

$$(4) \quad \text{To find } \sin 18^\circ.$$

$$\text{Let } \theta = 18^\circ, \text{ then } 5\theta = 90^\circ,$$

$$\therefore 2\theta = 90^\circ - 3\theta,$$

$$\therefore \sin 2\theta = \cos 3\theta,$$

$$\therefore 2 \sin \theta \cos \theta = 4 \cos^3 \theta - 3 \cos \theta,$$

$$\therefore 2 \sin \theta = 4 \cos^2 \theta - 3$$

$$= 1 - 4 \sin^2 \theta,$$

$$\therefore \sin^2 \theta + \frac{\sin \theta}{2} + \frac{1}{16} = \frac{1}{4} + \frac{1}{16} = \frac{5}{16},$$

$$\therefore \sin \theta = \frac{-1 \pm \sqrt{5}}{4},$$

$$\therefore \sin 18^\circ = \frac{\sqrt{5} - 1}{4},$$

since it must be positive. The other value of  $\sin \theta$  corresponds to the value  $\theta = -54^\circ$ , since in that case also  $\sin 2\theta = \cos 3\theta$ .

(5) Let  $\sin^{-1} x = \theta$ ,  $\sin^{-1} y = \phi$ , then  $\sin \theta = x$ ,

$$\cos \theta = \sqrt{1 - x^2}, \quad \sin \phi = y, \quad \cos \phi = \sqrt{1 - y^2};$$

$$\therefore \sin (\theta + \phi) = x \sqrt{1 - y^2} + y \sqrt{1 - x^2};$$

$$\therefore \sin^{-1} x + \sin^{-1} y = \theta + \phi = \sin^{-1} \{x \sqrt{1 - y^2} + y \sqrt{1 - x^2}\};$$

so also  $\cos^{-1} x + \cos^{-1} y = \cos^{-1} \{xy - \sqrt{1 - x^2} \sqrt{1 - y^2}\},$

and  $\tan^{-1} x + \tan^{-1} y = \tan^{-1} \frac{x + y}{1 - xy}.$

### EXAMPLES. (A).

1. Find the sine of  $15^\circ$ .

$$\begin{aligned} \sin 15^\circ &= \sin (45^\circ - 30^\circ) = \sin 45^\circ \cos 30^\circ - \cos 45^\circ \sin 30^\circ \\ &= \frac{1}{\sqrt{2}} \cdot \frac{\sqrt{3}}{2} - \frac{1}{\sqrt{2}} \cdot \frac{1}{2} = \frac{\sqrt{3} - 1}{2\sqrt{2}}. \end{aligned}$$

$$\text{Ans. } \sin 15^\circ = \frac{\sqrt{3} - 1}{2\sqrt{2}}.$$

Apply the formulæ in Art. 27, to prove the following.

2.  $\cos 15^\circ = \sin 75^\circ = \frac{\sqrt{3} + 1}{2\sqrt{2}}.$

3.  $\sin 15^\circ = \cos 75^\circ = \frac{\sqrt{3} - 1}{2\sqrt{2}}.$

4.  $\cot 15^\circ = \tan 75^\circ = 2 + \sqrt{3}.$

5.  $\tan 15^\circ = \cot 75^\circ = 2 - \sqrt{3}.$

$$6. \text{ If } \sin a = \frac{2}{3}, \sin \beta = \frac{3}{5}, \text{ then } \sin(a + \beta) = \frac{8 + 3\sqrt{5}}{15},$$

$$\cos(a + \beta) = \frac{4\sqrt{5} - 6}{15}.$$

$$7. \text{ If } \sin a = \frac{3}{4}, \cos \beta = \frac{1}{3}, \text{ then } \sin(a + \beta) = \frac{3 + 2\sqrt{14}}{12},$$

$$\cos(a + \beta) = \frac{\sqrt{7} - 6\sqrt{2}}{12}.$$

$$8. \text{ If } \sin a = \frac{5}{13}, \sin \beta = \frac{4}{5}, \text{ then } \sin(45^\circ + a + \beta) = \frac{79\sqrt{2}}{130}.$$

9. Prove the values found in questions 2, 3, 4, 5 by means of Art. 28.

$$10. \text{ If } \sin a = \frac{2}{3}, \sin \beta = \frac{3}{5}, \text{ then } \sin(a - \beta) = \frac{8 - 3\sqrt{5}}{15}.$$

$$11. \text{ If } \sin a = \frac{2}{3}, \text{ then } \sin 2a = \frac{4\sqrt{5}}{9}, \cos 2a = \frac{1}{9}.$$

$$12. \text{ Given } \sin 18^\circ = \frac{\sqrt{5} - 1}{4} \text{ shew that } \sin 36^\circ = \sqrt{\frac{5 - \sqrt{5}}{8}},$$

$$\cos 36^\circ = \frac{\sqrt{5} + 1}{4}, \sin 72^\circ = \frac{\sqrt{10 + 2\sqrt{5}}}{4}.$$

Prove the formulæ

$$13. \quad 1 - \frac{1}{2} \sin 2a \tan a = \cos^2 a.$$

$$14. \quad \sin 4a = 4 \cos 2a \cos a \sin a.$$

$$15. \quad \sin 8a = 8 \cos 4a \cos 2a \cos a \sin a.$$

$$16. \quad \sin(a + 2\beta) = \sin a - 2 \sin a \sin^2 \beta + 2 \cos a \sin \beta \cos \beta.$$

$$17. \quad \sin(a + 2\beta) = \sin a + 2 \sin \beta \cos(a + \beta).$$

$$18. \quad \sin(a - 2\beta) = \sin a - 2 \sin \beta \cos(a - \beta).$$

$$19. \quad \cos(a + 2\beta) = 2 \cos \beta \cos(a + \beta) - \cos a.$$

$$20. \quad \cos a \pm \sin a = \sqrt{1 \pm \sin 2a}.$$

$$21. \quad \cos^4 a - \sin^4 a = \cos 2a.$$

$$22. \cos^2 a + \sin^2 a = 1 - \frac{3}{4} \sin^2 2a = \frac{5 + 3 \cos 4a}{8}.$$

$$23. \cos^2 a - \sin^2 a = \cos 2a \left( \frac{7 + \cos 4a}{8} \right).$$

$$24. \frac{\cos 3a - \sin 3a}{\cos a + \sin a} = 1 - 2 \sin 2a.$$

$$25. \frac{\cos 3a + \sin 3a}{\cos a - \sin a} = 1 + 2 \sin 2a.$$

$$26. \cos 6a = \cos^2 3a - \sin^2 3a = \cos 2a (2 \cos 4a - 1).$$

$$27. \operatorname{cosec}^2 a - \sec^2 a = 4 \cos 2a \operatorname{cosec}^2 2a.$$

$$28. \frac{\operatorname{cosec} 2a - \cot 2a}{\operatorname{cosec} 2a + \cot 2a} = \tan^2 a.$$

$$29. \sin 2a + \cos a = \frac{2 \tan a + \sec a}{1 + \tan^2 a}.$$

$$30. \cot^2 a - \tan^2 a = \frac{8 \cos 2a}{1 - \cos 4a}.$$

$$31. \frac{\cos a - \cos 3a}{\sin 3a - \sin a} = \tan 2a.$$

32. By means of Art. 33, find  $\sin 75^\circ$ ,  $\cos 75^\circ$ , giving the proper signs to the radicals involved.

33. In Art. 33, give the proper signs to the radicals, (1) when  $a$  lies between  $180^\circ$  and  $270^\circ$ ; (2) when  $a$  lies between  $225^\circ$  and  $315^\circ$ ; (3) when  $a$  lies between  $360^\circ$  and  $450^\circ$ .

$$34. \frac{1 + \sin a}{\cos a} = \frac{1 + \tan \frac{a}{2}}{1 - \tan \frac{a}{2}}.$$

$$35. \frac{1 - \sin a}{\cos a} = \tan \left( 45^\circ - \frac{a}{2} \right).$$

$$36. \frac{1 + \sec a}{1 + \operatorname{cosec} a} = \frac{2 \tan a}{\left( 1 + \tan \frac{a}{2} \right)^2}.$$

$$37. \tan \frac{a}{2} + 2 \sin^2 \frac{a}{2} \cot a = \sin a.$$

38. By means of Art. 34, find  $\sin 22\frac{1}{2}^\circ$ ,  $\cos 22\frac{1}{2}^\circ$ .

$$\text{Ans. } \sqrt{\frac{\sqrt{2}-1}{2\sqrt{2}}}, \quad \sqrt{\frac{\sqrt{2}+1}{2\sqrt{2}}}.$$

39. Shew that  $\sin 9^\circ = \sqrt{\frac{4 - \sqrt{10 + 2\sqrt{5}}}{8}}$ ,

$$\cos 9^\circ = \sqrt{\frac{4 + \sqrt{10 + 2\sqrt{5}}}{8}}.$$

40. Find expressions for  $\sin 27^\circ$ ,  $\sin 63^\circ$ .

Apply the formulæ in Art. 35, to prove the following.

41.  $2 \sin a \cos 3a = \sin 4a - \sin 2a.$
42.  $2 \sin a \sin 3a = \cos 2a - \cos 4a.$
43.  $2 \sin 2a \cos a = \sin 3a + \sin a.$
44.  $2 \cos a \cos 2a = \cos 3a + \cos a.$
45.  $\cos 3a \cos 4a = \frac{1}{2} \cos 7a + \frac{1}{2} \cos a.$
46.  $4 \sin a \sin 2a \sin 3a = \sin 2a + \sin 4a - \sin 6a.$
47.  $4 \cos a \cos 2a \cos 3a = 1 + \cos 2a + \cos 4a + \cos 6a.$
48.  $4 \cos a \cos 2a \sin 3a = \sin 2a + \sin 4a + \sin 6a.$
49.  $8 \sin a \sin 2a \sin 3a \sin 4a = 1 - \cos 6a - \cos 8a + \cos 10a.$
50.  $\sin a + \sin 2a = 2 \sin \frac{3a}{2} \cos \frac{a}{2}.$
51.  $\sin 3a - \sin a = 2 \cos 2a \sin a.$
52.  $\cos 2a + \cos 3a = 2 \cos \frac{5a}{2} \cos \frac{a}{2}.$
53.  $\cos 3a - \cos 4a = 2 \sin \frac{7a}{2} \sin \frac{a}{2}.$
54.  $\sin \frac{3a}{2} + \sin 2a = 2 \sin \frac{7a}{4} \cos \frac{a}{4}.$
55.  $\sin 3a - \sin \frac{a}{2} = 2 \cos \frac{7a}{4} \sin \frac{5a}{4}.$
56.  $\sin a + \cos a = 2 \sin 45^\circ \cos (45^\circ - a).$

57.  $\cos a - \sin a = \sqrt{2} \sin (45^\circ - a)$ .
58.  $\cos a + \cos 2a + \cos 3a = 4 \cos \frac{a}{2} \cos a \cos \frac{3a}{2} - 1$ .
59.  $4 \cos a \cos (120 - a) \cos (120 + a) = \cos 3a$ .
60.  $\cos 9a + 3 \cos 7a + 3 \cos 5a + \cos 3a = 8 \cos^3 a \cos 6a$ .
61. If  $\tan \frac{a}{2} = 2 - \sqrt{3}$ , find  $a$ . *Ans.*  $a = 30^\circ$ .
62. If  $\cos 3\theta \cos \theta = \frac{5}{18}$ , then  $\sin \theta = \frac{1}{\sqrt{6}}$ .
63. If  $2 \sin \theta \sin 36^\circ = 1$ , then  $\sin \theta = \sqrt{\frac{5 + \sqrt{5}}{10}}$ .

EXAMPLES. (B).

- Find the Trigonometrical Ratios of  $81^\circ$ ,  $52\frac{1}{2}^\circ$ .
- Find the Trigonometrical Ratios of  $7\frac{1}{2}^\circ$ ,  $105^\circ$ .
- If  $\alpha + \beta + \gamma = 180^\circ$ ,

$$\tan \alpha + \tan \beta + \tan \gamma = \tan \alpha \tan \beta \tan \gamma;$$

we have

$$\tan (\alpha + \beta) = \tan (180^\circ - \gamma),$$

or

$$\frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta} = -\tan \gamma,$$

or

$$\tan \alpha + \tan \beta + \tan \gamma = \tan \alpha \tan \beta \tan \gamma. \quad \text{Q. E. D.}$$

Prove the formulæ,

$$4. \quad \tan (\alpha + 2\beta) = \frac{\tan \alpha + 2 \tan \beta - \tan \alpha \tan^2 \beta}{1 - 2 \tan \alpha \tan \beta - \tan^2 \beta}.$$

$$5. \quad \frac{\tan 3\alpha}{\tan \alpha} = \frac{3 - \tan^2 \alpha}{1 - 3 \tan^2 \alpha}.$$

$$6. \quad \cos 4\alpha = 8 \cos^4 \alpha - 8 \cos^2 \alpha + 1.$$

$$7. \quad \cos (\alpha + \beta) \sin (\alpha - \beta) = \sin \alpha \cos \alpha - \sin \beta \cos \beta.$$

$$8. \quad \sin (\alpha - \beta) \sin \gamma + \sin (\beta - \gamma) \sin \alpha + \sin (\gamma - \alpha) \sin \beta = 0.$$

$$9. \quad \sin (\alpha + \beta) \cos \alpha - \cos (\alpha + \beta) \sin \alpha = \sin \beta.$$

10.  $\cos a + \cos (a + 2\beta) = 2 \cos (a + \beta) \cos \beta.$
11.  $\cos \beta \cos (2a + \beta) = \cos^2 (a + \beta) - \sin^2 a.$
12.  $\frac{\sin (a + \beta) \sin (a - \beta)}{\cos^2 a \cos^2 \beta} = \tan^2 a - \tan^2 \beta.$
13.  $\sin (45^\circ + a) = \frac{1}{\sqrt{2}} (\sin a + \cos a) = \cos (45^\circ - a).$
14.  $\sin (60^\circ + a) = \frac{1}{2} (\sin a + \sqrt{3} \cos a) = \cos (30^\circ - a).$
15.  $\sin (30^\circ + a) + \sin (30^\circ - a) = \cos a.$
16.  $\sec (\theta \pm \phi) = \frac{\sec \theta \sec \phi}{1 \mp \tan \theta \tan \phi}.$
17.  $\frac{1 - \tan (45^\circ - \theta)}{1 + \tan (45^\circ - \theta)} = \tan \theta.$
18.  $\cos^2 (a - \beta) - \sin^2 (a + \beta) = \cos 2a \cos 2\beta.$
19.  $\sin^2 24^\circ - \sin^2 6^\circ = \frac{\sqrt{5} - 1}{8}.$
20.  $(\sin \theta - \sin \phi)^2 + (\cos \theta - \cos \phi)^2 = 2 \operatorname{vers} (\theta - \phi).$
21.  $\cos (30 - a) - \cos (30 + a) = \sin a.$
22.  $\cot a + \tan a = 2 \operatorname{cosec} 2a.$
23.  $\cot a - \tan a = 2 \cot 2a.$
24.  $\frac{\sec a - 1}{\sec a + 1} = \tan^2 \frac{a}{2}.$
25.  $\frac{\sin a + \sin 3a}{\cos a + \cos 3a} = \tan 2a.$
26.  $\tan a + \sec a = \tan \left( 45^\circ + \frac{a}{2} \right).$
27.  $2 (1 - \cot 2a \tan a) = \sec^2 a.$
28.  $\frac{1 + \sin a}{1 + \cos a} = \frac{1}{2} \left( 1 + \tan \frac{a}{2} \right)^2.$
29.  $\frac{1 + \sin a}{1 - \sin a} = \tan^2 \left( 45^\circ + \frac{a}{2} \right).$
30.  $\frac{\sin a + \sin 2a}{\cos a - \cos 2a} = \cot \frac{a}{2}.$

31.  $\tan \theta - \tan \frac{\theta}{2} = \tan \frac{\theta}{2} \sec \theta.$
32.  $\frac{2 \sin 2a - \sin 4a}{2 \sin 2a + \sin 4a} + 1 = \sec^2 a.$
33.  $\cos^2 (\theta + \phi) + \cos^2 (\theta - \phi) - \cos 2\theta \cos 2\phi = 1.$
34.  $\cot^2 a - \tan^2 a = 4 \operatorname{cosec} 2a \cot 2a.$
35.  $\sin a \tan a + \cos a \cot a = (\sin a + \cos a)(2 \operatorname{cosec} 2a - 1).$
36.  $\frac{\cos na - \cos (n+2)a}{\sin (n+2)a - \sin na} = \tan (n+1)a.$
37.  $\sin^4 \theta + 2 \cos a \sin^2 \theta \cos^2 \theta + \cos^4 \theta = 1 - \left( \sin \frac{a}{2} \sin 2\theta \right)^2.$
38.  $\sin 4\theta \tan^4 \theta + 4 \tan^2 \theta + 2 \sin 4\theta \tan^2 \theta - 4 \tan \theta + \sin 4\theta = 0.$
39.  $\frac{\cos 3a - 2 \cos a}{\sin 3a + 2 \sin a} \tan a = \frac{2 \cos 2a - 3}{2 \cos 2a + 3}.$
40.  $\frac{\tan \frac{a}{2} + \cot \frac{a}{2} + 2}{\tan \frac{a}{2} + \cot \frac{a}{2} - 2} = \frac{\sin^2 \left( 45^\circ + \frac{a}{2} \right)}{\sin^2 \left( 45^\circ - \frac{a}{2} \right)}.$
41.  $\cos a + \cos (a + 120^\circ) + \cos (a - 120^\circ) = 0.$
42.  $\cos^2 a + \cos^2 (a + 120^\circ) + \cos^2 (a - 120^\circ) = \frac{3}{2}.$
43.  $\sin 3a + \cos 3a = \sqrt{2} \left\{ \cos^2 \frac{3a - 45^\circ}{2} - \sin^2 \frac{3a - 45^\circ}{2} \right\}.$
44.  $1 + \frac{1}{2}(\tan a - \tan \beta)^2 + \frac{1}{2}\{\tan a - \cot(a + \beta)\}^2$   
 $+ \frac{1}{2}\{\tan \beta - \cot(a + \beta)\}^2 = \tan^2 a + \tan^2 \beta + \cot^2(a + \beta).$

Solve the equations :

45.  $\cos \theta + \cos 7\theta = \cos 4\theta.$  *Ans.*  $\theta = 20^\circ$ , or  $\frac{45^\circ}{2}.$
46.  $\cos \theta - \cos 3\theta = \sin 2\theta.$  *Ans.*  $\theta = 0$ , or  $30^\circ.$
47.  $\cos 4\theta + \cos 2\theta = \cos \theta.$  *Ans.*  $\theta = 20^\circ$ , or  $90^\circ.$
48.  $\sin 5x \cos 3x = \sin 9x \cos 7x.$  *Ans.*  $x = 0$ , or  $7\frac{1}{2}^\circ.$



49.  $\cos 2x + \sin x = 1$ . *Ans.*  $x = 0$ , or  $30^\circ$ .
50.  $4 \cot 2\theta = \cot^2 \theta - \tan^2 \theta$ . *Ans.*  $\theta = 0$ , or  $45^\circ$ .
51.  $\operatorname{cosec} \theta = 2 \cos 2\theta + 4 \cos^2 \theta$ . *Ans.*  $\theta = 10^\circ$ .
52.  $\sin x + \sqrt{3} \cos x = \sqrt{2}$ . *Ans.*  $x = 15^\circ$ , or  $75^\circ$ .
53.  $\frac{1}{\sin x} + \frac{\sqrt{3}}{\cos x} = 4$ . *Ans.*  $x = 30^\circ$ .
54.  $3(\sin^4 \theta - \cos^4 \theta) + 4 \cos^4 \theta = \cos^2 2\theta$ . *Ans.*  $\theta = 0$ .
55.  $\sin 9x + \sin 5x + 2 \sin^2 x = 1$ . *Ans.*  $x = 45^\circ$ , or  $2\frac{1}{2}^\circ$ .
56.  $\cos 4\theta + \cos 2\theta + \cos \theta = 0$ . *Ans.*  $\theta = 90^\circ$ , or  $40^\circ$ .
57.  $\cos ax \cos bx = \cos (a+c)x \cos (b+c)x$ .  
*Ans.*  $x = 0$ , or  $\frac{180^\circ}{c}$ , or  $\frac{180^\circ}{a+b+c}$ .
58. If  $\sin x = \sin a \sin (x+y)$ , then  $\tan x = \frac{\sin a \sin y}{1 - \sin a \cos y}$ .
59. If  $\tan \beta = \frac{2 \sin a \sin \gamma}{\sin (a+\gamma)}$ , then  $\cot a, \cot \beta, \cot \gamma$  are in A.P.
60. If  $\sin \beta = m \sin (2a + \beta)$ , then  $\tan (a + \beta) = \frac{1+m}{1-m} \tan a$ .
61. If  $\cot \theta = \frac{a}{b}$ , then  $a \sin \theta - b \cos \theta = 0$ ;  
and  $\frac{a \cos \theta - b \sin \theta}{a \cos \theta + b \sin \theta} = \frac{a^2 - b^2}{a^2 + b^2}$ .
62. If  $\sin x = \frac{\sin a}{\sin \gamma}$ ,  $\sin (90^\circ - x) = \frac{\sin \beta}{\sin \gamma}$ ,  
then  $\cos^2 a + \cos^2 \beta + \sin^2 \gamma = 2$ .
63. If  $a \cos \phi = b \cos \theta$ , then  $\cot \frac{\phi + \theta}{2} \cot \frac{\phi - \theta}{2} = \frac{a+b}{a-b}$ .
64. If  $\tan \beta = \sec a \sec \gamma + \tan a \tan \gamma$ , then  $\cos 2\beta$  is negative.
65. If  $\tan (a + \theta) \cot a = \tan (\beta + \phi) \cot \beta$ ,  
then  $\sin (2a + \theta) \operatorname{cosec} \theta = \sin (2\beta + \phi) \operatorname{cosec} \phi$ .
66. If  $\cos \theta = \tan \beta \cot \gamma$ ,  $\cot \theta = \sin \beta \cot a$ ,  
then  $\cos \gamma = \cos a \cos \beta$ .

67. If  $a \sin \alpha + x \cos \theta = b + x \tan \beta \sin \theta$ ,

and  $a \cos \alpha - x \sin \theta = x \tan \beta \cos \theta$ ,

$$\text{then } \tan \theta = \frac{a \cos (\beta - \alpha) - b \sin \beta}{a \sin (\beta - \alpha) + b \cos \beta}.$$

68. If  $\tan (\phi + \alpha)$ ,  $\tan \phi$ ,  $\tan (\phi + \beta)$  are in A.P. so also are  $\cot \alpha$ ,  $\tan \phi$ ,  $\cot \beta$ .

Prove the formulæ,

69.  $\sin^{-1} \frac{3}{5} + \sin^{-1} \frac{4}{5} = 90^\circ$ .

70.  $\cot^{-1} a - \cot^{-1} b = \cot^{-1} \frac{1 + ab}{b - a}$ .

71.  $\text{vers}^{-1} \frac{4a - x}{a} = 180^\circ - \text{vers}^{-1} \frac{x - 2a}{a}$ .

72.  $\tan^{-1} \frac{3}{5} + \cot^{-1} \frac{7}{3} = \cot^{-1} \frac{13}{18}$ .

73.  $\tan^{-1} \frac{2}{3} + \tan^{-1} \frac{1}{2} + \tan^{-1} \frac{1}{3} = \tan^{-1} 5$ .

74.  $45^\circ - 3 \tan^{-1} \frac{1}{4} - \tan^{-1} \frac{1}{20} = \tan^{-1} \frac{7}{467}$ .

75.  $\sin^{-1} \frac{4}{5} + \sin^{-1} \frac{5}{13} + \sin^{-1} \frac{16}{65} = 90^\circ$ .

76.  $\tan^{-1} \frac{1}{x-1} - \tan^{-1} \frac{1}{x} = \tan^{-1} \frac{1}{x^2 - x + 1}$ .

77.  $\tan (\sin^{-1} m + \cos^{-1} n) = \frac{mn + \sqrt{(1-m^2)(1-n^2)}}{n\sqrt{1-m^2} - m\sqrt{1-n^2}}$ .

78. If  $\tan^{-1} \frac{1}{x} - \tan^{-1} \frac{1}{x+2} = 30^\circ$ , then  $x = -1 \pm \sqrt{2\sqrt{3}}$ .

79. If  $\tan^{-1} \frac{x-1}{x-2} + \tan^{-1} \frac{x+1}{x+2} = 45^\circ$ , then  $x = \pm \frac{1}{\sqrt{2}}$ .

## EXAMPLES. (C).

Prove the formulæ,

1.  $\frac{\cos a + \sin a}{\cos a - \sin a} = \tan 2a + \sec 2a.$
2.  $\frac{\sin a + \sin (a + \beta) + \sin (a + 2\beta)}{\cos a + \cos (a + \beta) + \cos (a + 2\beta)} = \tan (a + \beta).$
3.  $16 \operatorname{cosec}^2 4a - 4 \sec^2 2a = \sec^2 a \operatorname{cosec}^2 a.$
4.  $\frac{\sin a + \sin 2a}{1 + \cos a + \cos 2a} = \tan a.$
5.  $\frac{\cos (a + 45^\circ)}{\cos (a - 45^\circ)} = \sec 2a - \tan 2a.$
6.  $4 \sin \frac{\theta}{3} \sin \frac{180 - \theta}{3} \sin \frac{180 + \theta}{3} = \sin \theta.$
7.  $\frac{\tan a + \sec a}{\cot a + \operatorname{cosec} a} = \tan \left( 45^\circ + \frac{a}{2} \right) \tan \frac{a}{2}.$
8.  $\frac{\sin (a + \beta + \gamma)}{\cos a \cos \beta \cos \gamma} = \tan a + \tan \beta + \tan \gamma - \tan a \tan \beta \tan \gamma.$
9.  $\tan 4a = \frac{4 \tan a - 4 \tan^3 a}{1 - 6 \tan^2 a + \tan^4 a}.$
10.  $\tan (30^\circ + a) \tan (30^\circ - a) = \frac{2 \cos 2a - 1}{2 \cos 2a + 1}.$
11.  $\frac{\sin^2 2a - 4 \sin^2 a}{\sin^2 2a + 4 \sin^2 a - 4} = \tan^4 a.$
12.  $2 \operatorname{vers} \frac{180^\circ + a}{2} \operatorname{vers} \frac{180^\circ - a}{2} = \operatorname{vers} (180^\circ - a).$
13.  $\tan a \tan (a - \beta) (\cot a + \tan \beta) = \tan a - \tan \beta.$
14.  $\frac{1 + \operatorname{cosec}^2 a \tan^2 \theta}{1 + \operatorname{cosec}^2 \beta \tan^2 \theta} = \frac{1 + \cot^2 a \sin^2 \theta}{1 + \cot^2 \beta \sin^2 \theta}.$
15.  $\sin^2 \beta + \sin^2 (a - \beta) + 2 \sin \beta \sin (a - \beta) \cos a = \sin^2 a.$
16.  $\frac{\sin^2 a - \sin^2 \beta}{\sin a \cos a - \sin \beta \cos \beta} = \tan (a + \beta).$

$$17. \quad 1 + \tan \alpha \tan \frac{\alpha}{2} = \sec \alpha.$$

$$18. \quad \left( \sin \frac{\alpha + \beta}{2} + \cos \frac{\alpha + \beta}{2} \right) \left( \sin \frac{\alpha - \beta}{2} + \cos \frac{\alpha - \beta}{2} \right) \\ = \sin \alpha + \cos \beta.$$

$$19. \quad \cos^2 (\alpha - \theta) + \cos^2 \theta - 2 \cos (\alpha - \theta) \cos \theta \cos \alpha = \sin^2 \alpha.$$

$$20. \quad \tan (45^\circ + \theta) - \tan (45^\circ - \theta) = \frac{2(\sec 2\theta - \cos 2\theta)}{\sin 2\theta}.$$

$$21. \quad \tan \frac{\theta}{3} \cot \frac{1}{3} (90^\circ - \theta) \cot \frac{1}{3} (90^\circ + \theta) = \tan \theta.$$

$$22. \quad \frac{\tan \phi - \cos \theta}{\tan \phi + \cos \theta} = \frac{1 - \tan^2 \frac{\theta}{2} \tan (45^\circ + \phi)}{\tan^2 \frac{\theta}{2} - \tan (45^\circ + \phi)}.$$

$$23. \quad (\cos \theta + \sin \theta)^2 = \frac{3}{2} (\cos \theta + \sin \theta) - \frac{1}{2} (\cos 3\theta - \sin 3\theta).$$

$$24. \quad 64 \sin^2 \theta \cos^2 \theta = 3 \sin \theta + 3 \sin 3\theta - \sin 5\theta - \sin 7\theta.$$

$$25. \quad \frac{\cos (45^\circ + \alpha) - \cos (45^\circ + 3\alpha)}{\cos (45^\circ - \alpha) + \cos (45^\circ - 3\alpha)} = \tan \alpha.$$

$$26. \quad \cos 3\theta \sin^2 \theta + \sin 3\theta \cos^2 \theta = \frac{3}{4} \sin 4\theta.$$

$$27. \quad \sin 3\theta \sin^2 \theta + \cos 3\theta \cos^2 \theta = \cos^3 2\theta.$$

$$28. \quad \frac{(\sec \theta \sec \phi + \tan \theta \tan \phi)^2 - (\tan \theta \sec \phi + \sec \theta \tan \phi)^2}{2(1 + \tan^2 \theta \tan^2 \phi) - \sec^2 \theta \sec^2 \phi} \\ = \frac{\sec 2\theta \sec 2\phi}{\sec^2 \theta \sec^2 \phi}.$$

$$29. \quad (\sec \theta + \operatorname{cosec} \theta + \sec \theta \operatorname{cosec} \theta) \left( 1 - \tan^2 \frac{\theta}{2} \right) \left( 1 - \tan^2 \frac{\theta}{4} \right) \\ = \left( \sec \frac{\theta}{2} + \operatorname{cosec} \frac{\theta}{2} \right) \sec^2 \frac{\theta}{4}.$$

$$30. \quad \tan \theta + \tan (\theta + 60^\circ) + \tan (\theta - 60^\circ) = 3 \tan 3\theta.$$

$$31. \quad (2 + \sin 2\theta) \sin (\theta - 45^\circ) + (2 - \sin 2\theta) \sin (\theta + 45^\circ) = 2\sqrt{2} \sin^2 \theta$$

$$32. \quad \tan^{-1} \frac{1}{\sqrt{2}} + \sin^{-1} \frac{1}{\sqrt{2}} = \tan^{-1} \frac{\sqrt{2} + 1}{\sqrt{2} - 1}.$$

Solve the equations,

$$33. \quad 5 \tan x = \tan 3x. \quad \text{Ans.} \quad \tan x = \pm \sqrt{\frac{1}{7}}, \text{ or } 0.$$

$$34. \quad 4 \sin \theta \sin (\theta - \alpha) = 2 \cos \alpha - 1. \quad \text{Ans.} \quad \theta = 30 + \frac{\alpha}{2}.$$

$$35. \quad 90^\circ + \sin^{-1} x = \tan^{-1} x. \quad \text{Ans.} \quad x = \sqrt{\frac{5-1}{2}}.$$

$$36. \quad \tan^{-1} 2x + \tan^{-1} 3x = 45^\circ. \quad \text{Ans.} \quad x = -1, \text{ or } \frac{1}{6}.$$

$$37. \quad \tan^{-1} \frac{\alpha+x}{a} + \tan^{-1} \frac{\alpha-x}{a} = 30^\circ. \quad \text{Ans.} \quad x = a \sqrt{2\sqrt{3}}.$$

$$38. \quad \tan^{-1} 3x + 2 \tan^{-1} 7x = 45^\circ. \quad \text{Ans.} \quad x = \frac{1}{21}.$$

$$39. \quad \tan^{-1} ax + \tan^{-1} \frac{x}{a} = 45^\circ. \\ \text{Ans.} \quad x = \frac{-(a^2+1) \pm \sqrt{a^4+6a^2+1}}{2a}.$$

$$40. \quad \tan^{-1} (x+1) \sqrt{2} - \tan^{-1} \frac{x-1}{\sqrt{2}} = \cot^{-1} 4 \sqrt{2}. \\ \text{Ans.} \quad x = 6, \text{ or } -2.$$

$$41. \quad \cos^{-1} x + \cos^{-1} (1-x) = \cos^{-1} (-x). \quad \text{Ans.} \quad x = 0, \text{ or } \frac{1}{2}.$$

$$42. \quad 3 \tan^{-1} t = \tan^{-1} 5t. \quad \text{Ans.} \quad t = 0, \text{ or } \pm \frac{1}{\sqrt{7}}.$$

$$43. \quad \cos^{-1} x + \cos^{-1} (1-x) = \cos^{-1} \sqrt{(x-x^2)}. \\ \text{Ans.} \quad x = 0, \text{ or } \frac{1}{2}, \text{ or } 1.$$

$$44. \quad \tan^{-1} \frac{1}{x-1} - \tan^{-1} \frac{1}{x+1} = \tan^{-1} a. \\ \text{Ans.} \quad x = \sqrt{\frac{2}{a}}.$$

$$45. \quad \tan (\alpha+x) \tan (\alpha-x) = \frac{1-2 \cos 2\alpha}{1+2 \cos 2\alpha}. \quad \text{Ans.} \quad x = 30^\circ.$$

$$46. \quad \cos^{-1} \frac{1-x^2}{1+x^2} + \tan^{-1} \frac{2x}{1-x^2} = 240^\circ. \\ \text{Ans.} \quad x = \sqrt{3}, \text{ or } -\frac{1}{\sqrt{3}}.$$

47.  $\{\cot(m+n)a + 1\} \cos 2ma = \{\cot(m+n)a - 1\} \cos 2na$ .  
*Ans.*  $a = \frac{45^\circ}{m-n}$ .
48.  $\sec(180-a) \cos(90+a) \cos(180-a)$ .  
 $= \tan(90-a) \sin(270+a)$ .  
*Ans.*  $a = \pm 45^\circ$ .
49.  $\cos na + \cos(n-2)a = \cos a$ . *Ans.*  $a = 90^\circ$ , or  $\frac{60^\circ}{n-1}$ .
50.  $\operatorname{cosec}^2 \frac{\theta}{2} - \sec^2 \frac{\theta}{2} = 2\sqrt{3} \operatorname{cosec}^2 \theta$ . *Ans.*  $\theta = 30^\circ$ .
51.  $\sin \theta - \cos \theta = 2\sqrt{2} \cos \theta \sin \theta$ .  
*Ans.*  $\theta = 15^\circ$ , or  $75^\circ$ , or  $135^\circ$ .
52.  $4 \cos \theta \cos 2\theta = 1$ . *Ans.*  $\theta = 120^\circ$ , or  $36^\circ$ , or  $108^\circ$ .
53.  $\sin^6 x + \cos^4 x = \frac{3}{8}$ . *Ans.*  $x = 45^\circ$ , or  $\sin^{-1} \sqrt{\frac{\sqrt{(29)}-3}{4}}$ .
54.  $\tan(a+\theta) \tan(a-\theta) = \tan^2 2a - \tan^2 \theta$ .  
*Ans.*  $\theta = \pm a$ , or  $\frac{1}{2} \cos^{-1} \left( \frac{-\cos 2a}{1 \pm \cos 2a} \right)$ .
55.  $\sec^2 \frac{\theta}{2} \sec \theta \left( \cot^2 \frac{\theta}{2} - \cot^2 \frac{3\theta}{2} \right) = 8 \left( 1 - \cot^2 \frac{3\theta}{2} \right)$ .  
*Ans.*  $\theta = 60^\circ$ .
56.  $\tan \theta + \cot \theta = 2 \operatorname{cosec} a (\sqrt{1 + \sin a} + \sqrt{1 - \sin a})$ .  
*Ans.*  $\theta = \frac{a}{4}$ , or  $45^\circ - \frac{a}{4}$ .
57.  $\sin(3x+a) + \cos(3x-a) = \cos(45^\circ+x)$ .  
*Ans.*  $x = 45^\circ$ , or  $\frac{1}{2} \sin^{-1} \frac{1}{4} \{\sec(45^\circ-a) - 2\}$ .
58.  $\frac{1}{\sqrt{1-\sin \theta}-1} + \frac{1}{\sqrt{1+\sin \theta}-1} = \frac{1}{\sin \theta}$ .  
*Ans.*  $\theta = 0, 180^\circ$ , or  $60^\circ$ .

59.  $\sin(\tan^{-1} x) + \tan(\sin^{-1} x) = mx.$

*Ans.*  $x = \frac{1}{m}(m^4 - m^2 - 2 \pm 2\sqrt{2m^2 + 1})^{\frac{1}{2}}.$

60. If  $\tan \beta + \cot \beta = 2 \sec 2\alpha$ , then  $\alpha + \beta = 45^\circ$ .

61. If  $\tan(\theta + \alpha) \tan(\theta - \beta) = \tan^2 \theta$ ,  
then  $\tan 2\theta = \frac{2 \sin \alpha \sin \beta}{\sin(\alpha - \beta)}.$

62. If  $\tan 2\alpha = 2 \tan \beta$ , and  $\tan \gamma = \tan^3 \alpha$ ,  
shew that  $\beta - \gamma = \alpha$ .

63. Shew that  $\tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{5} + \tan^{-1} \frac{1}{7} + \tan^{-1} \frac{1}{8} = 45^\circ$ .

64. Find  $\tan(\alpha + \beta + \gamma)$  in terms of  $\tan \alpha$ ,  $\tan \beta$ ,  $\tan \gamma$ ; and  
thence shew that if  $\alpha + \beta + \gamma = 180^\circ$ ,

$$\tan \alpha + \tan \beta + \tan \gamma = \tan \alpha \tan \beta \tan \gamma.$$

65. If  $\tan(\alpha + \theta) = n \tan(\alpha - \theta)$ , then  $\sin 2\theta = \frac{n-1}{n+1} \sin 2\alpha$ .

66. If  $\sin 2\theta = \tan \theta$ , then  $\csc \theta = \sqrt{2 - \sqrt{2}}$ .

67. If  $\sin \theta + \sin \phi = m$ ,  $\cos \theta + \cos \phi = n$ ,  
then  $m \sin \theta + n \cos \theta = \frac{m^2 + n^2}{2} = m \sin \phi + n \cos \phi$ .

68. If  $\tan^2 \theta = \frac{b}{a}$ , then  $\frac{a}{\cos \theta} + \frac{b}{\sin \theta} = (a^{\frac{2}{3}} + b^{\frac{2}{3}})^{\frac{3}{2}}$ .

69. Find  $\cos(\alpha + \beta + \gamma)$ , and if  $\alpha + \beta + \gamma = 180^\circ$ , shew that  
 $\cot \alpha + \cot \beta + \cot \gamma = \cot \alpha \cot \beta \cot \gamma + \operatorname{cosec} \alpha \operatorname{cosec} \beta \operatorname{cosec} \gamma$ .

If  $\alpha + \beta + \gamma = 180^\circ$ , then

70.  $\sin \alpha + \sin \beta + \sin \gamma = 4 \cos \frac{\alpha}{2} \cos \frac{\beta}{2} \cos \frac{\gamma}{2}.$

71.  $\cos \alpha + \cos \beta + \cos \gamma = 4 \sin \frac{\alpha}{2} \sin \frac{\beta}{2} \sin \frac{\gamma}{2} + 1.$

72.  $\cot \frac{\alpha}{2} + \cot \frac{\beta}{2} + \cot \frac{\gamma}{2} = \cot \frac{\alpha}{2} \cot \frac{\beta}{2} \cot \frac{\gamma}{2}.$

73.  $\cos^2 \frac{\alpha}{2} + \cos^2 \frac{\beta}{2} + \cos^2 \frac{\gamma}{2} = 2 \left( 1 + \sin \frac{\alpha}{2} \sin \frac{\beta}{2} \sin \frac{\gamma}{2} \right).$

$$74. \quad \frac{\tan a}{\tan \beta} + \frac{\tan \beta}{\tan a} + \frac{\tan a}{\tan \gamma} + \frac{\tan \gamma}{\tan a} + \frac{\tan \beta}{\tan \gamma} + \frac{\tan \gamma}{\tan \beta} \\ = \sec a \sec \beta \sec \gamma - 2.$$

$$75. \quad \tan \left( 45^\circ - \frac{a}{4} \right) \tan \left( 45^\circ - \frac{\beta}{4} \right) \tan \left( 45^\circ - \frac{\gamma}{4} \right) \\ = \frac{2 \cos \frac{a}{2} \cos \frac{\beta}{2} \cos \frac{\gamma}{2}}{\left( \cos \frac{a}{2} + \cos \frac{\beta}{2} + \cos \frac{\gamma}{2} \right)^2}.$$

$$76. \quad \sin 2a + \sin 2\beta + \sin 2\gamma = 4 \sin a \sin \beta \sin \gamma.$$

$$77. \quad \cos 2a + \cos 2\beta + \cos 2\gamma + 4 \cos a \cos \beta \cos \gamma + 1 = 0.$$

$$78. \quad \cos^2 a + \cos^2 \beta + \cos^2 \gamma + 2 \cos a \cos \beta \cos \gamma = 1.$$

$$79. \quad \cos \frac{a}{2} + \cos \frac{\beta}{2} + \cos \frac{\gamma}{2} = 4 \cos \frac{a+\beta}{4} \cos \frac{\beta+\gamma}{4} \cos \frac{a+\gamma}{4}.$$

$$80. \quad \sin 3a + \sin 3\beta + \sin 3\gamma = -4 \cos \frac{3a}{2} \cos \frac{3\beta}{2} \cos \frac{3\gamma}{2}.$$

$$81. \quad \sin^3 a + \sin^3 \beta + \sin^3 \gamma = 3 \cos \frac{a}{2} \cos \frac{\beta}{2} \cos \frac{\gamma}{2} + \cos \frac{3a}{2} \cos \frac{3\beta}{2} \cos \frac{3\gamma}{2}.$$

$$82. \quad \text{If } \cos^{-1} \frac{x}{a} = 2 \sin^{-1} \frac{y}{a}, \text{ then } ax + 2y^2 = a^2.$$

$$83. \quad \text{If } \cot(a-\gamma) - \cot(a-\theta) = \cot(a+\gamma) - \cot(a+\theta), \\ \text{then } \sin(a-\gamma) \sin(\theta-a) = \sin(a+\gamma) \sin(\theta+a).$$

$$84. \quad \text{If } \tan^2 x = \tan(a-x) \tan(a+x), \text{ then } \sin 2x = \sqrt{2} \cdot \sin a.$$

$$85. \quad \text{If } \sin(a-\beta) = \frac{1}{m} \sin(a+\beta), \\ \text{then } \cot(a+\beta) = \frac{1}{2} \left( \frac{m-1}{m} \cot \beta - \frac{m+1}{m} \tan \beta \right).$$

$$86. \quad \text{If } \cos \theta = \frac{\cos a + \cos \beta}{1 + \cos a \cos \beta}, \text{ then } \tan \frac{\theta}{2} = \tan \frac{a}{2} \tan \frac{\beta}{2}.$$

$$87. \quad \text{If } \cot \theta = m \cot(a-\theta), \\ \text{then } \theta = \frac{1}{2} \left\{ a - \sin^{-1} \left( \frac{m-1}{m+1} \sin a \right) \right\}.$$



88. If  $\tan \frac{a}{2} = \sqrt{\frac{1+e}{1-e}} \tan \frac{\beta}{2}$ , then  $\cos \beta = \frac{e + \cos a}{1 + e \cos a}$ .

89. If  $\frac{\sin(\theta + a)}{\cos(\theta - a)} = \cos 2\beta$ , then  $\tan(45^\circ - \theta) = \tan^2 \beta \cot(45^\circ - a)$ .

90. If  $2 \cot \beta = \cot a + \cot \gamma$ , then  $\cot(\beta - a) + \cot(\beta - \gamma) = 2 \cot \beta$ .

91. If  $\cot^2 \theta - \tan^2 \theta = 2 \sqrt{3} \cdot \sec 2\theta$ , then  $\sin 2\theta = \pm (\sqrt{3} - 1)$ .

92. If  $\cos \theta \cot \phi = \cos \theta_1 \cot \phi_1$ ,

$$\text{then } \cot \frac{\theta + \theta_1}{2} = \frac{\sin(\phi + \phi_1)}{\sin(\phi - \phi_1)} \tan \frac{\theta_1 - \theta}{2}.$$

93. If  $a + \beta + \gamma = 180^\circ$ , then

$$\cot ma \cot m\beta + \cot ma \cot m\gamma + \cot m\beta \cot m\gamma = 1,$$

where  $m$  is any integer.

94. If  $\cos \theta + \cos(\theta - \beta) = \cos(\theta - a) + \cos(\theta + a - \beta)$ ,

then each member of the equation is equal to  $\cos \theta + \cos(\theta - a)$ , or 0

95. If  $\tan(45^\circ + \theta) \tan(45^\circ + \phi) \tan(45^\circ + \psi) = 1$ ,

$$\text{then } \sin 2\theta \sin 2\phi \sin 2\psi + \sin 2\theta + \sin 2\phi + \sin 2\psi = 0.$$

96. If  $a + \beta + \gamma = 60^\circ$ , then

$$1 + 2 \cos(a + \beta - \gamma) + 2 \cos(a + \gamma - \beta) + 2 \cos(\beta + \gamma - a) = 8 \cos a \cos \beta \cos \gamma.$$

97. If  $\theta + \phi = 240^\circ$ , and  $\text{vers } \theta = 4 \text{ vers } \phi$ ,

$$\text{then } \tan \frac{\theta}{2} = -\frac{\sqrt{3}}{2}, \text{ or } \infty, \text{ and } \tan \frac{\phi}{2} = -\frac{\sqrt{3}}{5}, \text{ or } \frac{1}{\sqrt{3}}.$$

98.  $\sin(a - \beta) + \sin(\beta - \gamma) + \sin(\gamma - a)$

$$+ 4 \sin \frac{a - \beta}{2} \sin \frac{\beta - \gamma}{2} \sin \frac{\gamma - a}{2} = 0.$$

99.  $\sin(a + \beta) \sin(\beta + \gamma) - \sin a \sin \gamma = \sin(a + \beta + \gamma) \sin \beta$ .

100.  $4 \cos(a + \beta + \gamma) \cos(a + \beta - \gamma) \cos(a - \beta + \gamma) \cos(a - \beta - \gamma)$   
 $= \cos^2 2a + \cos^2 2\beta + \cos^2 2\gamma + 2 \cos 2a \cos 2\beta \cos 2\gamma - 1.$

101.  $\cos 2(a + \beta + \gamma) \sin 3(a + \beta) + \cos(\gamma - a) \sin(\gamma - \beta)$   
 $= \sin 2(a + \beta + \gamma) \cos 3(a + \beta) - \sin(\gamma - a) \cos(\gamma - \beta).$

$$102. \quad \sin \left( \frac{\theta + \phi}{2} - \psi \right) \cos \frac{\theta + \psi}{2} - \sin \left( \frac{\theta + \psi}{2} - \phi \right) \cos \frac{\theta + \phi}{2} \\ = \sin \frac{\phi - \psi}{2} (\cos \theta + \cos \phi + \cos \psi).$$

$$103. \quad \tan^{-1} \sqrt{\cos 60^\circ} + n \cos^{-1} (\tan^2 30^\circ) = \left( n + \frac{1}{2} \right) \cos^{-1} \frac{1}{3}.$$

$$104. \quad (2^n \operatorname{cosec} 2^n a)^2 + (2^n \sec 2^n a)^2 = (2^{n+1} \operatorname{cosec} 2^{n+1} a)^2.$$

$$105. \quad 2 \tan^{-1} \left\{ \sqrt{\frac{a-b}{a+b}} \cdot \tan \frac{x}{2} \right\} = \cos^{-1} \left\{ \frac{b+a \cos x}{a+b \cos x} \right\}.$$

$$106. \quad \cot \{ \theta + \tan^{-1} (\tan^2 \theta) \} = 2 \cot 2\theta.$$

$$107. \quad 3 \tan^{-1} \frac{1}{7} + \tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{26} - 45^\circ = \tan^{-1} \frac{1}{2057}.$$

$$108. \quad \tan^{-1} \frac{1}{4} + \tan^{-1} \frac{2}{9} + \tan^{-1} \frac{1}{5} + \tan^{-1} \frac{1}{8} = 45^\circ.$$

$$109. \quad \cos^2 \frac{45^\circ}{2} + \cos^2 \frac{135^\circ}{2} + \cos^2 \frac{225^\circ}{2} + \cos^2 \frac{315^\circ}{2} = \frac{17}{16}.$$

$$110. \quad \text{If } x = r \sin \frac{\theta - \alpha}{2}, y = r \sin \frac{\theta + \alpha}{2}, \\ \text{then } x^2 - 2xy \cos \alpha + y^2 = r^2 \sin^2 \alpha.$$

$$111. \quad \text{If } \cos \theta = \cot \beta \cot \gamma, \cos \phi = \cot \alpha \cot \gamma, \cos \psi = \cot \alpha \cot \beta, \\ \text{and } \theta + \phi + \psi = 180^\circ, \text{ then } \cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1.$$

Solve the equations :

$$112. \quad \frac{x}{a} \cos \frac{\phi + \theta}{2} + \frac{y}{b} \sin \frac{\phi + \theta}{2} = \cos \frac{\phi - \theta}{2}, \quad \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1.$$

$$\text{Ans.} \quad \begin{array}{l} x = a \cos \theta, \quad \text{or} \quad x = a \cos \phi, \\ y = b \sin \theta, \quad \text{or} \quad y = b \sin \phi. \end{array}$$

$$113. \quad \begin{array}{l} \sin^{-1} x + \sin^{-1} y = 120^\circ, \\ \cos^{-1} x - \cos^{-1} y = 60^\circ. \end{array} \quad \text{Ans.} \quad \begin{array}{l} x = \pm 1, \\ y = \pm \frac{1}{2}. \end{array}$$

$$114. \quad \text{If } \alpha, \beta, \gamma \text{ are the angles of a triangle, and}$$

$$\cos \beta = \frac{\sin \alpha}{2 \sin \gamma},$$

then the triangle is isosceles.

115. An object 6 feet high, placed on the top of a tower, subtends an angle  $\tan^{-1}.015$  at a place whose horizontal distance from the foot of the tower is 100 feet: find the height of the tower. *Ans.* 170 feet nearly.

116. A person observes the angle subtended by a tower and its spire to be the same; he knows the height  $h$  of the tower, and his distance  $a$  from it. The height of the spire is

$$\frac{a^2 + h^2}{a^2 - h^2} h.$$

117. A person at distance of 20 yards from the nearer of two towers in the same straight line with him, and 10 yards apart, observes them to subtend the same angle. Passing the nearer tower a certain distance, he observes them again subtend the same angle, the complement of the former. The heights of the towers are  $6\sqrt{5}$ ,  $4\sqrt{5}$  yards respectively.

118. A person at distance  $a$  from a tower, observes the elevation  $\alpha$  of the tower, and of the top of the flagstaff ( $90^\circ - \alpha$ ); the height of flagstaff is  $2a \cot 2\alpha$ .

If the distance from the tower is unknown, and on receding  $c$  feet the elevation of tower is  $\frac{\alpha}{2}$ ; then the height of flagstaff is  $c \operatorname{cosec} \alpha \cos 2\alpha$ .

119. From each of two stations in the same horizontal plane at distance  $D$  from each other, a pillar on a hill in the same vertical plane with the stations is observed to subtend the same angle at each of the two stations, and the elevations of the top of the pillar are  $\epsilon$ ,  $\epsilon'$  at the two stations respectively. The height of pillar is

$$\frac{D \cos (\epsilon' + \epsilon)}{\sin (\epsilon' - \epsilon)}.$$

120. The length of the shadow of a tower, height  $h$ , is observed twice in the same day to be  $a$ ,  $b$ , and the difference of the altitude of the sun on the two occasions was  $\alpha$ , shew that


$$h^2 + (a - b) h \cot \alpha + ab = 0;$$

and explain the two values of  $h$  obtained from this equation.

## SECTION IV.

RELATIONS BETWEEN THE SIDES AND ANGLES OF A TRIANGLE. THE RADII OF THE CIRCUMSCRIBED AND INSCRIBED CIRCLES.

36. We now proceed to establish some important relations between the sides and the Trigonometrical Ratios of the angles of a triangle.



Let  $ABC$  be a triangle, from  $A$  draw  $AD$  perpendicular to  $BC$  or  $BC$  produced : then denoting the angles of the triangle by  $\alpha, \beta, \gamma$ , and the sides opposite to them by  $a, b, c$  respectively, we have

$$\sin \beta = \frac{AD}{AB}, \text{ or } AD = c \sin \beta ;$$

$$\sin \gamma = \sin ACD = \frac{AD}{AC}, \text{ or } AD = b \sin \gamma.$$

Hence

$$c \sin \beta = b \sin \gamma,$$

or

$$\frac{\sin \beta}{b} = \frac{\sin \gamma}{c}.$$

Similarly, by dropping a perpendicular from  $B$  on  $AC$ , we have,

$$\frac{\sin \alpha}{a} = \frac{\sin \gamma}{c};$$

and therefore

$$\frac{\sin \alpha}{a} = \frac{\sin \beta}{b} = \frac{\sin \gamma}{c}.$$

B. T.

E

37. Again we have

$$BD = AB \cos \beta = c \cos \beta,$$

$$CD = AC \cos ACD = b \cos \gamma, \text{ if } \gamma \text{ is acute;}$$

$$= -b \cos \gamma, \text{ if } \gamma \text{ is obtuse.}$$

But  $a = BC = BD + CD$  in the first case,  
 $= BD - CD$  in second case;

and therefore in both cases,

$$a = c \cos \beta + b \cos \gamma.$$

This may be derived from the forms in the preceding Article as follows;

since  $a = 180 - (\beta + \gamma)$ ,

$$c \sin a = c \sin (\beta + \gamma) = c \sin \beta \cos \gamma + c \cos \beta \sin \gamma,$$

or

$$a \sin \gamma = b \sin \gamma \cos \gamma + c \cos \beta \sin \gamma;$$

and therefore

$$a = b \cos \gamma + c \cos \beta,$$

as before.

Similar forms may be proved for each of the other sides of the triangle.

38. Again  $(a - b \cos \gamma)^2 = (c \cos \beta)^2$ ;

$$\begin{aligned} \therefore a^2 - 2ab \cos \gamma + b^2 \cos^2 \gamma &= c^2 - c^2 \sin^2 \beta \\ &= c^2 - b^2 \sin^2 \gamma; \end{aligned}$$

$$\therefore c^2 = a^2 + b^2 - 2ab \cos \gamma,$$

and so for the other sides. This may also be put in the form

$$\cos \gamma = \frac{a^2 + b^2 - c^2}{2ab}.$$

This formula may be obtained geometrically as follows.

By Euclid, II. 13, if  $ACB$  is acute,

$$AB^2 = BC^2 + AC^2 - 2BC \cdot CD;$$

and by Euclid, II. 12, if  $ACB$  is obtuse,

$$AB^2 = BC^2 + AC^2 + 2BC \cdot CD.$$

In first case  $CD = b \cos \gamma$ , in second case  $CD = -b \cos \gamma$ .

Hence in both cases

$$c^2 = a^2 + b^2 - 2ab \cos \gamma.$$

39. From the forms in the last Article, we have

$$1 - \cos \gamma = 1 - \frac{a^2 + b^2 - c^2}{2ab} = \frac{c^2 - (a - b)^2}{2ab},$$

or 
$$2 \sin^2 \frac{\gamma}{2} = \frac{(c - a + b)(c + a - b)}{2ab}.$$

Let  $2s = a + b + c$ ,

and therefore  $s - a = \frac{b + c - a}{2}$ , and so for  $s - b$ ,  $s - c$ .

And we have 
$$\sin^2 \frac{\gamma}{2} = \frac{(s - a)(s - b)}{ab}.$$

Again, 
$$1 + \cos \gamma = \frac{(a + b)^2 - c^2}{2ab} = \frac{(a + b + c)(a + b - c)}{2ab};$$

$$\therefore \cos^2 \frac{\gamma}{2} = \frac{s(s - c)}{ab}.$$

So also 
$$\tan^2 \frac{\gamma}{2} = \frac{(s - a)(s - b)}{s(s - c)}.$$

And 
$$\sin^2 \gamma = 4 \sin^2 \frac{\gamma}{2} \cos^2 \frac{\gamma}{2} = 4 \frac{s(s - a)(s - b)(s - c)}{a^2 b^2},$$

or 
$$\sin \gamma = \frac{2}{ab} \sqrt{s(s - a)(s - b)(s - c)}.$$

Similar forms may be proved for the other angles.

From these forms the formulæ of Art. 36 may be directly deduced.

40. *To find the area of a triangle.* The area of a triangle is equal to the rectangle contained by half the base, and the perpendicular upon the base from the opposite angle, that is,

$$\text{area } ABC = \frac{1}{2} BC \cdot AD = \frac{1}{2} ab \sin \gamma;$$

this gives the area in terms of two sides and the included angle.


Substituting for  $\sin \gamma$  from the preceding Article, we have

$$\text{area } ABC = \sqrt{s(s - a)(s - b)(s - c)};$$

which is in terms of the sides only.

This expression is frequently denoted by  $S$ .

41. To find the radius of the circumscribed circle.



Let  $ABC$  be a triangle,  $BE$  the diameter of the circumscribed circle. Then angle  $BEC = BAC = \alpha$ ; and  $BCE$  is a right angle, and we have

$$\frac{BC}{BE} = \sin \alpha, \quad \text{or } 2R = \frac{a}{\sin \alpha};$$

which gives  $R$  in terms of one side and the angle opposite to it,  $R$  being the radius of the circle.

Substituting for  $\sin \alpha$  in terms of the sides we find

$$R = \frac{abc}{4S}.$$

These results may also be obtained from the construction for finding the centre of the circumscribed circle given in Euclid, iv. 5.

For  $BC$  being bisected in  $D$ , and  $O$  being the centre, angle<sup>1o</sup>  $BOC =$  twice the angle at circumference


$$= 2\alpha;$$

and therefore  $BOD = \alpha, \quad BD = \frac{a}{2},$

and  $\frac{a}{2} = R \sin \alpha,$  as before.

42. To find the radius of the inscribed circle.


Let  $O$  be the centre of the inscribed circle;  $OA'$ ,  $OB'$ ,  $OC'$  perpendicular to the sides and equal to  $r$ . Then the area of triangle



$BOC$  is  $\frac{1}{2}ra$ , and so also for triangles  $BOA$ ,  $COA$ . But area of triangle  $ABC$  equals the sum of these triangles,

or 
$$\frac{1}{2}ra + \frac{1}{2}rb + \frac{1}{2}rc = \text{area of } ABC = S;$$

$$\therefore r = \frac{S}{s}.$$



To find the radii of the escribed circles, that is, of the circles which touch one side, and the other two sides produced. Let  $O'$  be centre of the circle touching  $BC$ .



Then  $O'A' = O'B' = O'C' = r$ ,

and  $\Delta ABC = \Delta BAO' + \Delta CAO' - \Delta BCO'$ ,

or  $S = \frac{1}{2} r_a c + \frac{1}{2} r_a b - \frac{1}{2} r_a a$ ,

or  $r_a = \frac{S}{s-a}$ ;

so also  $r_b = \frac{S}{s-b}$ ,  $r_c = \frac{S}{s-c}$ .

### EXAMPLES. (A).

Prove the formulæ :

$$1. \frac{\sin \alpha + \sin \beta}{\sin \beta} = \frac{a+b}{b}.$$

$$2. \frac{\sin \alpha + \sin \beta}{\sin \alpha - \sin \beta} = \frac{a+b}{a-b}.$$

$$3. \frac{\sin \alpha + \sin \beta - \sin \gamma}{\sin \alpha - \sin \beta + \sin \gamma} = \frac{a+b-c}{a-b+c}.$$

$$4. \frac{\sin \alpha - \sin \beta}{a-b} = \frac{\sin \gamma}{c}.$$

$$5. \frac{a+c-b}{4c} = \frac{b \sin^2 \frac{\alpha}{2}}{a+b-c}.$$

$$6. \cos \gamma = \frac{\sin^2 \alpha + \sin^2 \beta - \sin^2 \gamma}{2 \sin \alpha \sin \beta}.$$

$$7. a+b+c = (a+b) \cos \gamma + (a+c) \cos \beta + (b+c) \cos \alpha.$$

$$8. (a+b)(1-\cos \gamma) = c(\cos \alpha + \cos \beta).$$

$$9. \frac{\sin \alpha + \sin \beta}{\cos \alpha + \cos \beta} = \frac{\sin \gamma}{1-\cos \gamma} = \cot \frac{\gamma}{2}.$$

$$10. c(1-\cos^2 \beta - \cos^2 \alpha) = \cos \gamma (a \cos \alpha + b \cos \beta).$$

$$11. \text{ If } a=b, \text{ shew that } \alpha=\beta.$$

12. If  $c \cos \beta = b \cos \gamma$ , shew that the triangle is isosceles ( $b = c$ ).
13. If  $a \sec \beta = 2c$ , the triangle is isosceles ( $b = c$ ).
14. If the triangle is isosceles, ( $a = b$ ),  $c = 2a \sin \frac{\gamma}{2}$ .
15. If  $c = 2a \sin \frac{\gamma}{2}$ , then the triangle is either isosceles, or  

$$c = \sqrt{a(a-b)}.$$
16. If  $(s-a)(s-b) = ab$ , the triangle is impossible.
17. If  $s(s-c) = \frac{ab}{2}$ , one angle is a right angle.
18. The area of a triangle whose sides are 4, 5, 7 inches respectively, is  $4\sqrt{6}$  square inches.
19. Two sides of a triangle are 8 and 10 inches, and the included angle is  $30^\circ$ ; the area is 20 square inches.
20. The sides of a triangle are in A.P. and their common difference is 2 inches. If the area is  $3\sqrt{15}$  square inches, find the sides.  
*Ans.* 4, 6, 8 inches.
21. If the area of a triangle is equal to 84 square inches, and two of its sides are 15 and 13 inches, find the third side.  
*Ans.* 14 inches.
22. The sides of a triangle are 3, 7, 8; compare the radii of the inscribed and circumscribed circles.  
*Ans.*  $R : r = 7 : 2$ .
23. If the area of a triangle is  $\frac{1}{2}ab$ , then the angle  $\gamma$  is a right angle.
24. If  $R = \frac{c}{2}$ , then  $\gamma = 90^\circ$ .
25. If  $r = s - c$ , then  $a^2 + b^2 = c^2$ .
26. Shew that  $r_a + r_b = c \cot \frac{\gamma}{2}$ .
27. If  $r_a = r_b = \frac{1}{2}r_c$ , then  $c = \frac{4}{3}a$ .

$$28. \text{ If } r_a + r_b = r_c, \text{ then } c = \frac{a + b \pm 2\sqrt{a^2 - ab + b^2}}{3}.$$

$$29. b \cos^2 \frac{\gamma}{2} + c \cos^2 \frac{\beta}{2} = \frac{a + b + c}{2}.$$

$$30. (b - c) \cos \frac{a}{2} = a \sin \frac{\beta - \gamma}{2}.$$

$$31. \frac{\sin^2 \frac{\beta}{2}}{b} + \frac{\sin^2 \frac{\gamma}{2}}{c} = \frac{s - a}{bc}.$$

$$32. \cot \frac{a}{2} + \cot \frac{\beta}{2} = \frac{c}{r}.$$

$$33. r = s \tan \frac{a}{2} \tan \frac{\beta}{2} \tan \frac{\gamma}{2}.$$

$$34. \frac{a}{b} \cdot \frac{1 - \cos \beta}{1 - \cos \alpha} = \tan \frac{\beta}{2} \cot \frac{a}{2}.$$

$$35. \frac{\sin \alpha}{\cot \frac{\beta}{2} + \cot \frac{\gamma}{2}} = 2 \sin \frac{a}{2} \sin \frac{\beta}{2} \sin \frac{\gamma}{2}.$$

$$36. \frac{b - c}{r_a} + \frac{c - a}{r_b} + \frac{a - b}{r_c} = 0.$$

$$37. (a + c) \sin \frac{\beta}{2} = b \cos \frac{a - \gamma}{2}.$$

$$38. \text{ If } \sin \alpha, \sin \beta, \sin \gamma \text{ are in A.P., then } \tan \frac{a}{2} \tan \frac{\gamma}{2} = \frac{1}{3}.$$

39.  $ABC$  is an isosceles right-angled triangle, one of whose base angles  $B$  is trisected by the lines  $BD, BE$ . Shew that

$$AD : DE : EC = \sqrt{3} : 2 : \sqrt{3} - 1.$$

40. If  $AD$  is the perpendicular from  $A$  on the side  $BC$  of triangle  $ABC$ , and  $DE, DF$  are drawn perpendicular to  $AB$ , respectively, shew that  $EF \times R = \text{area of triangle } ABC$ .

EXAMPLES. (B).

Prove the following relations between the sides and angles of a triangle.

$$1. \quad a^2 + b^2 + c^2 = 2ab \cos \gamma + 2ac \cos \beta + 2bc \cos \alpha.$$

$$2. \quad a(b \cos \gamma - c \cos \beta) = b^2 - c^2.$$

$$3. \quad \cos \alpha + \cos \beta = 2 \frac{a+b}{c} \sin^2 \frac{\gamma}{2}.$$

$$4. \quad \tan \beta = \frac{b \sin \gamma}{a - b \cos \gamma}.$$

$$5. \quad a^2 \sin 2\beta + b^2 \sin 2\alpha = 2ab \sin \gamma.$$

$$6. \quad \cot \alpha - \cot \beta = \frac{b^2 - a^2}{ab \sin \gamma}.$$

$$7. \quad R = \frac{1}{8} (a + b + c) \sec \frac{\alpha}{2} \sec \frac{\beta}{2} \sec \frac{\gamma}{2}.$$

$$8. \quad \frac{1}{a} \cos^2 \frac{\alpha}{2} + \frac{1}{b} \cos^2 \frac{\beta}{2} + \frac{1}{c} \cos^2 \frac{\gamma}{2} = \frac{(a + b + c)^2}{4abc}.$$

$$9. \quad 1 - \tan \frac{\alpha}{2} \tan \frac{\beta}{2} = \frac{2c}{a + b + c}.$$

$$10. \quad \frac{\text{vers } \alpha}{\text{vers } \beta} = \frac{a + c - b}{b + c - a} \cdot \frac{a}{b}.$$

$$11. \quad \frac{\cot \frac{\beta}{2} + \cot \frac{\gamma}{2}}{\cot \frac{\alpha}{2}} = \frac{2a}{b + c - a}.$$

$$12. \quad 2Rr = \frac{abc}{a + b + c}.$$

$$13. \quad 4R \sin \frac{\alpha}{2} \sin \frac{\beta}{2} \sin \frac{\gamma}{2} = r.$$

$$14. \quad r_a r_b + r_b r_c + r_c r_a = s^2.$$

$$15. \quad a \cos \alpha + b \cos \beta + c \cos \gamma = 4R \sin \alpha \sin \beta \sin \gamma.$$

$$16. \quad \tan^2 \frac{\alpha}{2} = \frac{r r_a}{r_b r_c}.$$

$$17. \quad r_a r_b r_c = r s^3.$$

$$18. \quad r_a + r_b + r_c - r = 4R.$$

$$19. \quad \frac{\sin(\beta - \gamma)}{\sin \alpha} = \frac{b^2 - c^2}{a^2}.$$

$$20. \quad \cot \frac{\alpha}{4} - \operatorname{cosec} \frac{\alpha}{2} : \cot \frac{\beta}{2} + \cot \frac{\gamma}{2} = b + c - a : 2a.$$

Prove the following expressions for the area of a triangle.

$$21. \quad (r r_a r_b r_c)^{\frac{1}{2}}.$$

$$22. \quad Rr(\sin \alpha + \sin \beta + \sin \gamma).$$

$$23. \quad \frac{a^2 - b^2}{2} \frac{\sin \alpha \sin \beta}{\sin(\alpha - \beta)}.$$

$$24. \quad \frac{2abc}{a+b+c} \cos \frac{\alpha}{2} \cos \frac{\beta}{2} \cos \frac{\gamma}{2}.$$

$$25. \quad \frac{2s^2 \sin \alpha \sin \beta \sin \gamma}{(\sin \alpha + \sin \beta + \sin \gamma)^3}.$$

$$26. \quad \frac{a^2}{4} \left\{ \frac{\cos(\beta - \gamma)}{\sin \alpha} + \cot \alpha \right\}.$$

$$27. \quad s(s - a) \tan \frac{\alpha}{2}.$$

$$28. \quad \frac{1}{2} \sqrt{a^2 bc \cos \beta \cos \gamma + b^2 ac \cos \alpha \cos \gamma + c^2 ab \cos \alpha \cos \beta}.$$

$$29. \quad \frac{1}{4} \sqrt{2a^2 b^2 c^2 (\sin 2\alpha + \sin 2\beta + \sin 2\gamma)}.$$

30. The areas of all triangles described about the same circle vary as their perimeters.

31. If  $a^2, b^2, c^2$  be in A.P., then

$$\frac{\sin 3\beta}{\sin \beta} = \left( \frac{a^2 - c^2}{2ac} \right)^2.$$

32. If  $a, b, c$  be in A.P., then will  $r_a, r_b, r_c$  be in H.P.

33. Shew that  $R, r$  have a common measure if  $a, b, c$  have one.

34. The distances between the centres of the inscribed and the centres of the escribed circles are

$$4R \sin \frac{\alpha}{2}, 4R \sin \frac{\beta}{2}, 4R \sin \frac{\gamma}{2},$$

respectively.

35. If  $p_a, p_b, p_c$  be the perpendiculars from the angles on the opposite sides, then

$$p_a p_b p_c = \frac{8s^2 r^2}{abc}.$$

36. If  $l$  be the length of the line bisecting the angle  $a$ , then

$$l = \frac{2\sqrt{bcs \cdot (s-a)}}{b+c}.$$

37. If the perpendiculars from the angles of a triangle  $ABC$  on the opposite sides meet in  $O$ , then

$$\tan a = \frac{AO}{BC},$$

and  $AO : BO : CO = \sin a \tan a : \sin \beta \tan \beta : \sin \gamma \tan \gamma.$

38. If  $\tan a, \tan \beta, \tan \gamma$  are in G.P., then

$$a^4 + c^4 = b^4 (a^2 + c^2).$$

39. If  $a, b, c$  be the sides of a triangle inscribed in a circle radius  $R$ , then

$$R^2 (a^4 + b^4 + c^4 - 2a^2 b^2 - 2a^2 c^2 - 2b^2 c^2) + a^2 b^2 c^2 = 0.$$

## EXAMPLES. (C).

In any triangle, prove that

1.  $\cot \alpha + \cot \beta + \cot \gamma = \frac{a^2 + b^2 + c^2}{4S}.$
2.  $b^2 \sin 2\gamma - 2bc \sin (\beta - \gamma) - c^2 \sin 2\beta = 0.$
3.  $b^2 \cos 2\gamma + 2bc \cos (\beta - \gamma) + c^2 \cos 2\beta = a^2.$
4.  $(b^2 - c^2) \cot \alpha + (c^2 - a^2) \cot \beta + (a^2 - b^2) \cot \gamma = 0.$
5.  $\tan \frac{\alpha}{2} + \tan \frac{\beta}{2} + \tan \frac{\gamma}{2} + \frac{s}{r} = 4R \left( \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right).$
6.  $s = r \left( \cot \frac{\alpha}{2} + \cot \frac{\beta}{2} + \cot \frac{\gamma}{2} \right) = R (\sin \alpha + \sin \beta + \sin \gamma).$
7.  $a^2 \tan^2 \beta - 2ab \tan \beta \sin \gamma + b^2 = b^2 \cos^2 \gamma \sec^2 \beta.$
8.  $4r^2 s = abc (\cos \alpha + \cos \beta + \cos \gamma - 1).$
9.  $\frac{a}{r_a} - \frac{b}{r_b} = \tan \frac{\beta}{2} - \tan \frac{\alpha}{2}.$
10.  $\frac{\cos \frac{\alpha}{2} \cos \frac{\beta - \gamma}{2}}{b + c} = \frac{\cos \frac{\beta}{2} \cos \frac{\alpha - \gamma}{2}}{a + c} = \frac{\cos \frac{\gamma}{2} \cos \frac{\alpha - \beta}{2}}{a + b}.$
11.  $\cos^{-1} \sqrt{\frac{r_b r_c}{bc}} + \cos^{-1} \sqrt{\frac{r_a r_c}{ac}} + \cos^{-1} \sqrt{\frac{r_a r_b}{ab}} = 90^\circ.$
12.  $\frac{a^2 \sin (\beta - \gamma)}{\sin \alpha} + \frac{b^2 \sin (\gamma - \alpha)}{\sin \beta} + \frac{c^2 \sin (\alpha - \beta)}{\sin \gamma} = 0.$
13.  $(a + b + c) (\cos \alpha + \cos \beta + \cos \gamma)$   
 $= 2a \cos^2 \frac{\alpha}{2} + 2b \cos^2 \frac{\beta}{2} + 2c \cos^2 \frac{\gamma}{2}.$
14.  $S = s^2 \tan \frac{\alpha}{2} \tan \frac{\beta}{2} \tan \frac{\gamma}{2}.$

$$15. S = \sin \frac{a}{2} \sin \frac{\beta}{2} \sin \frac{\gamma}{2} \left\{ \frac{a^2}{\sin a} + \frac{b^2}{\sin \beta} + \frac{c^2}{\sin \gamma} \right\}.$$

16. The distance of the foot of the perpendicular from the angle  $A$  on the side  $a$ , from the middle point of  $a$ , is  $\frac{b^2 - c^2}{2a}$ .

17. If  $\cot a$ ,  $\cot \beta$ ,  $\cot \gamma$  are in arithmetical progression, so also are  $a^2$ ,  $b^2$ ,  $c^2$ .

18. If  $p_a$ ,  $p_b$ ,  $p_c$  be the perpendiculars from the angles on the opposite sides, then

$$\frac{1}{p_a} + \frac{1}{p_b} + \frac{1}{p_c} = \frac{1}{r}.$$

$$19. \text{ Shew that } r = \frac{a \sin \beta \sin \gamma}{\sin a + \sin \beta + \sin \gamma} = \frac{a \sin \frac{\beta}{2} \sin \frac{\gamma}{2}}{\cos \frac{a}{2}}.$$

$$20. \text{ Shew that } r_a = \frac{a \sin \beta \sin \gamma}{\sin \beta + \sin \gamma - \sin a} = \frac{a \cos \frac{\beta}{2} \cos \frac{\gamma}{2}}{\cos \frac{a}{2}}.$$

$$21. \text{ Shew that } r_a + r_b + r_c = 2R \left( \cos^2 \frac{a}{2} + \cos^2 \frac{\beta}{2} + \cos^2 \frac{\gamma}{2} \right).$$

22. If  $\gamma$  be a right angle,

$$\cos (2a - \beta) = \frac{a}{c^2} (3c^2 - 4a^2).$$

23. The perpendicular from  $C$  on the opposite side  $AB$  is

$$\frac{ab}{2} \cdot \frac{a \sin a + b \sin \beta + c \sin \gamma}{ab \cos \gamma + ac \cos \beta + bc \cos a}.$$

$$24. \text{ If } a : \beta : \gamma = 2 : 3 : 4, \text{ then } \frac{a+c}{2b} = \cos \frac{a}{2}.$$

25. The length of the line drawn bisecting the angle  $a$ , and meeting the side  $a$  is  $\frac{2bc}{b+c} \cos \frac{a}{2}$ .



26. In any triangle the perpendicular from the angle  $A$  on the side  $BC$  is equal to  $\frac{b^2 \sin \gamma \pm c^2 \sin \beta}{b \pm c}$ .

27. If  $A'B'C'$  be the triangle formed by joining the feet of the perpendiculars from the angles  $A, B, C$  on the opposite sides, then  $BC' = R \sin 2\alpha$ .

28. If  $a'b'c'$  be the sides of the triangle  $A'B'C'$ , then

$$\frac{a'}{a^2} + \frac{b'}{b^2} + \frac{c'}{c^2} = \frac{a^2 + b^2 + c^2}{2abc},$$

and  $a' + b' + c' = \frac{abc}{2R^2}$ .

29. If  $a^2, b^2, c^2$  be in A.P., then  $a \sec \alpha, b \sec \beta, c \sec \gamma$  are in H.P.

30. If  $\tan \frac{\alpha}{2} \tan \frac{\beta}{2} = \frac{n-1}{n+1}$ , then

$$\tan \frac{\alpha}{2} + \tan \frac{\beta}{2} + \tan \frac{\gamma}{2} = \frac{\sin \alpha}{n - \cos \alpha} + \frac{2}{n+1} \cdot \frac{n - \cos \alpha}{\sin \alpha}.$$

31. The distance between the centres of the escribed circles touching the sides  $a, b$  is

$$\frac{2c \sqrt{ab}}{\sqrt{c^2 - (a-b)^2}}.$$

32. If  $s, s'$  be the semi-perimeters of two triangles, and  $r_a = r'_a$ ; then  $\frac{s-a}{s} : \frac{s'-a'}{s'}$  is the ratio of the radii of the inscribed circles.

33. Prove that  $\frac{\sin \beta}{\sin(\alpha - \beta)} = \frac{b}{a \cos \beta - b \cos \alpha}$ .

34. If  $p, q, r$  be the perpendiculars from the angles upon the opposite sides  $a, b, c$ , respectively; then  $\frac{p^2}{qr} = \frac{bc}{a^2}$ .

35. The distance of the points of contact of the inscribed circle with the sides  $AB, AC$  from the angle  $A$  are  $s - a$ .

36. If  $a, b, c$  be in arithmetical progression, then

$$\tan \frac{a}{2} \tan \frac{\gamma}{2} = \frac{1}{3}.$$

37. If the sides of a triangle are in arithmetical progression, then the cotangents of the half angles are in arithmetical progression.

38. In a right-angled triangle ( $\gamma$  the right angle),

$$\sin \frac{\beta}{2} = \sqrt{\left(\frac{c-a}{2c}\right)}.$$

39. If  $O$  be the centre of the circles inscribed in a triangle  $ABC$ , and  $R_a, R_b, R_c$  the radii of the circles circumscribed about the triangles  $BOC, AOC, AOB$ , respectively, then

$$\frac{R_a}{\sin \frac{a}{2}} = \frac{R_b}{\sin \frac{\beta}{2}} = \frac{R_c}{\sin \frac{\gamma}{2}}.$$

40. Shew also that the distance of the chords common to these circles and the inscribed circle are distant from  $O$ ,

$$\frac{r^2}{2R_a}, \frac{r^2}{2R_b}, \frac{r^2}{2R_c},$$

respectively.

41. If  $O$  be the centre and  $D, E, F$  the points of contact of the circle inscribed in a triangle  $ABC$ , then

$$OA \cdot OB \cdot OC (AF + BD + CE) = 4R \cdot AF \cdot BD \cdot CE.$$

42. If  $O$  be the centre of the inscribed circle of a triangle  $ABC$ , and  $O_a, O_b, O_c$  the centres of the escribed circles, shew that

$$\frac{AO}{AO_a} + \frac{BO}{BO_b} + \frac{CO}{CO_c} = 1.$$

43. If  $R_a, R_b, R_c$  be the radii of the circles circumscribed about  $AEF, BDF, CDE$ , (Ex. 41) respectively, then

$$aR_a^2 + bR_b^2 + cR_c^2 = \frac{abc}{4}.$$

44. If  $AD$  be drawn through  $O$  the centre of the inscribed circle then the radii of the circles circumscribing  $ADB, ADC$  are together equal to  $2R \cos \frac{a}{2}$ .

45. If  $\rho_a, \rho_c$  be the radii of circles passing through  $A$  and touching the side  $BC$  of the triangle  $ABC$  in the points  $B, C$ , then  $\rho_a \rho_c = R^2$ , and the radius of circle touching  $BC$  in its middle point is

$$\frac{2(b^2 + c^2) - a^2}{8b \sin \gamma}.$$

46. If  $A'B'C'$  be points in the sides  $BC, AC, AB$  of a triangle such that

$$BA' = \frac{1}{3}a, \quad CB' = \frac{1}{3}b, \quad AC' = \frac{1}{3}c;$$

then the area of the triangle  $A'B'C'$  is  $\frac{S}{3}$ .

47. An isosceles triangle is described about two circles (radii  $r, r'$ ) which touch each other, so that its equal sides touch both the circles, and its base the larger circle; shew that its base is  $2r \sqrt{\frac{r}{r'}}$ , and its sides are  $r \cdot \frac{r+r'}{r-r'} \cdot \sqrt{\frac{r}{r'}}$ .

48. If  $O$  be the centre of the circumscribed circle, and  $r_1, r_2, r_3$  the radii of the circles circumscribed about  $COB, AOC, BOA$  respectively, then

$$r_1 \cos \alpha = r_2 \cos \beta = r_3 \cos \gamma.$$

49. If  $D$  be any point in the sides  $BC$  of a triangle  $ABC$ , and if  $k$  be the harmonic mean between the radii of the circles described about the triangles  $ADB, ADC$ , then

$$\frac{AD}{k} = \sin \beta + \sin \gamma.$$

50.  $ABC$  is an isosceles triangle ( $AB=AC$ ): on  $BC$  as diameter a circle is described meeting  $AB, AC$  in  $D, E$ ;  $DC, EB$  intersect in  $O$ . Shew that area  $ABOC = \frac{1}{2}a^2 \cot \alpha$ .

51. A tangent drawn from a point outside a circle (radius  $r$ ) makes an angle  $2\theta$  with the diameter through that point. Shew that the distance of the point from the circle is

$$\frac{r(1 - \tan \theta)^2}{2 \tan \theta}.$$

52. If sides  $BA$ ,  $CA$  of a triangle be produced to  $D$ ,  $E$  so that  $AD = AE$ , and the area of the triangle  $ADE$  equals the area of the triangle  $ABC$ : shew that the radius of the circle inscribed in  $DAE$  is equal to

$$\frac{S}{\sqrt{bc} \left( 1 + \sin \frac{\alpha}{2} \right)}.$$

53. If lines be drawn bisecting the angles of a parallelogram, the area of the rectangle formed by these lines equals  $\frac{1}{2}(a-b)^2 \sin \alpha$ , where  $a$ ,  $b$  are the sides of the parallelogram, and  $\alpha$  the angle between them.

54. If a triangle have sides, which vary inversely as the Harmonic means of the radii of the escribed circles taken two and two, it is similar to the original triangle.

55. Find the length of the line bisecting the triangle  $ABC$  and making equal angles with the sides  $AB$ ,  $AC$ , and find the condition that the problem is possible.

*Ans.* length =  $\sqrt{2 \cdot s - b \cdot s - c}$ , and one side must not be greater than twice the other.

56. Three circles (radii  $r_1$ ,  $r_2$ ,  $r_3$ ) touch one another: shew that the radius of the circle passing through the three points of contact is

$$\sqrt{\frac{r_1 r_2 r_3}{r_1 + r_2 + r_3}}.$$

Also if  $a$ ,  $b$ ,  $c$  be the chords joining the points of contact of the three circles, then

$$\frac{8}{abc} = \left( \frac{1}{r} + \frac{1}{r_2} \right) \left( \frac{1}{r_2} + \frac{1}{r_1} \right) \left( \frac{1}{r_3} + \frac{1}{r_1} \right).$$

## SECTION V.

SOLUTION OF TRIANGLES. FOUR CASES. THEIR TREATMENT  
IN EUCLID. APPLICATION.

43. EVERY triangle has six parts, three angles and three sides ; these parts are not independent of each other, for the three angles are always equal to two right angles, that is, we always have

$$\alpha + \beta + \gamma = 180^\circ \dots\dots\dots \text{I.}$$

Also between the sides and angles two other relations independent of each other can be proved to exist. Take for instance the relations proved in Art. 36,

$$\frac{\sin \alpha}{a} = \frac{\sin \beta}{b} \dots\dots\dots \text{II.}$$

$$\frac{\sin \alpha}{a} = \frac{\sin \gamma}{c} \dots\dots\dots \text{III.}$$

From I. II. III. all other relations between the sides and angles may be deduced. If we had taken any other two of the relations proved in the last section which are independent of each other, we could with equation I. have deduced all the rest.

Hence since we have three and only three independent equations amongst the six parts of a triangle, if three parts are given, we are enabled to determine the other three. Also three parts at least must be given ; but if more were, we should find one or more equations between them to express the condition that these parts may belong to the same triangle.

We may observe also, that the three given parts must not all be angles ; for by equation I. if we know two angles we also know the other, and so having three angles given would really amount to knowing only two parts ; one part then at least must be a side. Also we must not have two sides given us less than the third side, nor such as to make the sines and cosines involved in our formulæ greater than unity. (See Art. 46.)

The process of finding the other parts of a triangle, when three of them are known, is called solving the triangle.

44. In solving triangles we shall find four distinct cases present themselves.

CASE I. When the three sides are given.

CASE II. When two sides and an angle opposite to one of them are given.

CASE III. When two sides and the included angle are given.

CASE IV. When two angles and a side are given.

We shall treat each of these in order.

45. CASE I. Let  $a, b, c$  be given; to find  $\alpha, \beta, \gamma$ , we have, Art. 38,

$$\cos \alpha = \frac{b^2 + c^2 - a^2}{2bc},$$

$$\cos \beta = \frac{a^2 + c^2 - b^2}{2ac},$$

$$\cos \gamma = \frac{a^2 + b^2 - c^2}{2ab};$$

these equations completely determine the angles  $\alpha, \beta, \gamma$ , but are not used because they are not adapted for logarithmic computation, as will be shewn hereafter (Art. 61). In practice we use the forms derived from them in Art. 39.

46. CASE II. Let  $a, b, \alpha$  be known: to find  $c, \beta, \gamma$ :

we have  $\sin \beta = \frac{b}{a} \sin \alpha,$

from which we find  $\beta$ : we then know


$$\gamma = 180 - (\alpha + \beta),$$

which gives  $\gamma$ : and we then know  $c$  from the equation

$$c = a \frac{\sin \gamma}{\sin \alpha}.$$

This case presents a peculiarity which we will now point out.

Let the angle  $BAC = \alpha$ ;  $AC = b$ ;  $CE$ , any line drawn from  $C$ , equal to  $a$ . With centre  $C$  and distance  $CE$  describe a circle cutting  $AB$  in  $B, B'$ : join  $CB, CB'$ ; draw  $CD$  perpendicular to  $AB$ .



If  $CE < CD$ , the circle will not cut  $AB$  at all, and there will be no triangle with the parts proposed, that is, there is no solution.

If  $CE > CD$  and  $< CA$ , two triangles will be formed,  $CAB$ ,  $CAB'$ , each having the angle  $CAB = \alpha$ , the sides  $CA = b$  and  $CB$  or  $CB'$  equal to  $a$ : that is to say, there are two solutions of the triangle.

If  $CE > CA$ , then the circle will meet  $AB$  in points  $B_1$ ,  $B_1'$ . Now  $CAB_1$  is not a solution, inasmuch as the angle  $CAB$  or  $\alpha$  is not a part of it, but  $CAB_1$  or  $180^\circ - \alpha$  instead. Hence in this case there is only one solution  $CAB_1'$ .

Now let us examine how these cases are indicated by the Trigonometrical formulæ. We have  $\sin \beta = \frac{b \sin \alpha}{a}$ .

Now  $CD = b \sin \alpha$ ,  $CE = a$ , and if  $CE < CD$ ,  $\sin \beta$  is greater than unity, which is impossible, and therefore there can be no solution in this case.

If  $CE > CD$  and  $< CA$ , then  $\sin \beta$  is possible, and  $b > a$  and therefore  $\beta > \alpha$ . Now two values  $\beta$  and  $180 - \beta$  will satisfy the equation

$$\sin \beta = \frac{b}{a} \sin \alpha,$$

and since  $\beta$  may be either greater or less than  $90^\circ$  both these values will give solutions of the triangle.

If  $CE > CA$ ,  $b < a$ , and therefore  $\beta < \alpha$  and  $\beta$  must be less than  $90^\circ$ ; and only the smaller of the values  $\beta$  and  $180 - \beta$  will give a solution of the triangle.

From the fact that when  $b > a$  there are two solutions of the triangle, this case is called the ambiguous case.

When  $a$  is obtuse,  $\beta$  must be acute, and  $a > b$ , and there is no ambiguity.

47. CASE III. Let  $a, b, \gamma$  be known: to find  $c, \alpha, \beta$  we have

$$c^2 = a^2 + b^2 - 2ab \cos \gamma;$$

this gives the side  $c$ : we can find  $\alpha$  and  $\beta$  from the formulæ

$$\sin \alpha = \frac{a}{c} \sin \gamma,$$

$$\sin \beta = \frac{b}{c} \sin \gamma.$$

In determining  $\alpha, \beta$  in this manner we should have to consider the order of magnitude of  $\alpha, \beta, \gamma$  from knowing that of  $a, b, c$ ; and so determine whether  $\alpha$  or  $180 - \alpha, \beta$  or  $180 - \beta$  are the true solutions. No ambiguity will really present itself.

The following method will avoid this difficulty as well as be better adapted for logarithms, as will be shewn hereafter.

Since 
$$\frac{\sin \alpha}{\sin \beta} = \frac{a}{b},$$

we have 
$$\frac{\sin \alpha - \sin \beta}{\sin \alpha + \sin \beta} = \frac{a - b}{a + b},$$

or 
$$\frac{2 \sin \frac{\alpha - \beta}{2} \cos \frac{\alpha + \beta}{2}}{2 \cos \frac{\alpha - \beta}{2} \sin \frac{\alpha + \beta}{2}} = \frac{a - b}{a + b},$$

or 
$$\tan \frac{\alpha - \beta}{2} = \frac{a - b}{a + b} \tan \frac{\alpha + \beta}{2}.$$

Now 
$$\frac{\alpha + \beta}{2} = 90^\circ - \frac{\gamma}{2};$$

$$\therefore \tan \frac{\alpha - \beta}{2} = \frac{a - b}{a + b} \cot \frac{\gamma}{2}.$$



This equation will give  $\frac{a-\beta}{2}$ , and since  $a+\beta$  is known we can find  $a$  and  $\beta$ .

The side  $c$  will then be found from the equation

$$c = a \frac{\sin \gamma}{\sin \alpha}.$$

48. CASE IV. First, let  $\alpha, \beta, \gamma$  be given; to find  $b, c, a$ . We have

$$\alpha = 180^\circ - (\beta + \gamma),$$

which gives  $\alpha$ , and then  $b, c$  are determined from the equations

$$b = a \frac{\sin \beta}{\sin \alpha},$$

$$c = a \frac{\sin \gamma}{\sin \alpha}.$$

Secondly, let  $\alpha, a, \beta$  be given; to find  $b, c, \gamma$ . We have

$$\gamma = 180^\circ - (\alpha + \beta),$$

and then  $b, c$  are found as before.

49. It will perhaps be of advantage to the student to compare these cases with the corresponding propositions in Euclid.


Euclid (I. 8) proves that if two triangles have their sides equal, they have their angles equal; in other words, that if three sides of a triangle are fixed, then the angles cannot vary; this agrees with Case I.

Euclid (I. 4) in the same way agrees with Case III., and (I. 26) with the two parts of Case IV.

Case II. we have already considered geometrically, but the student would do well to compare it with Euclid, VI. 7.

50. We give the following as an example of the application of the rules in this section to practical purposes.

Let it be required to find the distance between the summits  $P, Q$  of two inaccessible mountains. At  $A$  let the observer measure the angles  $PAB = \alpha$ ,  $QAB = \alpha'$ ,  $PAQ = \gamma$ ; at  $B$  let him



measure  $PBA = \beta$ ,  $QBA = \beta'$ , and also let him measure the distance  $AB = a$ . Let  $AP = x$ ,  $AQ = y$ . Then he would proceed with his calculations as follows :

$$AP = x = a \frac{\sin \beta}{\sin (\alpha + \beta)}, \quad y = a \frac{\sin \beta'}{\sin (\alpha' + \beta')},$$

which give  $x$  and  $y$ .  $PQ$  is then known from the equation

$$PQ^2 = x^2 + y^2 - 2xy \cos \gamma.$$

EXAMPLES. (A).

1. If  $a = 2$ ,  $b = 3$ ,  $c = 4$  ; solve the triangle.  
*Ans.*  $\alpha = 28^\circ. 57'$  ;  $\beta = 46^\circ. 34'$  ;  $\gamma = 104^\circ. 29'$ .
2. If  $a = 4$ ,  $b = 5$ ,  $c = 6$  ; solve the triangle.  
*Ans.*  $\alpha = 41^\circ. 36'$  ;  $\beta = 55^\circ. 46'$  ;  $\gamma = 82^\circ. 38'$ .
3. If  $a = 3$ ,  $b = 7$ ,  $c = 5$  ; solve the triangle.  
*Ans.*  $\alpha = 21^\circ. 47'$ ,  $\beta = 120^\circ$ ,  $\gamma = 38^\circ. 13'$ .
4. If  $a = 5$ ,  $b = 6$ ,  $c = 7$  ; solve the triangle.  
*Ans.*  $\alpha = 44^\circ. 25'$ ,  $\beta = 57^\circ. 7'$ ,  $\gamma = 78^\circ. 28'$ .
5. If  $a = 11$ ,  $b = 15$ ,  $\gamma = 78^\circ. 28'$  ; solve the triangle.  
*Ans.*  $c = 16.7332$  nearly,  $\alpha = 40^\circ. 6'$ ,  $\beta = 61^\circ. 26'$ .
6. If  $a = 9$ ,  $b = 10$ ,  $\gamma = 60^\circ$ , find  $c$ . *Ans.*  $c = 9.539$ .
7. If  $a = 21$ ,  $b = 15\sqrt{2}$ ,  $\gamma = 45^\circ$  ; solve the triangle.  
*Ans.*  $c = 16.1554$ ,  $\alpha = 66^\circ. 48'$ ,  $\beta = 68^\circ. 12'$ .
8. If  $\alpha = 68^\circ. 13'$ ,  $\beta = 45^\circ. 35'$ ,  $a = 25$  ; find  $b$ .  
*Ans.*  $b = 19.2$  nearly.

9. If  $\alpha = 60^\circ$ ,  $b = 20$ ,  $c = 10$ ; solve the triangle.

*Ans.*  $\beta = 90^\circ$ ,  $\gamma = 30^\circ$ ,  $a = 17.32$ .

10. If  $\alpha = 38^\circ.13'$ ,  $\beta = 21^\circ.47'$ ,  $c = 77$ ; solve the triangle.

*Ans.*  $a = 55$ ,  $b = 33$ ,  $\gamma = 120^\circ$ .

11. If  $\beta = 30^\circ$ ,  $\gamma = 45^\circ$ ,  $a = 7$ ; solve the triangle.

*Ans.*  $\alpha = 105^\circ$ ,  $b = 3.62$ ,  $c = 5.12$ .

12. If  $b = d\sqrt{2}$ ,  $c = (\sqrt{3} + 1)d$ ,  $\alpha = 45^\circ$ ; solve the triangle.

*Ans.*  $\alpha = 2d$ ,  $\beta = 30^\circ$ ,  $\gamma = 105^\circ$ .

13. Having given the lines  $p$ ,  $q$ ,  $r$  drawn from the angles  $A$ ,  $B$ ,  $C$  of a triangle to bisect the opposite sides, shew that

$$a = \frac{2}{3} \sqrt{2(q^2 + r^2) - p^2}.$$

If  $\gamma$  is a right angle, shew that  $p^2 + q^2 = 5r^2$ .

14. A lake is bounded by a vertical cliff whose height is equal to the breadth of the lake ( $a$ ): from a balloon above the lake they subtend the same angle  $\alpha$ . Shew that the height of the balloon is

$$\frac{a}{2 \sin \alpha} (\sin \alpha + \cos \alpha).$$

15. The courses of two ships are N., and E., and their rates of sailing are as  $1 : \sqrt{2}$ . The bearing of the former to the latter was E.N.E., but after the latter had sailed 4 miles, it was N.W. What was the original distance of the ships?

*Ans.*  $2\sqrt{20 - 14\sqrt{2}}$  miles.

16. If in a triangle  $ABC$ ,  $AD$ ,  $BE$ ,  $CF$  are drawn perpendicular to the sides from the opposite angles, and a new triangle is formed whose sides are  $AD$ ,  $BE$ ,  $CF$ , shew that the cosine of the angle opposite to  $AD$  is  $\frac{a^2b^2 + a^2c^2 - b^2c^2}{2a^2bc}$  where  $a$ ,  $b$ ,  $c$  are the sides of  $ABC$ .

17. If in a triangle, whose sides are  $a$ ,  $b$ ,  $c$ , the lines drawn from the angles to bisect the opposite sides are  $p_1$ ,  $p_2$ ,  $p_3$ , then

$$p_1^2 + p_2^2 + p_3^2 = \frac{3}{4}(a^2 + b^2 + c^2).$$

EXAMPLES. (B).

1. If  $a = 3$ ,  $b = 5$ ,  $\alpha = 21^\circ.47'$ ; find  $c$ .

*Ans.*  $c = 7$ , or  $2.28$  nearly.

2. If  $a = 5$ ,  $b = 6$ ,  $\alpha = 32^\circ.53'$ ; solve the triangle.

*Ans.*  $\beta = 40^\circ.39'$  or  $139^\circ.21'$ ;  $\gamma = 106^\circ.28'$  or  $7^\circ.46'$ ;

$c = 8.8$  or  $1.245$ .

3. In the ambiguous case, if  $a$ ,  $b$  be given, shew that the two values  $\gamma$ ,  $\gamma'$  of the third angle may be found from the equations

$$\gamma + \gamma' = 180^\circ - 2\alpha, \quad \cos \frac{\gamma - \gamma'}{2} = \frac{b}{a} \sin \alpha.$$

4. If  $c$ ,  $c'$  be the two values of the third side in the last question, then  $c + c' = 2b \cos \alpha$ ,  $c - c' = 2a \cos \beta$ .

Shew also that  $(c - c')^2 + (c + c')^2 \tan^2 \alpha = 4a^2$ .

Also  $cc' = b^2 - a^2$ .

5. In the ambiguous case if  $a + b = 2c$ , shew that  $a \sim b = \frac{c'}{2}$ .

6. In the ambiguous case if  $\sin \alpha = \frac{a}{2b}$ , shew that the angle between the two positions of  $a$  is  $120^\circ$ .

7. In the ambiguous case if  $k$ ,  $k'$  be the areas of the two triangles formed, shew that

$$k + k' = \frac{b^2}{2} \sin 2\alpha, \quad kk' = \frac{1}{4} b^2 (b^2 - a^2) \sin^2 \alpha.$$

8. The sides of a triangle are in A.P. and its area : the area of an equilateral triangle of the same perimeter =  $3 : 5$ ; shew that its sides are as  $3 : 5 : 7$  and one of its angles is  $120^\circ$ .

9. If  $\cos \gamma = \frac{\sin \alpha}{2 \sin \beta}$ , and  $\sin^2 \alpha = \sin^2 \beta + \sin^2 \gamma$ , find the angles.

*Ans.*  $\alpha = 90^\circ$ ,  $\beta = \gamma = 45^\circ$ .

10. If  $AD$  be drawn bisecting the angle  $A$  of a triangle, and  $\beta$ ,  $ADC = \delta$ ,  $DC = d$ , be known, solve the triangle.

11. If a triangle be formed whose sides are  $a + b$ ,  $a + c$ ,  $b + c$  respectively, and  $\gamma'$  is the angle opposite to  $a + b$ , then

$$\sin^2 \frac{\gamma'}{2} = \sin \frac{a}{2} \sin \frac{\beta}{2} \operatorname{cosec} \left( a + \frac{\beta}{2} \right) \operatorname{cosec} \left( \beta + \frac{a}{2} \right).$$

12. If  $a$ ,  $a$ ,  $b + c$  are known, shew that  $\beta$  and  $\gamma$  may be found from the equation

$$\cos \frac{\beta - \gamma}{2} = \frac{b + c}{a} \sin \frac{a}{2}.$$

13. Given  $\alpha$ ,  $\beta$ , and  $a + b$ ; solve the triangle.

14. Given the perimeter  $2s$ , the area  $S$ , and the angle  $\alpha$  of a triangle, find the side  $a$ .

$$\text{Ans. } a = s - \frac{S}{s} \cot \frac{\alpha}{2}.$$

15. The angles of a triangle are as 3, 4, 5, and the least side is  $\alpha$ ; find all the sides.

$$\text{Ans. } \frac{\alpha}{2} \sqrt{6}, \frac{\alpha}{2} (\sqrt{3} + 1).$$

16. Given the vertical angle, the base, and the difference of the two sides of the triangle; find the other angles.

17. A straight line of length  $p$  bisects the angle  $BAC$ , and divides the side  $BC$  into parts  $m$ ,  $n$ . Shew that

$$p^2 = bc - mn.$$

18. If perpendiculars be drawn from the angles of a triangle upon the opposite sides, and the feet of these perpendiculars be joined to form a new triangle, its sides are  $a \cos \alpha$ ,  $b \cos \beta$ ,  $c \cos \gamma$  respectively.

19. Find the height of a cloud by observing its elevation  $\alpha$ , and its depression  $\beta$  when seen by reflection in a lake from a station at a height  $h$  above the surface of the lake

$$\text{Ans. Height above the lake is } h \frac{\sin (\alpha + \beta)}{\sin (\beta - \alpha)}.$$

20.  $AB, CD$  are two towers in the same horizontal plane. The height of  $AB$  is  $h$ , the elevation of  $D$  at  $A$  is  $\alpha$ , at  $B$  is  $\beta$ .

Show that the height of  $CD$  is  $\frac{h \sin \alpha \cos \beta}{\sin (\alpha - \beta)}$ ;

and the distance  $AC$  is  $\frac{h \cos \alpha \cos \beta}{\sin (\alpha - \beta)}$ .

21. Two ships are lying at anchor at a known distance from each other: find by observations made on board the two ships the distance from either of them of an object on the shore.

22. Find  $c$  in the equation

$$b^2 + c^2 - 2bc \cos \alpha = a^2,$$

and shew that the ambiguous case is involved in the result.

23. From the top  $A$  of a tower  $AB$  (height  $h$ ) the angles of depression of two objects  $C, D$ , in the horizontal plane with the foot of the tower are observed to be  $45^\circ - \alpha, 45^\circ + \alpha$  respectively, and the angle subtended by  $CD$  is  $2\alpha$ : shew that  $CD = 2h \tan 2\alpha$ , and  $\angle ACD = 45^\circ - \alpha$ .

### EXAMPLES. (C.)

1. In any triangle  $\tan \left( \frac{\alpha}{2} + \beta \right) = \frac{c+b}{c-b} \tan \frac{\alpha}{2}$ .

2. A circle of radius  $r$  is inscribed in a sector of a circle of radius  $a$ , the chord of the sector being equal to  $2c$ ; shew that

$$\frac{1}{r} = \frac{1}{a} + \frac{1}{c}.$$

3. Two inaccessible objects  $C, D$  are observed from two stations  $A, B$  in the same plane with them, so that

$$CAD = \alpha = CBD, \quad BAD = \beta, \quad ABC = \gamma, \quad AB = c:$$

$$\text{shew that } CD = \frac{c \sin \alpha}{\sin (\alpha + \beta + \gamma)}.$$

4. In the ambiguous case if  $c, c'$  be the ambiguous sides and  $\theta$  the angle between the two positions of  $a$ , then

$$\tan \theta = \frac{(c^2 - c'^2) \sin 2a}{2cc' - (c^2 + c'^2) \cos 2a}.$$

5. In the ambiguous case ( $a, b, a$  given), if one triangle be  $n$  times the other, then

$$\frac{a}{b} = \frac{1}{n+1} \sqrt{n^2 + 1 - 2n \cos 2a}.$$

6.  $A, B, C, D$  are four trees in a straight row, such that  $AB, BC, CD$  subtend equal angles at a point  $P$ . If  $AB = 40$  ft.,  $BC = 20$  ft.,  $CD = 60$  ft., find  $PA, PB, PC, PD$ .

$$\text{Ans. } PA = 24 \sqrt{5} \text{ ft. } PB = 8 \sqrt{10} \text{ ft. } PC = 12 \sqrt{5} \text{ ft. } PD = 24 \sqrt{10} \text{ ft.}$$

7. In what direction must a road be carried up a hill, whose inclination to the horizon is  $\alpha$ , in order that the ascent may be 1 in  $n$ ? *Ans.* An angle  $\operatorname{cosec}^{-1}(n \sin \alpha)$  with the base of the hill.

8. Two towers (heights  $a, b$ ) stand on a horizontal plane at a distance  $c$  from each other: if  $\theta$  be the angle subtended by the tower ( $a$ ) at the top of the other, then  $\cot \theta = \frac{b^2 + c^2 - ab}{ac}$ .

9. A tower  $AB$  stands on an inclined plane; at a point  $C$  on the plane the tower is observed to subtend an angle  $\alpha$ ; and on proceeding to a point  $D$  in the line  $AC$ , so that  $CD = CA$ , it subtends an angle  $\beta$ . If  $\phi$  be the angle between the tower and the line  $AC$ , then  $\cot \phi = \cot \beta - 2 \cot \alpha$ .

10. A person in the valley between two hills observes the altitude  $\alpha$  of one of them. He then ascends the other through a vertical height  $a$ , and finds the angle of depression of his former station  $\beta$ , and the altitude of the hill  $(\alpha - \beta)$ . The height of the observed hill is  $a \frac{\sin^2 \alpha}{\sin^2 \beta}$ .

11. A tower situated on a horizontal plane leans towards the north: at two points due south and distant  $a, b$  respectively from the foot of the tower, the altitudes of the tower are observed to be

$\alpha, \beta$  respectively: the vertical height of the tower is  $\frac{b-a}{\cot \beta - \cot \alpha}$ ,  
and its inclination to the horizon is  $\tan^{-1} \frac{b-a}{b \cot \alpha - a \cot \beta}$ .

12. A quadrilateral figure is inscribed in a semicircle (diameter  $a$ ), and the sides drawn from the extremities of the diameter are inclined to each other at an angle  $\alpha$ . Shew that the side of the quadrilateral opposite to the diameter is equal to  $a \cos \alpha$ .

13. A person at  $A$  observes the elevation  $\alpha$  of the summit  $C$  of a mountain: he then proceeds up a slope (inclination  $\gamma$ ) directly towards the summit to  $B$ ; at  $B$  the elevation of the mountain is  $\beta$ : if  $AB = a$ , shew that the height of the mountain is

$$a \sin \alpha \sin (\beta - \gamma) \operatorname{cosec} (\beta - \alpha).$$

14. A person observes two telegraph posts in the same line, which is at right angles to the road on which he is walking. After walking  $a$  yards the nearer post is in a straight line with a church; after  $b$  more yards the further post and church are in the same line, and the two posts subtend the same angle as they did at his first station. If the distance of the nearer post from the first station is  $l$ , shew that the distance of the church is  $\frac{lab}{l^2 - 2a^2}$ .



## SECTION VI.

LOGARITHMS. THEIR USE. ADVANTAGE OF LOGARITHMS  
 BASE 10. ARRANGEMENT OF THE TABLES. LOGARITHMS  
 OF THE TRIGONOMETRICAL RATIOS.

51. We have established in the preceding sections certain relations between the sides and angles of triangles by means of which when we know certain parts of a triangle, we are enabled to calculate the rest: and our examples have illustrated the way in which these relations may be applied to calculate the heights and distances of objects which cannot be directly measured. In most practical cases these methods would involve tedious numerical calculations, and it becomes a matter of great importance to shorten them. This is done by means of what are called logarithms, as we shall now proceed to explain.

52. In the equation  $N = a^x$ , where  $N$ ,  $a$ ,  $x$  are any quantities satisfying this relation,  $a$  is called the *base*, and  $x$  the *logarithm* of the number  $N$  to the base  $a$ . This relation is also thus expressed,  $x = \log_a N$ .

Hence we obtain the following definition. The logarithm of a number to a given base, is the index of the power to which the base must be raised in order to be equal to the given number.

53. In the equation  $N = a^x$ , let  $a$  be some constant quantity greater than unity: then as  $x$  gradually increases,  $N$  gradually increases;

when  $x = 0$ ,  $N = 1$ ;    when  $x = 1$ ,  $N = a$ ;  
 when  $x = 2$ ,  $N = a^2$ ;    and when  $x = \infty$ ,  $N = \infty$ .

As  $x$  increases from 0 to 1,  $N$  increases from 1 to  $a$ ; and whatever value  $x$  may have between 0 and 1, there must be some corresponding value of  $N$  between 1 and  $a$ . And generally, whatever positive value be given to  $x$ , there must be some corresponding value of  $N$  between 1 and infinity. And if we give  $x$  all possible values between 0 and  $\infty$ , the corresponding values of  $N$  will comprise all possible values between 1 and  $\infty$ .

Again, if  $x$  receive all possible values from 0 to  $-\infty$ ,  $N$  must receive all fractional values between  $\frac{1}{a^0}$  and  $\frac{1}{a^\infty}$ , that is, between and 0.

Hence, by giving  $x$  all values from  $-\infty$  to  $+\infty$ ,  $N$  will have all values from 0 to  $\infty$ , that is, all positive values.

And conversely, whatever positive value  $N$  may have, there must be some real value of  $x$ , such that  $a^x = N$ .

If  $a$  be less than unity, let  $a = \frac{1}{b}$ , then  $a^x = b^{-x} = N$ ; and by giving  $x$  all values from  $+\infty$  to  $-\infty$ , we give  $b^{-x}$  all values from 0 to  $\infty$ , and therefore  $a^x$  or  $N$  all values from 0 to  $\infty$ .

If  $a = 1$ , all real powers of  $a$  are equal to unity, and we cannot make  $a^x = N$ .

If  $a$  is negative, the values of  $a^x$  are sometimes positive, sometimes negative, sometimes impossible, and so we cannot always make it equal to  $N$ . So also if  $N$  is negative we cannot always find  $x$ , such that  $a^x = N$ .

Hence we conclude, that if  $a$  is any positive quantity other than unity, we can always find a quantity  $N$  equal to  $a^x$  whatever real quantity  $x$  may be, and we can always find a value of  $x$ , such that  $a^x = N$ , whatever positive quantity  $N$  may be.

Now in order to make logarithms of any practical advantage, we must have the logarithms of all numbers up to a certain magnitude, calculated to a given base, arranged in tables; such an arrangement is called a table of logarithms, and the series of numbers and their corresponding logarithms is called a system of logarithms. Hence we see that the base of our system of logarithms must be positive, and the logarithms of negative numbers must be excluded.

We shall assume in the present treatise that the logarithms of all numbers to any positive base other than unity can be calculated to any degree of accuracy, and arranged in tables. The rest is all that falls within our province.

54. Let  $x, y$  be the logarithms of the same number  $N$  to the bases  $a, b$  respectively. Then

$$x = \log_a N, \quad y = \log_b N,$$

and we have

$$a^x = N = b^y;$$

$$\therefore a^{\frac{x}{y}} = b;$$

$$\therefore \frac{x}{y} = \log_a b;$$

$$\therefore \log_b N = \frac{1}{\log_a b} \cdot \log_a N.$$

Hence if we know the logarithm of any number  $N$  to base  $a$ , and also the logarithm of  $b$  to the same base  $a$ , we can find the logarithm of  $N$  to the base  $b$ , by multiplying  $\log_a N$  by the fraction  $\frac{1}{\log_a b}$ . This multiplier is called the *modulus* of the system whose base is  $b$  to the system whose base is  $a$ , because by means of it we can reduce a system of logarithms with base  $a$ , to a system with base  $b$ .

To prove that  $\log_a b \times \log_b a = 1$ .

Let  $x = \log_a b, \quad y = \log_b a,$   
 then  $a^x = b, \quad b^y = a,$   
 $\therefore a^{xy} = (a^x)^y = b^y = a,$   
 $\therefore xy = 1;$

or  $\log_a b \times \log_b a = 1,$

which was to be proved.

55. To find the logarithm of a product or quotient; of power or root.

Let  $x = \log_a M, \quad y = \log_a N,$   
 then  $M = a^x, \quad N = a^y,$

$$\therefore MN = a^x \cdot a^y = a^{x+y},$$

$$\therefore \log_a (MN) = x + y = \log_a M + \log_a N;$$

similarly  $\log_a (MNP \dots) = \log_a M + \log_a N + \log_a P + \dots$

Again, 
$$\frac{M}{N} = \frac{a^x}{a^y} = a^{x-y},$$

$$\therefore \log_a \left( \frac{M}{N} \right) = x - y = \log_a M - \log_a N.$$

Again, since 
$$M = a^x,$$

$$\therefore M^r = a^{rx},$$

$$\therefore \log_a M^r = rx = r \log_a M.$$

This is true whether  $r$  be whole or fractional,

$$\therefore \log_a \sqrt[r]{M} = \log_a M^{\frac{1}{r}} = \frac{1}{r} \log_a M.$$

It is on these properties that the great utility of logarithms depends—for the logarithm of the product of two numbers, we here see, is equal to the sum of the logarithms of the numbers, and so a process of multiplication is reduced to one of simple addition. And in the same way we facilitate the operations of division, involution, and evolution, by reducing them to subtraction, multiplication, and division respectively.

**Ex.** To find the value of  $\sqrt[16]{7985683}$ .

We find from the tables that

$$\log_{10} 7985683 = 6.9023122,$$

$$\therefore \log_{10} \sqrt[16]{7985683} = \frac{6.9023122}{16} = .4313945;$$

and from the tables we have

$$.4313945 = \log_{10} 2.700191,$$

$$\therefore \sqrt[16]{7985683} = 2.700191.$$

56. Let the logarithm of any number  $N$  to base 10 be  $n + d$ , where  $n$  is some integer, positive, or negative, or zero; and  $d$  a decimal fraction; then

$$\log_{10} N = n + d,$$

$$\text{and } \log_{10} 10^m N = \log_{10} 10^m + \log_{10} N = m + n + d.$$

**R T.**

**Q**

Hence we see that the logarithms of  $N$  and  $10^m N$  to base 10 have the same decimal part, whatever integer  $m$  is, positive or negative. And if we have a system of logarithms calculated to base 10, the same decimal part of the logarithm will correspond to all numbers which differ from each other only in the position of the decimal point, or which can be derived from each other by multiplying or dividing by any power of 10. Numbers of this kind are said to have the same *significant* digits. It is from this property that 10 is chosen as the base of the system of logarithms in common use; and for the future we shall speak of the logarithm of a number to the base 10, as simply the logarithm of the number, and we shall leave out the suffix in writing it; thus by  $\log N$  we shall mean  $\log_{10} N$ .

57. The integral part of a logarithm is called its *characteristic*, the decimal part its *mantissa*. We have shewn that all numbers having the same significant digits have the same mantissa, we now proceed to shew how the characteristic may be determined in any particular case by simple inspection.

Let  $N$  be a number which has one digit before the decimal point, that is to say, some number between 1 and 10: then  $\log N$  must lie between 0 and 1, and is therefore some decimal.

$$\text{Let} \qquad \qquad \qquad \log N = d,$$

then  $10^{n-1} N$  is a number which has  $n$  figures before the decimal point, and its logarithm is  $(n-1) + d$ . Hence we see that if a number has  $n$  figures before the decimal point, its characteristic is  $(n-1)$ .

Again,  $\frac{N}{10^{n+1}}$  has  $n$  cyphers after the decimal point, before the first significant digit, and its logarithm is  $-(n+1) + d$ . Hence, if there are  $n$  cyphers after the decimal point, the characteristic is  $-(n+1)$ .

This last case requires some explanation:

$$-(n+1) + d = -n - (1-d) = -n - d' = -(n+d'),$$

where  $d' = 1 - d$  and is some decimal. Here the logarithm is negative, and the integral part  $-n$ : also the mantissa  $d'$  of  $\log \frac{N}{10^{n+1}}$  is not the same as  $d$  the mantissa of  $\log N$ , which seems to contradict what was before observed: but it is found more convenient to consider all numbers as having logarithms with positive mantissæ, and it is with this convention that we say that  $\log N$  and  $\log \frac{N}{10^{n+1}}$  have the same mantissæ, and that the characteristic of the latter is  $-(n+1)$ . Without this convention we should be obliged to have two tables, one for numbers greater, and the other for numbers less than unity.

$$\text{Ex.} \quad \log 5.4952 = .7399835 ;$$

$$\therefore \log 5495.2 = 3.7399835,$$

$$\text{and} \quad \log .0054952 = \bar{3}.7399835,$$

the  $-$  sign placed above the 3 in the last case signifying that the characteristic alone is negative.

Thus  $\bar{3}.7399835$  and  $-3.7399835$  are different: in fact

$$\bar{3}.7399835 = -3 + .7399835 = -2.2600165.$$

58. In the tables in common use the mantissæ of the logarithms of all numbers from 1 to 100000, calculated to seven decimal places, are arranged in order. Thus when we have any number containing only five significant digits we find the mantissa of its logarithm directly from the tables, and prefix the characteristic by inspection; and if we have a logarithm, we can find the logarithm nearest to it in the tables, and so find the corresponding number correctly to five significant digits.

Numbers of more than five digits must be taken from the tables by the use of proportional parts, for an explanation of which the student is referred to Section IX. in this treatise. We will only observe that in determining the side of a triangle of about a mile in length to five digits, the error cannot exceed one inch; a degree of accuracy which is fully equal to that of the ordinary instruments for surveying.

59. Since the Trigonometrical Ratios of angles less than  $90^\circ$  are real positive quantities, they also can have logarithms, just in the same way as other quantities. The logarithms of the sines, cosines, tangents, cotangents, secants, cosecants of all angles are arranged in ordinary tables at intervals of one minute. In some tables they are given for every second. They are calculated to seven decimal places as for common numbers; but since they are mostly negative 10 is added to them, to make them positive: hence the tabular logarithm of any Trigonometrical Ratio of an angle is equal to its real logarithm increased by 10. Thus

$$L \sin \alpha = \log \sin \alpha + 10,$$

where  $L \sin \alpha$  stands for the tabular logarithm, and  $\log \sin \alpha$  for the real logarithm of  $\sin \alpha$ ; and so also for the other Ratios.

60. Also since  $\sin \alpha = \cos (90 - \alpha)$ , therefore

$$\log \sin \alpha = \log \cos (90 - \alpha).$$

Hence if we have the log-sines of all angles from  $0^\circ$  to  $45^\circ$ , we also have the log-cosines of all angles from  $90^\circ$  to  $45^\circ$ : and if we have the log-cosines of all angles from  $0^\circ$  to  $45^\circ$ , we have also the log-sines of all angles from  $90^\circ$  to  $45^\circ$ . Hence a complete table of log-sines and log-cosines from  $0^\circ$  to  $45^\circ$  is also a complete table to  $90^\circ$ . This is taken advantage of in the arrangement of the tables. The column headed sines and counted downwards from the top of the page is headed cosines at the bottom of the page and counted upwards, as may be best understood by inspecting the tables. The same arrangement applies to the tangents and cotangents, the secants and cosecants.

### EXAMPLES. (A).

1. Shew that  $\log 2 + \log 3 + \log 5 = \log 30$ .
2. Shew that  $\log 8 + \log 25 = 2 + \log 2$ .
3. Shew that  $\log \frac{144}{35} = 5 \log 2 + 2 \log 3 - \log 7 - 1$ .

4. Shew that  $\log \sin 30^\circ = -\log 2$ ;  $\log \cos 30^\circ = \frac{1}{2} \log 3 - \log 2$ ;  
 $\log \sin 45^\circ = -\frac{1}{2} \log 2$ .
5. What will be the characteristics of the logarithms of the following numbers, 2750, 27·564, 28325·7, ·000042?
6. Find the value of  $\sqrt[3]{30901\cdot7}$ . *Ans.* 7·90672.
7. Find the value of  $(3727593)^{\frac{2}{3}}$ . *Ans.* 425·1787.
8. Find the value of  $(401\cdot3116)^{\frac{2}{3}}$ , and verify the result by arithmetic. *Ans.* 11.
9. Find the volume of a cube whose edges are 7·39148 inches. *Ans.* 403·826 cubic inches.
10. Find the volume of a rectangular paralleliped whose edges are 4·23665, 7·39148, 3·652281 inches respectively. *Ans.* 114·3716 cubic inches.
11. How many figures will  $2^{40}$  contain? *Ans.* 13.
12. Given  $\log 2$  find  $\log \sqrt[14]{\frac{2^{60}}{5^{20}}}$ . *Ans.* ·2916.
13. What is the characteristic of 476 to base 8? *Ans.* 2.
14. What is the characteristic of ·0156 to base 3? *Ans.* -4.

EXAMPLES. (B).

1. Find the logarithms of  
309·017, ·0000309017, 309017000.
2. Find the logarithms of all numbers from 1 to 10.
3. Find the logarithms of the Trigonometrical Ratios of  $30^\circ$  and  $45^\circ$ .
4. Find the logarithms of 12,  $\sqrt{(45)}$ ,  $\frac{\sqrt{2}}{\sqrt[3]{3}}$ .  
*Ans.* 1·0791812, ·8266062, 1·9914746.
5. Find the value of  $\frac{\sqrt[3]{3}}{\sqrt[4]{4} \cdot \sqrt[5]{5}}$ . *Ans.* ·739148.



6. Find the value of  $\frac{3}{7} \left( \frac{5}{2} \right)^{\frac{1}{3}}$ . *Ans.* 1.69408.
7. Find the value of  $(.0035)^{\frac{1}{7}}$ . *Ans.* .445813
8. If  $20^\circ = 100$ , find  $x$ . *Ans.* 1.537...
9. If  $7^\circ = 2$ , find  $x$ . *Ans.* .356...
10. Find the value of  $\log \frac{1}{8}$  in terms of  $\log 25$ .  
*Ans.*  $\frac{3}{2} (\log 25 - 2)$ .
11. Given  $\log 2$ , find  $\log .00016$ ,  $\log (.000016)^{\frac{1}{5}}$ .  
*Ans.*  $\bar{4}.20412$ ,  $\bar{1}.4671244$ .
12. Find the number of cyphers between the decimal point and the first significant digit of  $\frac{1}{5^w}$ . *Ans.* 27
13. Find  $L \sin 36^\circ$ ,  $L \tan 18^\circ$ .  
*Ans.* 9.7692187, 9.511776

## EXAMPLES. (C).

1. Find the value of  $\left( \frac{1}{3} \right)^{\frac{187}{14}}$ . *Ans.* .00000042366
2. Find the value of  $\left( \frac{1}{100} \right)^{\frac{19}{7}}$ . *Ans.* .00000372759
3. Find the value of  $\sin 18^\circ$ ; and shew that the value found above (page 47) is equal to it.
4. If  $a^x = e^y$ , find the fraction  $\frac{x}{y}$ .
5. Shew that  $\log_a x : \log_b x = \log_c b : \log_c a$ .
6. Transform  $\sqrt[4]{3 \sin \alpha} = \frac{1}{5} \sqrt[4]{\left( \frac{\cos^2 \beta}{\sin \gamma} \right)}$ ,  
and  $4 \sqrt[4]{\left\{ \left( \frac{\sin \gamma}{\cos \beta} \right)^2 \right\}} = 3 \sqrt[4]{\left( \frac{\sin^2 \beta}{\cos^4 \gamma} \right)}$ ,  
into equations between the tabular logarithms of the quantities involved.

7. Assuming that  $\log 250$  and  $\log 256$  differ by  $\cdot 0103$ , shew that  $\log 2 = \cdot 30103$ .

8. Find the logarithm of 9 to base  $3\sqrt{3}$ , and of 125 to base  $\sqrt{5} \cdot \sqrt[3]{5}$ .

*Ans.*  $1\cdot\bar{3}$ ,  $3\cdot\bar{6}$ .

9. Given  $\log_{10} 2$ ,  $\log_{10} 3$ , find  $\log 16$  to base 15.

*Ans.*  $1\cdot 023832$ .

10. Given  $\log_{10} 2$ ,  $\log_{10} 3$ , find  $\log_8 3240$ .

*Ans.*  $5\cdot 02244$ .

11. Find  $\log \frac{10\sqrt[3]{\cdot 045}}{\sqrt[5]{720}}$ .

*Ans.*  $\bar{1}\cdot 9796043$ .

12. Given  $\log 2$ ,  $\log 3$ , find  $\log 10\cdot\bar{6}$ .

*Ans.*  $1\cdot 0280287$ .

13. Given  $\log 6 = a$ ,  $\log 15 = b$ , find  $\log 8$ ,  $\log 9$ .

*Ans.*  $\log 8 = \frac{3}{2}(a - b + 1)$ ,  $\log 9 = a + b - 1$ .

14. If a series of numbers are in G.P., shew that their logarithms are in A.P.

15. If  $x$ ,  $y$  are the logarithms of two numbers  $M$ ,  $N$ , shew that  $\log \sqrt{MN} = \frac{1}{2}(x + y)$ . Hence shew that  $1\cdot 5$  is the logarithm of  $31\cdot 622\dots$

16. Shew (from last question) that  $x + \frac{r(y-x)}{n}$  is the logarithm of  $M\left(\frac{N}{M}\right)^{\frac{r}{n}}$ .

17. If in last question  $N = M + 1$ , a number of 5 digits, and  $n = 10$ ,  $r < 10$ , shew that the equation  $M\left(\frac{N}{M}\right)^{\frac{r}{n}} = M + \frac{r}{10}$  is true to four decimal places.

18. If  $\log 48753 = 4\cdot 6880013$ ,  $\log 48754 = 4\cdot 6880103$ , shew that  $\log 487537 = 5\cdot 6880076$ .

19. Prove that if the bases of different systems of logarithms increase in G.P. their moduli to any base decrease in H.P.

## SECTION VII.

### APPLICATION OF LOGARITHMS TO THE SOLUTION OF TRIANGLES. ILLUSTRATION.

61. We have seen that when we know the logarithms of any numbers we can at once find the logarithm of their product and quotient, but that we cannot apply logarithms directly to the determination of the sum or difference of numbers. Hence in calculating any parts of a triangle from the others by logarithms we must use the formulæ which connect the parts by means of *factors* only; if we wish to use a formula involving *terms* we must first transform it into the form of a product.

Thus in calculating  $a$  from the formulæ

$$\cos a = \frac{b^2 + c^2 - a^2}{2bc},$$

we cannot apply logarithms directly, but if we transform this equation to

$$\cos \frac{a}{2} = \sqrt{\left( \frac{s \cdot s - a}{bc} \right)}$$

logarithms become immediately applicable.

We may observe by the way, that in using the first equation we should have to perform four operations in multiplication and one in division; in using the latter we should have to look out five logarithms. If  $a, b, c$  were simple numbers the first operation would probably give the least trouble, but as in practice they are generally quantities of five or six digits, the latter is much the shorter.

62. In most cases of solution of triangles the formulæ to be employed are in the form of products: the case which offers most difficulty is when two sides are given and the included angle; *this we accordingly proceed to discuss.*

Let  $a, b, \gamma$  be given; it is required to find the other angles  $\alpha, \beta$ , and the side  $c$ .

Since 
$$\tan \frac{\alpha - \beta}{2} = \frac{a - b}{a + b} \cot \frac{\gamma}{2}, \quad \text{Art. 47,}$$

we have

$$\log \tan \frac{\alpha - \beta}{2} = \log (a - b) - \log (a + b) + \log \cot \frac{\gamma}{2},$$

(supposing  $a > b$ ) which gives  $\frac{\alpha - \beta}{2}$ ;  $\frac{\alpha + \beta}{2}$  is already known, and therefore we can easily determine  $\alpha$  and  $\beta$ .

Having found  $\alpha$  and  $\beta$  we can find  $c$  from the equation

$$c = a \frac{\sin \gamma}{\sin \alpha}$$

or

$$\log c = \log a + \log \sin \gamma - \log \sin \alpha.$$

63. It sometimes happens that in solving this triangle we may know  $\log a, \log b$ , but not  $a$  and  $b$ , and it becomes an object to save the trouble of looking them out in the tables.

$$\tan \frac{\alpha - \beta}{2} = \frac{a - b}{a + b} \cot \frac{\gamma}{2}$$

$$= \frac{1 - \frac{b}{a}}{1 + \frac{b}{a}} \cot \frac{\gamma}{2}.$$

Let  $\frac{b}{a} = \tan \phi$ , an assumption which is always allowable, since the tangent of an angle may have any value;

then 
$$\tan \frac{\alpha - \beta}{2} = \frac{1 - \tan \phi}{1 + \tan \phi} \cot \frac{\gamma}{2}$$

$$= \tan (45^\circ - \phi) \cot \frac{\gamma}{2}. \quad \text{See Art. 35 bis, 3.}$$

Now

$$\log \tan \phi = \log b - \log a,$$

and is known at once, and therefore  $\phi$  may be found from the

Hence

$$45^\circ - \phi = 9^\circ. 19',$$

and

$$\begin{aligned} L \tan \frac{P-Q}{2} &= 9.2149894 + 10.8395431 - 10 \\ &= 10.0545325 \\ &= L \tan 48^\circ. 35'. \end{aligned}$$

Hence

$$\begin{aligned} P - Q &= 97^\circ. 10', \\ P + Q &= 163^\circ. 32'; \\ \therefore Q &= 33^\circ. 11'. \end{aligned}$$

Now

$$PQ = \frac{\sin \gamma}{\sin Q} x;$$

$$\begin{aligned} \therefore \log PQ &= L \sin \gamma - L \sin Q + \log x \\ &= 9.4524879 - 9.7382412 + 2.4311124 \\ &= 2.1453591 \\ &= \log 139.75. \end{aligned}$$

And therefore  $PQ = 139.75$  yards, the distance required.

## EXAMPLES. (A).

1. Adapt the formulæ for the solution of Case I, Art. 45, to tabular logarithms.
2. Adapt the formulæ for the solutions of Cases II, III, IV, to tabular logarithms.
3. If  $\alpha = 22$ ,  $b = 23$ ,  $c = 25$ , then  $\beta = 58^\circ. 10'. 43''$ .
4. If  $\alpha = 230$ ,  $b = 240$ ,  $c = 12$ , then  $\beta = 145^\circ. 37'. 35''$ .
5. If  $\alpha = 23$ ,  $\beta = 18^\circ$ ,  $\gamma = 23^\circ. 42'. 43''$ ,  
then  $b = 10.68185$ ,  $c = 13.9009$ .
6. If  $b = 159.0643$ ,  $\beta = 62^\circ. 6'. 51''$ ,  $\gamma = 53^\circ. 27'. 20''$ ,  
then  $\alpha = 162.335$ ,  $c = 144.58$ .
7. If  $\alpha - b = 4013.116$ ,  $\alpha + b = 7906.72$ ,  $\gamma = 36^\circ$ ,  
then  $\alpha = 129^\circ. 22'. 26''.8$ ,  $\beta = 14^\circ. 37'. 33''.2$ .

8. If  $a = 2820.9385$ ,  $b = 1430.8485$ ,  $\gamma = 157^\circ. 27'. 40''$ ,  
then  $\alpha = 14^\circ. 59'. 49''. 03$ ,  $\beta = 7^\circ. 32'. 30''. 97$ .
9. If  $\log a = 3.2978321$ ,  $\log b = 4.0135140$ ,  $\gamma = 36^\circ. 52'. 12''$ ,  
then  $\beta = 135^\circ. 21'. 29''. 59$ ,  $\alpha = 7^\circ. 46'. 18''. 41$ .
10. Find the value of  $\sqrt{(16233.5)^2 - (10681.85)^2}$  by means  
of a subsidiary angle. *Ans.* 12223.9.

EXAMPLES. (B).

1. If the sides of a triangle are 32, 40, 66, find the greatest  
angle. *Ans.*  $132^\circ. 34'. 34''$ .
2. The sides  $a$ ,  $b$ ,  $c$  of a triangle are in the ratio 4, 5, 6; find  
 $\beta$ . *Ans.*  $\beta = 55^\circ. 46'. 18''$ .
3. If  $a = 45^\circ$ ,  $a = 14000$ ,  $b = 15906.43$ , find the other angles.  
*Ans.*  $\beta = 53^\circ. 27'. 20''$ ,  $\gamma = 81^\circ. 32'. 40''$ ,  
 $\text{or } \beta = 126^\circ. 32'. 40''$ ,  $\gamma = 8^\circ. 27'. 20''$ .
4. Two sides of a triangle are to each other as 9 : 7, and the  
included angle is  $64^\circ. 12'$ ; find the other angles.  
*Ans.*  $69^\circ. 10'. 10''$ ,  $46'. 37'. 50''$ .
5. If  $a = 23^\circ. 42'. 43''$ ,  $\beta = 18^\circ$ ,  $a = 207$ , find  $b$ .  
*Ans.*  $b = 159.0643$ .
6. If  $a = 60^\circ$ ,  $\beta = 57^\circ. 53'. 9''$ ,  $c = 3727.593$ , find  $a$ .  
*Ans.*  $a = 3652.28$ .
7. If  $\log a$ ,  $\log b$  are given, find  $\log(a + b)$  by means of a  
subsidiary angle.

Find also  $\log(a - b)$ , where  $a > b$ .

8. Shew how to find the value of the following expressions by  
means of subsidiary angles :

$$\sqrt{(a^2 \pm b^2)}, \quad a \pm \sqrt{(a^2 - b^2)}, \quad \sqrt{(a + b)} \pm \sqrt{(a - b)}.$$

9. Solve by means of a subsidiary angle

$$\sin x - a \text{ vers } x = b.$$

$$\text{Ans. } \sin(x + \theta) = (a + b) \cos \theta, \text{ where } \tan \theta = a.$$

## EXAMPLES. (C).

1. If two sides of a triangle are 70, 35, feet respectively, and the included angle is  $36^{\circ}.52'.12''$ , find the remaining angles.

*Ans.*  $116^{\circ}.33'.54''$ ,  $26^{\circ}.33'.54''$ .

2. If the angles of a triangle be in arithmetical progression, and the greatest side is to the least in the ratio 5 : 4, find the angles.

*Ans.*  $70^{\circ}.53'.36''$ ,  $49^{\circ}.6'.24''$ .

3. One angle of a triangle is  $60^{\circ}$ , and the ratio of the side opposite to it to the difference of the sides containing it, is  $9\sqrt{3} : 2$ ; find the remaining angles.

*Ans.*  $66^{\circ}.22'.45''$ ,  $53^{\circ}.37'.15''$ .

4. If  $x^2 = a^2 + b^2 + ab$ , shew that subsidiary angles may be used to determine  $x$  from  $x = \sqrt{(ab)} \sec \phi$ , where

$$\tan \theta = \frac{b}{a},$$

and

$$\tan^2 \phi = 2 \operatorname{cosec} (2\theta).$$

5. If  $a, b, \gamma$  be given, shew that

$$\tan \beta = \frac{2b}{a-b} \tan \frac{\gamma}{2} \cos^2 \phi, \quad \tan \alpha = -\frac{2a \sec 2\phi}{a-b} \tan \frac{\gamma}{2} \cos^2 \phi,$$

where

$$\tan \phi = \sqrt{\frac{a+b}{a-b}} \tan \frac{\gamma}{2}.$$

6. If

$$\sin \theta \sqrt{1 + \tan^2 \alpha \tan^2 \beta} + \cos \theta \sqrt{1 - \tan^2 \alpha \tan^2 \beta} = \tan \alpha + \tan \beta,$$

find  $\theta$  in a form adapted to logarithmic computation.

*Ans.*  $\sin (\theta + \phi) = \frac{\sin (\alpha + \beta)}{\sqrt{2} \cos \alpha \cos \beta}$ , where  $\cos 2\phi = \tan^2 \alpha \tan^2 \beta$ .

## SECTION VIII.


### CIRCULAR MEASURE. POLYGONS INSCRIBED, AND CIRCUM- SCRIBED ABOUT A CIRCLE. AREA OF A CIRCLE.

65. **THERE** is another system of measuring angles besides that by degrees or grades, which on account of its importance in all the higher branches of mathematics, it will be necessary to consider in this treatise.

We must premise the following propositions.

The ratio of the circumference of a circle to its diameter is the same for all circles.

To prove this, we shall assume, what is proved by Newton and appears almost self-evident, that if in a curved figure a polygon be inscribed, and the number of sides of the polygon be increased and the magnitude of each side diminished without limit, the perimeter of the polygon becomes more and more nearly equal to that of the curvilinear figure, and is ultimately equal to it.



Let  $ABCD$ ,  $abcd$  be two circles whose centres are  $O, o$ : and in  $ABCD$  let a polygon  $ABCD\dots$  be inscribed; in  $abcd$  let a similar polygon  $abcd\dots$  be inscribed, then we have

$$AB : AO = ab : ao,$$

$$BC : AO = bc : ao,$$

$$CD : AO = cd : ao,$$

$$\text{And so on.}$$



Hence  $AB + BC + CD + \dots : AO = ab + bc + cd + \dots : ao$ , or  
 Perimeter of polygon  $ABCD \dots : AO = \text{Per. of pol. } abcd : ao$ . Now  
 this proportion is true however many sides the polygons have; let  
 their number be increased, and the magnitudes of each of them  
 diminished indefinitely; then the perimeters of the polygons be-  
 come the circumferences of the circles, and we have

$$\text{circumference } ABCD : AO = \text{circum. } abcd : ao,$$

or the ratio of the circumference of a circle to its diameter is the  
 same for all circles.

This ratio is generally denoted by the symbol  $\pi$ ; so that if  $r$   
 be the radius of a circle, and  $2r$  its diameter, its circumference is  
 $2\pi r$ . The value of  $\pi$  has been calculated, and is found to be  
 $3.14159\dots$  or the fraction  $\frac{22}{7}$  nearly.

66. We must next prove that the angle subtended at the  
 centre of any circle by an arc equal to the radius of the circle is of  
 invariable magnitude.

Let  $AOP$  (fig. Art. 7) be an angle at the centre of the circle  
 $ABA'B'$ , such that the arc  $AP$  is equal to the radius  $AO$ : then,  
 since angles at the centre of a circle are as the arcs on which they  
 stand (Euclid, VI. 33),

$$\frac{\text{angle } AOP}{4 \text{ right angles}} = \frac{\text{arc } AP}{\text{circumference } ABA'B'} = \frac{r}{2\pi r},$$

(since arc  $AP$  = radius, and circumference =  $2\pi r$ ),

$$\text{or} \quad \text{angle } AOP = \frac{4 \text{ right angles}}{2\pi} = \frac{2 \text{ right angles}}{\pi}.$$

Now  $\pi$  being invariable, and this expression independent of  $r$ , the  
 angle  $AOP$  is the same whatever be the size of the circle.

Hence it appears that this angle is a proper standard by which  
 to measure other angles. If this angle be taken for unity, any  
 other angle which is  $\theta$  times as great will be represented by  $\theta$ .  
 $\theta$  is called the circular measure of the angle.

67. To shew that  $\theta$ , the circular measure of an angle, is equal  
 to the arc subtended by that angle, divided by the radius.

Let  $\angle AOP$  be the angle whose arc  $AP$  = radius  $AO$ ,  $\angle AOP$ , any other angle whose circular measure is  $\theta$ , then

$$\theta = \frac{\text{angle } AOP}{\text{angle } AOP} = \frac{\text{arc } AP}{\text{arc } AP} = \frac{\text{arc}}{\text{radius}}.$$

Since the arc subtended by four right angles is  $2\pi r$ , the circular measure of four right angles is  $\frac{2\pi r}{r} = 2\pi$ . Hence the measure of a right angle is  $\frac{\pi}{2}$

68. To pass from circular measure to degrees and grades and vice versa, we have, as in Art. 5,

$$\frac{D}{90} = \frac{G}{100} = \frac{2\theta}{\pi}.$$


To find the number of degrees in the unit of circular measure, put  $\theta = 1$  in this equation, then we have

$$D = \frac{180^\circ}{\pi} = \frac{180^\circ}{3.14159} = 57^\circ.29577.$$

69. It is usual to represent angles when degrees are used by the Roman letters A, B, C, ..., when circular measure is used, by the Greek letters  $\alpha, \beta, \gamma, \dots, \theta, \phi, \dots$ . We have used the latter throughout this treatise, even when the student had no idea of circular measure. Our reasons for so doing are mentioned in the preface: whilst we are merely dealing with the relations amongst the Trigonometrical Ratios of various angles, it is of no consequence what measure we employ; but when we trace relations between the angles themselves and their Trigonometrical Ratios, we must always use circular measure as in the following article.

70. To shew that  $\frac{\sin \theta}{\theta}, \frac{\tan \theta}{\theta}$  are equal to unity, when  $\theta = 0$ .

Let  $\angle AOB = \angle AOB' = \theta$ , and draw  $BCB'$  perpendicular to  $BD$  and let  $BAB'$  be the arc of a circle whose centre is  $O$ : draw  $B'D$  to touch this arc at  $B$ ,  $B'$ .



Then

$$\theta = \frac{1}{2} \frac{\widehat{BAB'}}{OA} = \frac{\widehat{BA}}{OA},$$

$$\tan \theta = \frac{BD}{OB} = \frac{BD}{OA},$$

$$\sin \theta = \frac{BC}{OB} = \frac{BC}{OA}.$$

Hence, since

$$BD + DB' > \text{arc } BAB',$$

and

$$BB' < \text{arc } BAB',$$

we have

$$\tan \theta > \theta,$$

and

$$\sin \theta < \theta;$$

Hence  $\tan \theta$ ,  $\theta$ ,  $\sin \theta$  are in descending order of magnitude; hence

$$\frac{\sin \theta}{\tan \theta}, \frac{\sin \theta}{\theta}, \frac{\sin \theta}{\sin \theta},$$

or

$$\cos \theta, \frac{\sin \theta}{\theta}, 1$$


are in ascending order of magnitude. But when  $\theta = 0$ ,  $\cos \theta = 1$

therefore also  $\frac{\sin \theta}{\theta}$  must in this case equal unity.

Also

$$\frac{\tan \theta}{\theta} = \frac{1}{\cos \theta} \cdot \frac{\sin \theta}{\theta} = 1 \text{ when } \theta = 0.$$

71. To find the perimeters and areas of regular polygons circumscribed about, and inscribed in a given circle.



Let  $AB$ ,  $ab$  be sides of regular polygons of  $n$  sides circumscribed about and inscribed in a circle whose centre is  $O$  and radius  $r$ .

Then

$$\angle AOB = \angle aOb = \frac{2\pi}{n},$$

$$\therefore \angle aOb = \frac{\pi}{n}.$$

Then

$$AB = 2aB = 2r \tan \frac{\pi}{n};$$

perimeter of circumscribed polygon  $= n \cdot AB$

$$= 2nr \tan \frac{\pi}{n};$$

area of circumscribed polygon

$$= n \cdot aB \cdot Oa = nr^2 \tan \frac{\pi}{n}.$$

Again, perimeter of the inscribed polygon  $= 2nr \sin \frac{\pi}{n}.$

Area of inscribed polygon  $= n \cdot \Delta aOb = n \cdot \frac{1}{2} aO \cdot bO \sin aOb$

$$= \frac{1}{2} nr^2 \sin \frac{2\pi}{n}.$$

72. *To find the area of a circle.*

This equals the area of the inscribed polygon when the number of sides is made infinite.

Let  $\frac{2\pi}{n} = \theta$ , then  $n = \frac{2\pi}{\theta}$ , and when  $n = \infty$ ,  $\theta = 0$ .

The area of polygon becomes

$$\frac{\pi}{\theta} r^2 \sin \theta = \pi r^2 \frac{\sin \theta}{\theta},$$

when  $\theta = 0$ ,  $\frac{\sin \theta}{\theta} = 1$ ;

hence, the area of the circle  $= \pi r^2$ .

The volume of the cylinder is the area of its base, multiplied by its height.

73. Referring the student to Art. 22, we will investigate a general expression for all the angles which satisfy the relation

$$\sin \theta = a.$$

Let  $AOP_1$  (fig. Art. 7) or  $a$  be the smallest angle whose sine is equal to  $a$ , then all the values of  $\theta$  are bounded either by  $OP_1$ , or  $OP_2$ , that is,  $\theta$  is equal to  $a$ , or  $(\pi - a)$  taken positively, or these angles increased or diminished by any multiple of  $2\pi$ ; or again,  $\theta$  is equal to  $-(\pi + a)$ ,  $-(2\pi - a)$  taken negatively, or these angles increased or diminished by any multiple of  $2\pi$ . Hence all the values of  $\theta$  are included in the forms  $2n\pi + a$ ,  $2n\pi + (\pi - a)$ ,  $2n\pi - (\pi + a)$ ,  $2n\pi - (2\pi - a)$ , where  $n$  is any integer positive or negative.

Now when the coefficient of  $\pi$  in these forms is even (as in the first and fourth),  $a$  has the positive sign, and when odd (as in the second and third) the negative sign before it. Hence the form  $m\pi + (-1)^m a$ , where  $m$  is any integer positive or negative, includes all these forms, since when  $m$  is even  $(-1)^m$  is positive, and when  $m$  is odd  $(-1)^m$  is negative; hence  $\theta = m\pi + (-1)^m a$ .

74. *To find all the values of  $\theta$  which satisfy the equation*

$$\cos \theta = a.$$

Let  $\angle OP_1$  or  $a$  be the smallest of these values. Then all the values of  $\theta$  must be bounded either by  $OP_1$  or  $OP_2$ , that is, in the positive direction they will be  $a$  and  $(2\pi - a)$ , and these angles increased or diminished by any multiple of  $2\pi$ ; and in the negative direction they will be  $-a$ , and  $-(2\pi - a)$ , and these angles increased or diminished by any multiple of  $2\pi$ . Hence all the values of  $\theta$  are included in the forms  $2n\pi + a$ ,  $2n\pi + (2\pi - a)$ ,  $2n\pi - a$ ,  $2n\pi - (2\pi - a)$ . Observing in these forms that the coefficients of  $\pi$  are all even, whilst  $a$  is either positive or negative, we find as the general value of  $\theta$ , that  $\theta = 2m\pi \pm a$ , when  $m$  is any integer positive or negative.

75. To find all the values of  $\theta$  which satisfy the equation

$$\tan \theta = a.$$

In the same way as in the last two Articles, we may shew that all the values in the positive direction are included in the forms  $n\pi + a$ ,  $2n\pi + (\pi + a)$ , and those in the negative direction in the forms  $2n\pi - (\pi - a)$ ,  $2n\pi - (2\pi - a)$ ; since in these forms  $a$  is always positive, all the values of  $\theta$  are included in the form

$$\theta = m\pi + a.$$

It is not necessary in these cases that  $a$  should be the *smallest* value of  $\theta$ .

76. By the help of these forms we may explain the double value of  $\sin \frac{\theta}{2}$  when expressed in terms of  $\cos \theta$  (see Art. 34).

Let  $a$  be one value of  $\theta$  which satisfies the equation,  $\cos \theta = a$ ,

$$\text{then } \theta = 2n\pi \pm a, \text{ and } \frac{\theta}{2} = n\pi \pm \frac{a}{2}.$$

Now if the value found for  $\frac{\theta}{2}$  is as general as the value of  $\theta$  or  $\cos^{-1}a$  from which it is derived,  $\sin \frac{\theta}{2}$  must contain all the values of  $\sin \left( n\pi \pm \frac{a}{2} \right)$ .

Now  $\sin\left(n\pi \pm \frac{a}{2}\right) = \sin n\pi \cos \frac{a}{2} \pm \cos n\pi \sin \frac{a}{2} = \pm \sin \frac{a}{2}$ , (since  $\sin n\pi = 0$ ;  $\cos n\pi = +1$  or  $-1$ , as  $n$  is even or odd).

Hence  $\sin \frac{\theta}{2}$  must have two values, equal and of opposite sign.

Similarly we may shew that  $\cos \frac{\theta}{2}$  when expressed in terms of  $\sin \frac{\theta}{2}$  will have two values,  $\pm \cos \frac{a}{2}$ .

77. The four values of  $\sin \frac{\theta}{2}$  when expressed in terms of  $\sin \frac{a}{2}$  (Art. 33) may be thus explained.

Let  $a$  be one value of  $\theta$  which satisfies the equation,  $\sin \theta = \sin a$ ; then  $\theta = n\pi + (-1)^n a$ , and  $\frac{\theta}{2} = \frac{n\pi}{2} + (-1)^n \frac{a}{2}$ .

$$\text{Now } \sin\left(\frac{n\pi}{2} + (-1)^n \frac{a}{2}\right) = \sin \frac{n\pi}{2} \cos (-1)^n \frac{a}{2} + \cos \frac{n\pi}{2} \sin (-1)^n \frac{a}{2}$$

If  $n$  be even,  $\sin \frac{n\pi}{2} = 0$ ,  $\cos \frac{n\pi}{2} = \pm 1$ ,  $\sin (-1)^n \frac{a}{2} = \sin \frac{a}{2}$ , and therefore  $\sin \frac{\theta}{2} = \pm \sin \frac{a}{2}$ .

If  $n$  be odd,  $\sin \frac{n\pi}{2} = \pm 1$ ,  $\cos (-1)^n \frac{a}{2} = \cos \frac{a}{2}$ ,  $\cos \frac{n\pi}{2} = 0$ , and therefore  $\sin \frac{\theta}{2} = \pm \cos \frac{a}{2}$ .

Hence  $\sin \frac{\theta}{2}$  will have four values, two and two of opposite sign.

$$\begin{aligned} \text{Similarly } \cos \frac{\theta}{2} &= \cos \frac{n\pi}{2} \cos (-1)^n \frac{a}{2} - \sin \frac{n\pi}{2} \sin (-1)^n \frac{a}{2} \\ &= \pm \cos \frac{a}{2}, \text{ when } n \text{ is even,} \\ &= \pm \sin \frac{a}{2}, \text{ when } n \text{ is odd.} \end{aligned}$$

Similar generalizations apply to all the formulæ proved in Chapters II. and III.

78. Example.  $\sin \frac{\theta}{3}$  will have three values when expressed in terms of  $\sin \theta$  from the cubic equation

$$4 \sin^3 \frac{\theta}{3} - 3 \sin \frac{\theta}{3} + \sin \theta = 0, \text{ (see Art. 32).}$$

This will also appear as follows :

$$\sin \frac{\theta}{3} \text{ must contain all the values of } \sin \left( \frac{n\pi}{3} + (-1)^n \frac{\alpha}{3} \right).$$

Now  $n$  is of the form  $3m$  or  $3m \pm 1$ .

Let  $n = 3m$ , then

$$\sin \frac{\theta}{3} = \sin \left( m\pi + (-1)^{2m} \frac{\alpha}{3} \right) = \cos m\pi \sin (-1)^{2m} \frac{\alpha}{3} = \sin \frac{\alpha}{3},$$

whether  $m$  be even or odd.

Let  $n = 3m \pm 1$ , then

$$\begin{aligned} \sin \frac{\theta}{3} &= \sin \left( m\pi \pm \frac{\pi}{3} + (-1)^{2m \pm 1} \frac{\alpha}{3} \right) \\ &= \cos m\pi \sin \left( \pm \frac{\pi}{3} + (-1)^{2m \pm 1} \frac{\alpha}{3} \right) = \sin \left( \pm \frac{\pi}{3} - \frac{\alpha}{3} \right), \end{aligned}$$

whether  $m$  be even or odd.

Hence  $\sin \frac{\theta}{3}$  has 3 values,

$$\sin \frac{\alpha}{3}, \sin \left( \frac{\pi}{3} - \frac{\alpha}{3} \right), -\sin \left( \frac{\pi}{3} + \frac{\alpha}{3} \right).$$

From the form of the equation for determining  $\sin \frac{\theta}{3}$ , we see that the sum of these values must equal zero, and their product  $\sin \alpha$ ; which results may be easily verified.



## EXAMPLES. (A).

In the following examples the value of  $\pi$  is supposed to be  $\frac{22}{7}$ .

1. Find the number of degrees in an angle of a regular hexagon. *Ans.* 120

2. Find the number of degrees in an angle of a regular pentagon. *Ans.* 108

3. Find the number of degrees in an angle of a polygon of  $n$  sides.

$$\text{Ans. } \frac{n-2}{n} \cdot 180$$

4. How many sides has a polygon, whose angles are  $135^\circ$ ?

*Ans.*

5. If the area of the circumscribed polygon equals four times the area of the inscribed polygon, find the number of sides.

*Ans.*

6. Find the circular measure of  $35^\circ$ ,  $35'$ . *Ans.* .61, .5

7. Find the degrees &c. in an angle whose circular measure is 1.21. *Ans.*  $69^\circ.11'$

8. The number of sides of one polygon is four times that of another, and the angles of the first one-half as large again as those of the second. How many sides have they? *Ans.* 20,

9. Find the number of grades in the unit of circular measure

*Ans.*  $63^\circ.66'$

10. A tree is 12 feet round, what is the width of the largest plank which can be cut from it? *Ans.* 3 feet  $9\frac{1}{2}$  inches

11. A tree is 18 feet in circumference, what is the thickness of the greatest square plank which can be cut from it?

*Ans.* 4 feet nearly

12. A circle of three inches diameter is cut out of a circle of 5 inches diameter; shew that the area of the remainder is equal to a circle of 4 inches diameter.

13. How many cubic feet of timber are there in a tree 16 feet in circumference and 22 feet in height? *Ans.* 448.

14. A round tower is 42 feet high: its wall is 3 feet thick, and its interior 24 feet in diameter. How many bricks 9 inches long,  $4\frac{1}{2}$  inches wide, and 3 inches thick, were required to build it? *Ans.* 152064.

15. Three equal circles, radii  $r$ , are placed in contact; find the area between them. *Ans.*  $\left(\sqrt{3} - \frac{\pi}{2}\right)r^2$ .

Find the general values of  $\theta$  in the equations

16.  $\cos 7\theta + \cos \theta = 0$ . *Ans.*  $\theta = \frac{2n+1}{8}\pi$ , or  $\frac{2n+1}{6}\pi$ .

17.  $\sin 7\theta - \sin \theta = \sin 3\theta$ . *Ans.*  $\theta = \frac{n\pi}{3}$ , or  $\frac{n\pi}{2} \pm \frac{\pi}{12}$ .

18.  $\sin \theta = \cos n\theta$ . *Ans.*  $\theta = \frac{2m\pi \pm \frac{\pi}{2}}{n \pm 1}$ .

19.  $\cos \theta + \cos 3\theta = \frac{1}{2}$ .  
*Ans.*  $\theta = 2\left(n\pi \pm \frac{\pi}{3}\right)$ ,  $2n\pi \pm \frac{\pi}{5}$ ,  $2n\pi \pm \frac{3\pi}{5}$ .

20.  $\tan 3\theta \pm \tan \theta = 0$ . *Ans.*  $\theta = n\pi$  or  $n\pi \pm \frac{\pi}{4}$ .

21.  $\tan^2 \theta = a^2$ . *Ans.*  $\theta = n\pi \pm \tan^{-1} a$ .

22.  $\sin^2 2\theta - \sin^2 \theta = \sin^2 \frac{\pi}{6}$ . *Ans.*  $\theta = \left(n \pm \frac{1}{5} \pm \frac{1}{10}\right)\pi$ .

23.  $\sin^2 \theta \frac{\cos 3\theta}{3} + \cos^2 \theta \frac{\sin 3\theta}{3} = \frac{m}{4}$ .  
*Ans.*  $\theta = \frac{1}{4}\{n\pi + (-1)^n \sin^{-1} m\}$ .

## EXAMPLES. (B).

1. Find the circular measure of  $\pi$  grades. *Ans.*  $\frac{\pi^2}{200}$
2. At what distance from the eye will a shilling whose diameter is .91 inches exactly hide the moon whose diameter is  $30' 12''$ ? *Ans.*  $\frac{4095}{151\pi}$  feet
3. The radius of a circle is  $r$ ; find the length of an arc which subtends  $A^\circ$  at the centre. *Ans.*  $\frac{\pi r A}{180}$
4. The radius of the Earth being 4000 miles, what is the length of  $1^\circ$  of the meridian? *Ans.*  $69\frac{2}{3}$  miles
5. How far does a person at the equator travel in a second by reason of the Earth's rotation about its axis? *Ans.* 512 yards nearly
6. The diameter of the Earth's orbit about the Sun being 192,000,000 miles, how far in space does the Earth travel in one second? *Ans.* 19 miles nearly
7. The greatest square possible is cut out of a circle whose radius is  $r$ , the area of the remainder is  $(\pi - 2)r^2$ .
8. The area of a square inscribed in a circle : the area of an equilateral triangle inscribed in the same circle, as  $8 : 3\sqrt{3}$ .
9. If  $a_1, a_2, a_3$  be respectively the sides of a regular pentagon, hexagon, and decagon inscribed in a circle, then
 
$$a_1^2 = a_2^2 + a_3^2.$$
10. The area of inscribed polygon of  $m$  sides : area of circumscribed polygon of  $n$  sides  $= m \cos \frac{\pi}{m} : n$ , if the sides of the two polygons are equal.

11. A circular sector whose angle is  $\theta$  is formed into a right cone. Shew that the vertical angle of the cone is  $\sin^{-1} \frac{\theta}{2\pi}$ .

12. A regular hexagon is inscribed in a circle (radius  $r$ ), and its angular points are joined two and two so as to form another hexagon; shew that the area of this latter is  $\frac{r^2\sqrt{3}}{2}$ .

13. In a sector of a circle (angle  $2\alpha$ , radius  $r$ ) an equilateral triangle is inscribed so that one angle is at the middle point of the arc of the sector; shew that the sides of the triangle are each equal to  $\frac{2r}{\cot \alpha + \sqrt{3}}$ .

14. If  $2(\sin 2\theta + \sin 2\phi) = 1 = 2\sin(\theta + \phi)$ , find  $\theta$  and  $\phi$ .

$$\text{Ans. } \theta = \frac{n\pi}{2} + (-1)^n \frac{\pi}{12} \pm \frac{\pi}{6}, \quad \phi = \frac{n\pi}{2} + (-1)^n \frac{\pi}{12} \pm \frac{\pi}{6}.$$

15. If  $\sin \theta + \cos \theta < \frac{1+\sqrt{3}}{2}$ , find the limits of  $\theta$ .

$$\text{Ans. } \theta \text{ must not lie between } 2n\pi + \frac{\pi}{6} \text{ and } 2n\pi + \frac{\pi}{3}.$$

16. If  $\tan(\pi \cot \theta) = \cot(\pi \tan \theta)$ , then

$$\tan \theta = \frac{2n+1 \pm \sqrt{4n^2+4n-15}}{4},$$

where  $n$  is any integer positive or negative.

17. If  $\tan \theta + \tan m\theta + \tan n\theta = \tan \theta \tan m\theta \tan n\theta$ , then  $\theta = \frac{r\pi}{m+n+1}$ , where  $r$  is any integer.

18. Find the least values of  $\phi$  and  $\theta$  which satisfy the equations

$$\cos(\theta + 3\phi) = \sin(2\theta + \phi), \quad \sin(\phi + 3\theta) = \cos 2(\theta + \phi).$$

$$\text{Ans. } \theta = \phi = \frac{\pi}{16}.$$

19. If  $\sin^2 \theta \tan \theta + \cos^2 \theta \cot \theta = \frac{5}{4} \operatorname{cosec} 2\theta$ , then

$$\theta = (3n \pm 1) \frac{\pi}{6}.$$

20.  $\tan (n \cot \theta) = \cot (n \tan \theta)$ , then

$$\theta = \frac{m\pi}{2} + (-1)^m \frac{2n}{(2r+1)\pi},$$

where  $m, r$  are any integers.

### EXAMPLES. (C).

1. Find the length of an arc subtending an angle of  $60^\circ$  in a circle whose radius is 105 feet. *Ans.* 110 feet

2. If the length of an arc of  $60^\circ$  is 11 feet, then the radius of the circle is 10 feet 6 inches.

3. If a right angle were divided into 80 parts, and each of these into 80, find the number of the latter parts contained in an angle whose circular measure is .001. *Ans.*  $4\frac{1}{8}$

4. Shew that  $\frac{\operatorname{vers} \theta}{\theta \sin \theta} = \frac{1}{2}$ , when  $\theta = 0$ .

5. Shew that  $\frac{\sin n\theta}{\sin m\theta} = \frac{n}{m}$ , when  $\theta = 0$ .

6. A circular flat object at distance  $a$  subtends an angle of  $30'$ , its area is  $\frac{\pi^2 a^2}{(720)^2}$  very nearly.

7. If  $\alpha$  be the angle whose arc is  $r_1$  and radius  $r_1$ , and  $\alpha$  the angle whose arc is  $r_2$  and radius  $r_2$ , then

$$\frac{\alpha}{\alpha'} = \left( \frac{r_1}{r_2} \right)^2.$$

8. If  $R, r$  be radii of circles circumscribed about and inscribed in a regular polygon whose side is  $2a$ , then

$$R^2 - r^2 = a^2.$$

9. The Earth's radius being  $R$  and the height of a tower  $h$ ; find how far an object at sea must be from the foot of the tower so as just to be visible from the top.

$$\text{Ans. } R \cos^{-1} \frac{R}{R+h}.$$

If  $h$  be so small that  $\frac{h^2}{R^2}$  may be neglected, shew that this answer becomes  $\sqrt{2hR}$ .

10. The height of one ship is  $n$  times that of another, and the top of one is just visible from the masthead of the other at the distance of  $a$  miles. Given  $R$  the Earth's radius, shew that the height of one is given by the equation

$$\cos^{-1} \frac{R}{R+x} + \cos^{-1} \frac{R}{R+nx} = \frac{a}{R}.$$

11. The radius of the circle of latitude  $60^\circ$  is 2000 miles.

12. What distance is travelled in half an hour by a person situated in latitude  $60^\circ$ , by reason of the Earth's rotation?

$$\text{Ans. } 261\frac{1}{2} \text{ miles.}$$

13. If two plumb-lines suspended from points at a given distance apart on the Earth's surface are inclined at an angle of  $m''$ , and at  $n''$  when at an elevation  $h$ , then the radius of the Earth is  $\frac{nh}{m-n}$  very nearly.

14. A lighthouse 60 feet high is just seen from the deck of a ship 12 feet above the water, how far is the ship from the lighthouse?

$$\text{Ans. } 13.79 \text{ miles.}$$

If the height of the deck be raised 1 inch, how much further off may the lighthouse be seen?

$$\text{Ans. } .0148 \text{ miles.}$$

15. An observer from the deck of a ship 20 feet above the sea can just see the top of a lighthouse: at the masthead (80 feet above the sea) he can just see the door of the lighthouse, which is  $\frac{1}{4}$ th of the height of the top. Find the height of the lighthouse.

$$\text{Ans. } 80 \text{ feet.}$$

16. A person ascending in a balloon observes the dip of horizon  $\alpha$ ,  $\alpha'$  at successive heights  $h$ ,  $h'$ . Shew that

$$\frac{h}{h'} = \frac{(1 - \cos \alpha) \cos \alpha'}{(1 - \cos \alpha') \cos \alpha}.$$

17. Find the circular measure of an angle whose complement contains as many degrees, as the supplement of an angle  $n$  times as large contains grades.

$$\text{Ans. } \frac{11}{16} \pi$$

18. The centres of two wheels (radii  $r$ ,  $r'$ ) are at a distance  $a$  from each other: find the length of a strap passing round them both but not crossing between them.

$$\text{Ans. } \pi(r + r') + 2(r - r') \sin^{-1} \frac{r - r'}{a} + 2\sqrt{a^2 - (r - r')^2}.$$

19. Shew that  $\cos \frac{x}{2} \cos \frac{x}{2^2} \cos \frac{x}{2^3} \dots \dots \cos \frac{x}{2^n}$  is equal to  $\frac{\sin x}{x}$ , when  $n$  becomes infinite.

20. If with the angular points of a regular polygon of  $n$  sides (length of side  $a$ ) as centres, circles are drawn so that the points where the circles meet the sides being joined form a regular polygon of  $2n$  sides, shew that the radii of the circles must be

$$\frac{a}{4} \sec^2 \frac{\pi}{2n} \text{ or } \frac{a}{4} \left( 1 + 2 \cos \frac{\pi}{n} \right) \sec^2 \frac{\pi}{2n}.$$

21. In last question prove that area of new polygon : area of original polygon =  $\cos^2 \frac{\pi}{n} : \cos^2 \frac{\pi}{2n}$ .

22. In an equilateral triangle two hexagons are inscribed, one having three of its sides coinciding with sides of the triangle, and the other three of its angles bisecting the sides of the triangle. Shew that their areas are as 4 : 3.

23. Find the three values of  $\cos \frac{\theta}{3}$  when determined from the equation  $\cos \theta = a$ .

24. Find the three values of  $\tan \frac{\theta}{3}$  when determined from the equation  $\tan \theta = a$ , and shew that the sum of the three values found together with three times their product is equal to zero.

## SECTION IX.

## ON THE USE OF PROPORTIONAL PARTS IN LOGARITHMIC AND TRIGONOMETRICAL TABLES.

79. We have shewn in the preceding Sections VI. and VII. how logarithms are used for shortening calculations involving lengthy products and powers : but we were only able to use the Tables of logarithms with exactness, when the numbers or angles were found in the Tables, and in other cases we could only extract the nearest numbers or angles. The Tables generally give the logarithms of all numbers of five digits, and of all Trigonometrical Ratios of angles for every minute of the quadrant. We proceed in this section to shew how the Tables can almost always be applied with exactness to numbers of seven digits ; and to angles within the hundredth part of a second, by means of the principle of proportional parts.

80. First, as regards the logarithms of numbers. If we open the Tables at random, we find, for instance, that the logarithms of all the numbers from 26629 to 26673 increase uniformly by  $\cdot 0000163$ , from 26674 to 26794 increase by  $\cdot 0000163$  or  $\cdot 0000162$ , from 26795 to 26822 uniformly by  $\cdot 0000162$  ; and generally the logarithms of successive numbers increase gradually, nearly uniformly, with a difference which slowly diminishes as the numbers increase. Now this being true for the numbers found in the Tables, it will be equally true if we insert 100 numbers between two consecutive numbers. Thus we find by the Tables

$$\log 2662900 = 6\cdot 4253549,$$

and

$$\log 2663000 = 6\cdot 4253712,$$

the difference between the two logarithms being  $\cdot 0000163$ . Now if we take the 100 numbers between 2662900 and 2663000, we may conclude that the difference between each successive number is  $\cdot 000000163$  ; and if we wish to find the logarithm of any intermediate number, as  $\log 2662937$ , we must multiply  $\cdot 000000163$



by 37, and add the result to 6·4253549 the logarithm of 2662900,

$$\begin{aligned} \text{i.e.} \quad \log 2662937 &= \log 2662900 + \cdot 000000163 \times 37 \\ &= 6\cdot4253549 + \cdot 0000060 \\ &= 6\cdot4253609, \end{aligned}$$

omitting the figures beyond the seventh decimal place.

81. The trouble of this multiplication is saved in the Tables. In a column by the side of the logarithms in each page is placed the difference of two consecutive logarithms, and in another column the proportional parts, or tenths of these differences, as shewn in the annexed columns. Observe that the cyphers after the decimal point are omitted for brevity: the numbers 16, 33, 49, &c., are 163 multiplied by 1, 2, 3, &c., with the last figure cut off for the division by 10; and 33 is placed opposite to 2, rather than 32, because 33 is nearer to 32·6 than 32 is, and so for 49, 82, 98, 147. When we require hundredths we cut off one more figure, and increase the last figure by unity, when the figure cut off is greater than 4. Thus to multiply 163 by 37 we take for the figures required  $49 + 11 = 60$ , which gives the same result as above.

D.	Pro.	
	163	
163	1	16
	2	33
	3	49
	4	65
	5	82
	6	98
	7	114
	8	130
	9	147

Our work may be arranged as follows:—

From the Tables

$$\begin{array}{rcl} \log 2662900 & = & 6\cdot4253549 \\ \text{diff. for} \quad 30 & = & 49 \\ \text{,,} \quad 7 & = & 11 \\ \hline \therefore \log 2662937 & = & 6\cdot4253609 \end{array}$$

82. The reverse process of finding a number, whose logarithm is given, more exactly than can be found directly from the Tables is similar. Thus to find the number whose logarithm is 3·4253685: we find  $\log 2662900 = 6\cdot4253549$ , difference is 136, and difference for 100 is 163, omitting the cyphers. Therefore we have to add  $\frac{136}{163}$  of 100, or 84 nearly, to 2662900 to obtain

the required number; therefore 6.4253685 is the logarithm of 2662984, and 3.4253685 is the logarithm of 2662.984 the required number.

Making use of the table of proportional parts the work may be exhibited as follows :—

6.4253685 is log. of required number,

6.4253549 is log. of 2662900

136

130 is diff. of 80

6 is diff. of 4

∴ required number = 2662984

In practice the characteristic of the logarithm and the decimal point of the number may be omitted till the work is complete.

83. The principle above applied to the logarithms of numbers, applies equally well to the determination of natural sines, &c. In the latter case however the change of successive sines is not so uniform. There are 90000 numbers with 5 digits, the mantissæ of whose logarithms all lie between 0 and 1, whereas the Tables give the Trigonometrical Ratios of 5400 angles, whose sines and cosines lie between 0 and 1, so that the differences of successive sines and cosines are usually larger and less uniform than the differences of the logarithms of successive numbers; and what is said of the sines and cosines applies with still greater force to the other trigonometrical ratios, since their values extend between greater limits. Hence the differences only are given in the Tables, and the proportional parts are not calculated for them, as in the logarithms of common numbers, since to do so would render the Tables inconveniently bulky.

The mode of using the Tables will be learnt from the following examples, cyphers being omitted after decimal points for brevity.

To find  $\sin 14^\circ. 18'. 33''. 56$ .

From the Tables  $\sin 14^\circ. 19' = .2472809$

$\sin 14^\circ. 18' = .2469990$

diff. for  $1' = 2819$

$$\therefore \text{required diff.} = \frac{33.56}{60} \times 2819 = 1576 \text{ nearly,}$$

$$\therefore \sin 14^\circ. 18'. 33''.56 = .2469990 + .0001576 \\ = .2471566.$$

To find  $\cos 14^\circ. 18'. 33''.56$ ,

$$\cos 14^\circ. 18' = .9690157$$

$$\cos 14^\circ. 19' = .9689438$$

$$\text{diff. for } 1' = 719$$

$$\therefore \text{required diff.} = \frac{33.56}{60} \times 719 = 402 \text{ nearly;}$$

$$\therefore \cos 14^\circ. 18'. 33''.56 = .9690157 - .000402 \\ = .9689755.$$

Notice that in this case the difference is subtracted, because the cosine of an angle diminishes as the angle itself increases. The same observation applies to the cosecant and cotangent.

Given  $\tan \theta = .75$ , to find  $\theta$ .

From the Tables  $\tan 36^\circ. 53' = .7503665$

$$\tan 36^\circ. 52' = .7499119$$

$$\text{diff. for } 1' = 4546,$$

and  $\tan \theta - \tan 36^\circ. 52' = .75 - .7499119 = 881,$

$$\therefore \text{required seconds} = \frac{881}{4546} \text{ of } 60'' = 11''.63,$$

and  $\theta = 36^\circ. 52'. 11''.63.$

Given  $\text{cosec } \theta = 1.6 = 1.6666667$  nearly, to find  $\theta$ .

From the Tables  $\text{cosec } 36^\circ. 52' = 1.6667920$

$$\text{cosec } 36^\circ. 53' = 1.6661458$$

$$\text{diff. for } 1' = 6462$$

$$\text{cosec } 36^\circ. 52' - \text{cosec } \theta = 1.6667920 - 1.6666667 = 1253$$

$$\therefore \text{required seconds} = \frac{1253}{6462} \text{ of } 60'' = 11''.63;$$

$$\therefore \theta = 36^\circ. 52'. 11''.63.$$

In this case since  $\tan^{-1} .75 = \text{cosec}^{-1} 1.6$  the results ought to agree exactly. The results obtained above do agree, but if the operation

had been carried further  $\tan^{-1} \cdot 75$  would have been found slightly less, and  $\operatorname{cosec}^{-1} 1 \cdot 6$  slightly greater than the results obtained. The Tables being exact to 7 decimal places, the true values of the Ratios are slightly in excess or defect of the values given, and the true differences likewise. The example above shews that the method employed may be generally relied upon to give a true result as far as the hundredth part of a second.

84. The same principle also applies to the determination of the logarithms of the Trigonometrical Ratios not found exactly in the Tables, and the reverse. The following examples illustrate the method.

To find  $L \sec 27^\circ. 13'. 15''. 34$ .

From the Tables  $L \sec 27^\circ. 14' = 10 \cdot 0510248$   
 $L \sec 27^\circ. 13' = 10 \cdot 0509598$   
 diff. for  $1' = 650$

$$\therefore \frac{15 \cdot 34}{60} \times 650 = 166 ;$$

$$\therefore L \sec 27^\circ. 13'. 15''. 34 = 10 \cdot 0509598 + \cdot 0000166 \\ = 10 \cdot 0509764.$$

In this case the difference for  $1'$  is so small that each hundredth of a second does not affect the result; thus  $15''. 33$  to  $15''. 37$  inclusive would give the same result.

Given  $L \cot \theta = 10 \cdot 1789937$ , to find  $\theta$ .

From the Tables  $L \cot 33^\circ. 30' = 10 \cdot 1792171$   
 $L \cot 33^\circ. 31' = 10 \cdot 1789426$   
 diff. for  $1' = 2745$ .

Also  $L \cot 33^\circ. 30' = 10 \cdot 1792171$   
 $L \cot \theta = 10 \cdot 1789937$   
 diff.  $= 2234$ .

And  $\frac{2234}{2745}$  of  $60'' = 48''. 83$  ;

$$\therefore \theta = 33^\circ. 30'. 48''. 83.$$

85. In the preceding Articles we have explained the method of employing proportional parts : it only remains to point out that this

method may fail from two causes, either the successive differences may be too irregular, or they may be too small. The second cause of failure applies partially to the determination of the second additional figure in numbers from 85000 to 100000; and the Trigonometrical Ratios of angles very small or very nearly  $90^\circ$ , and their logarithms, cannot be determined by this method from one or both causes.

For a theoretical proof of the principle of proportional parts the student is referred to more advanced treatises on the subject. A proof of the principle for the logarithms of common numbers is indicated in the Examples 16, 17, 18 C. of Section VI.

### EXAMPLES. (A).

1. Given  $\log 1752 = 3.2435341$ ,  $\log 1752.1 = 3.2435589$ , find  $\log 1752.87$ . *Ans.* 3.2435557.

2. Given  $\log 48664 = 4.6872078$ ,  
 $\log 48665 = 4.6872167$ .

Calculate the table of proportional parts, and by means of it determine  $\log 4866.439$ ,  $\log 48.66407$ , and the numbers whose logs are  $\bar{1}.6872100$ ,  $8.6872135$ .

*Ans.* 3.6872113, 1.6872084, .4866425, 486646500.

3. Given  $\sin 16^\circ.24' = .2823415$ ,  $\sin 16^\circ.25' = .2826205$ , find  $\sin 16^\circ.24'.17'' \cdot 3$ ,  $\cos 73^\circ.35'.22'' \cdot 5$ ,  $\sin^{-1}.2824907$ .

*Ans.* .2824219, .2825159,  $16^\circ.24'.32'' \cdot 08$ .

4. Given  $\cot 32^\circ.36' = 1.5636564$ , diff. for  $1' = 10016$ , find  $\cot 32^\circ.36'.27'' \cdot 59$ ,  $\cot^{-1}.5628793$ .

*Ans.* 1.5631958,  $32^\circ.36'.46'' \cdot 55$ .

5. Given  $L \tan 34^\circ.36' = 9.8387571$ , diff. for  $1' = 2702$ , find  $L \tan 34^\circ.36'.27'' \cdot 3$ , and angle whose  $L \tan$  is 9.8388972.

*Ans.* 9.8388800,  $34^\circ.36'.31'' \cdot 11$ .

6. Given  $L \operatorname{cosec} 40^\circ.48' = 10.1848072$ , diff. for  $1' = 1463$ , find  $L \operatorname{cosec} 40^\circ.48'.17'' \cdot 9$ , and angle whose  $L \sec$  is 10.1847832.

*Ans.* 10.1847636,  $49^\circ.41'.50'' \cdot 16$ .

7. Given  $a=18$ ,  $b=12$ ,  $c=10$ , find  $\alpha$ ,  $\beta$ ,  $\gamma$ .

Given  $\sqrt{2}=1.4142136$ ,  $\tan 54^\circ.44'=1.4140943$ ,  
 $\tan 54^\circ.45'=1.4149673$ ,  $\tan 15^\circ.47'=.2826573$ ,  
 $\tan 15^\circ.48'=.2829715$ .

*Ans.*  $\alpha=109^\circ.28'.16''.39$ ,  $\gamma=31^\circ.35'.10''.8$ ,  $\beta=38^\circ.56'.32''.81$ .

8. If  $a=5$ ,  $b=20$ ,  $\gamma=90^\circ$ , find  $\alpha$ ,  $\beta$ .

Given  $\log 2$ ,  $L \tan 75^\circ.58'=10.6021537$ ,  
 $L \tan 75^\circ.57'=10.6016170$ .

*Ans.*  $\alpha=14^\circ.2'.10''.48$ ,  $\beta=75^\circ.57'.49''.52$ .

9. If  $a=18$ ,  $b=2$ ,  $\gamma=55^\circ$ , find  $\alpha$ ,  $\beta$ .

Given  $\log 2$ ,  $L \cot 27^\circ.30'=10.2835233$ ,  
 $L \tan 56^\circ.56'=10.1863769$ , diff. for  $1'=2763$ .

*Ans.*  $\alpha=119^\circ.26'.51''.33$ ,  $\beta=5^\circ.33'.8''.67$ .

#### EXAMPLES. (B).

1. If  $\gamma=90^\circ$ ,  $a=141$ ,  $b=193$ , solve the triangle.

Given  $\log 141=2.1492191$ ,  $\log 193=2.2855573$ ,  
 $\log 2390=3.3783979$ ,  $\log 2391=3.3785796$ ;  
 $L \tan 36^\circ.9'=9.8636500$ , diff. for  $1'=2652$ ,  
 $L \sin 36^\circ.9'=9.7707793$ , diff. for  $1'=1729$ .

*Ans.*  $c=239.018$ ,  $\alpha=36^\circ.9'.2''.7$ ,  $\beta=53^\circ.50'.57''.3$ .

2. If  $a=254$ ,  $\beta=16^\circ$ ,  $\gamma=64^\circ$ , find  $b$ .

Given  $\log 2.54=.4048337$ ,  $L \sin 80^\circ=9.9933515$ ,  
 $L \sin 16^\circ=9.4403381$ ,  $\log 7.109=.8518085$ ,  
 $\log 7.11=.8518696$ . *Ans.*  $b=71.0919$ .

3. If  $b=246.35$ ,  $a=197.63$ ,  $\alpha=34^\circ.27'$ , find  $\beta$ .

Given  $\log 24635=4.3915526$ ,  $\log 19763=4.2958529$ ,  
 $L \sin 44^\circ.50'=9.8482180$ ,  $L \sin 44^\circ.51'=9.8483450$ ,  
 $L \sin 34^\circ.27'=9.7525761$ .

*Ans.*  $\beta=44^\circ.50'.27''.3$ , or  $135^\circ.8'.32''.7$ .

4. If  $a = 254.13$ ,  $b = 375.16$ ,  $\beta = 61^\circ.10'.4''.2$ , find  $\alpha$ .

Given  $\log 25413 = 4.4050559$ ,  $\log 37516 = 4.5742165$ ,

$L \sin 61^\circ.10' = 9.9425171$ , diff. for  $1' = 695$ ,

$L \sin 36^\circ.24' = 9.7733614$ . *Ans.*  $\alpha = 36^\circ.24'$

5. If  $a = 53^\circ.24'$ ,  $\beta = 66^\circ.27'$ ,  $c = 338.65$ , find  $\alpha$ .

Given  $L \sin 53^\circ.24' = 9.9046168$ ,  $\log 33865 = 4.5297511$ ,

$L \sin 60^\circ.9' = 9.9381851$ ,  $\log 31346 = 4.4961821$ ,

$\log 31347 = 4.4961960$ .

*Ans.*  $\alpha = 313.460$

6. If  $a = 275.35$ ,  $b = 189.28$ ,  $c = 301.47$ , find  $\alpha$ .

Given  $\log 38305 = 4.5832555$ ,  $\log 8158 = 3.9115837$ ,

$\log 10770 = 4.0322157$ ,  $\log 19377 = 4.2872865$ ,

$L \tan 31^\circ.45' = 9.7915635$ ,  $L \tan 31^\circ.46' = 9.7918458$ .

*Ans.*  $\alpha = 63^\circ.30'.57''.8$

7. If  $a = 673.12$ ,  $b = 415.89$ ,  $\gamma = 90^\circ$ , find  $\alpha$ ,  $\beta$ .

Given  $\log 67312 = 4.8280925$ ,  $L \tan 58^\circ.17' = 10.2090013$ ,

$\log 41589 = 4.6189785$ ,  $L \tan 58^\circ.18' = 10.2092839$ .

*Ans.*  $\alpha = 58^\circ.17'.23''.92$ ,  $\beta = 31^\circ.42'.36''.08$

8. If  $a = 456.12$ ,  $b = 296.86$ ,  $\gamma = 74^\circ.20'$ , find  $\alpha$ ,  $\beta$ .

Given  $L \cot 37^\circ.10' = 10.1202593$ ,  $\log 15926 = 4.2021067$ ,

$L \tan 15^\circ.35' = 9.4454352$ ,  $\log 75298 = 4.8767834$ ,

$L \tan 15^\circ.36' = 9.4459232$ .

*Ans.*  $\alpha = 68^\circ.25'.18''.12$ ,  $\beta = 37^\circ.14'.41''.88$

9. If  $a = 576.12$ ,  $c = 873.14$ ,  $\gamma = 90^\circ$ : find  $\alpha$ ,  $\beta$ .

Given  $\log 57612 = 4.7605054$ ,  $\log 87314 = 4.9410839$ ,

$L \sin 41^\circ.17' = 9.8194012$ , diff. for  $1' = 1438$ .

*Ans.*  $\alpha = 41^\circ.17'.8''.47$ ,  $\beta = 48^\circ.42'.51''.58$

10. If  $a = 4439$ ,  $b = 4861$ ,  $c = 8583$ , find  $\beta$ .

Given  $\log 8.9415 = .9514104$ ,  $\log 4.439 = .6472851$ ,

$\log 4.0805 = .6107027$ ,  $\log 8.583 = .9336391$ ,

$L \cos 11^\circ.52' = 9.9906180$ ,  $L \cos 11^\circ.53' = 9.9905914$ .

*Ans.*  $\beta = 23^\circ.45'.46''.40$

11. If  $\alpha = 38$ ,  $\beta = 48^\circ$ ,  $\gamma = 54^\circ$ , find  $b$ ,  $c$ .

Given  $L \sin 54^\circ = 9.9079576$ ,  $L \sin 78^\circ = 9.9904044$ ,

$L \sin 48^\circ = 9.8710735$ ,  $\log 38 = 1.5797836$ ,

$\log 2.8870 = .4604468$ ,  $\text{diff. for } 1 = 150$ ,

$\log 3.1429 = .4973306$ ,  $\text{diff. for } 1 = 138$ .

*Ans.*  $b = 28.87039$ ,  $c = 31.42945$ .

12. One side of a triangle is double the other, and the included angle is a right angle, find the other angles.

Given  $\log 2$ ,  $L \cot 26^\circ.34' = 10.3009994$ ,  $\text{diff. for } 1' = 3159$ .

*Ans.*  $26^\circ.33'.54''.19$ ,  $63^\circ.26'.5''.81$ .

13. If  $a = 4$ ,  $b = 5$ ,  $c = 6$ , find  $\beta$ .

Given  $\log 2$ ,  $L \cos 27^\circ.53' = 9.9464040$ ,  $L \cos 27^\circ.54' = 9.9463371$ .

*Ans.*  $\beta = 55^\circ.46'.16''.16$ .

14. A rope dancer wishes to ascend a tower 100 feet high by a rope 196 feet long; at what inclination must he be able to walk up the rope?

Given  $\log 2$ ,  $\log 7$ ,  $L \sin 30^\circ.40' = 9.7076064$ ,  $\text{diff. for } 1' = 2130$ .

*Ans.*  $30^\circ.40'.38''.76$ .

15. If  $L \sin 16^\circ.26' = 9.4516322$ ,  $L \sin 26^\circ.27' = 9.4520603$ , find  $\sin 16^\circ.26'.46''$ .

Given  $\log 2831.1 = 3.4519552$ ,  $\log 2.8312 = .4519705$ .

*Ans.*  $.2831133$ .

# EXAMPLES. (C).

1. If  $b = 354$ ,  $c = 426$ ,  $\alpha = 49^\circ.16'$ , find  $\beta$ ,  $\gamma$ .

Given  $\log 2$ ,  $\log 3$ ,  $\log 13$ ,  $L \cot 24^\circ.38' = 10.3386231$ ,

$L \tan 11^\circ.22' = 9.3032609$ ,  $L \tan 11^\circ.23' = 9.3039143$ .

*Ans.*  $\beta = 53^\circ.59'.4''.9$ ,  $\gamma = 76^\circ.44'.55''.1$ .

2. If  $\alpha = 60^\circ$ ,  $b = 14$ ,  $c = 11$ , find  $\beta$ ,  $\gamma$ .

Given  $\log 2$ ,  $\log 3$ ,  $L \tan 11^\circ.44' = 9.3174299$ ,  $\text{diff. for } 1' = 6341$ .

*Ans.*  $\beta = 71^\circ.44'.29''.52$ ,  $\gamma = 48^\circ.15'.30''.48$ .



3. If  $a = 135$ ,  $b = 105$ ,  $\gamma = 60^\circ$ , find  $\alpha$ ,  $\beta$ .

Given  $\log 2$ ,  $\log 3$ ,  $L \tan 12^\circ. 12' = 9.3348711$ ,

$L \tan 12^\circ. 13' = 9.3354823$ .

*Ans.*  $\alpha = 72^\circ. 12'. 58''.8$ ,  $\beta = 47^\circ. 47'. 1'' - 2$

4. If  $a = 5$ ,  $b = 12$ ,  $c = 13$ , find  $\alpha$ ,  $\beta$ ,  $\gamma$ .

Given  $\log 2$ ,  $L \tan 11^\circ. 18'. 30'' = 9.3009670$ ,

$L \tan 11^\circ. 18'. 40'' = 9.3010764$ .

*Ans.*  $\alpha = 22^\circ. 37'. 11''.52$ ,  $\beta = 67^\circ. 22'. 48''.48$ ,  $\gamma = 90^\circ$ .

5. Two sides of a triangle are as 5 : 9, and the included angle is a right angle, find the other angles.

Given  $\log 2$ ,  $\log 3$ ,  $L \tan 29^\circ. 3' = 9.7446453$ ,

$L \tan 29^\circ. 4' = 9.7449428$ .

*Ans.*  $29^\circ. 3'. 16''.55$ ,  $60^\circ. 56'. 43''.45$ .

6. Prove that  $\tan\left(\beta + \frac{\alpha}{2}\right) = \frac{c+b}{c-b} \tan \frac{\alpha}{2}$ ; and if  $b : c = 21 : 29$ , and  $\alpha = 46^\circ. 37'. 24''$ , find  $\beta$ .

Given  $\log 2$ ,  $L \tan 23^\circ. 18' = 9.6341426$ , diff. for  $1' = 3477$ ,

$L \tan 69^\circ. 37' = 10.4299645$ ,  $L \tan 69^\circ. 38' = 10.4303516$ .

*Ans.*  $\beta = 46^\circ. 19'. 4''.73$ .

7. If  $a = 140.5$ ,  $b = 170.6$ ,  $\alpha = 40^\circ$ , find  $\beta$ ,  $\gamma$ .

Given  $L \sin 40^\circ = 9.8080675$ ,  $\log 1405 = 3.1476763$ ,

$\log 1706 = 3.2319790$ ,

$L \sin 51^\circ. 18' = 9.8923342$ , diff. for  $1' = 1012$ .

*Ans.*  $\beta = 51^\circ. 18'. 21''.34$ ,  $\gamma = 88^\circ. 41'. 38''.66$ .

8. If  $a = 1.56234$ ,  $b = .43766$ ,  $\gamma = 58^\circ. 42'. 6''.1$ , find  $\alpha$ ,  $\beta$ .

Given  $\log 56234 = 4.750150$ ,  $\log \cot 29^\circ. 21' = .2500150$ ,

$\log \cot 29^\circ. 22' = .2497194$ .

*Ans.*  $\alpha = 105^\circ. 38'. 56''.95$ ,  $\beta = 15^\circ. 38'. 56''.95$ .

9. If  $a = 3$ ,  $b = 1$ ,  $\gamma = 53^\circ. 7'. 48''$ , find  $c$  without determining  $\alpha$  and  $\beta$ .

Given  $\log 2$ ,  $\log 25298 = 4.4030862$ , diff. for  $1 = 172$ ,

$L \cos 26^\circ. 33'. 54'' = 9.9515452$ ,  $L \tan 26^\circ. 33'. 54'' = 9.6989700$ -

*Ans.*  $c = 2.529823$ .

10. If  $\tan \theta = \frac{2\sqrt{ab} \sin \frac{\gamma}{2}}{a-b}$ , and  $a = 5$ ,  $b = 2$ ,  $\gamma = 120^\circ$ , find  $\theta$ .

Given  $\log 3$ ,  $L \tan 61^\circ.17' = 10.2613287$ ,

$L \tan 61^\circ.18' = 10.2616286$ .

*Ans.*  $\theta = 61^\circ.17'.22''.13$ .

11. One angle of a triangle is  $60^\circ$ , and the side opposite is to the difference of the two sides including it as  $9 : 2$ ; find the remaining angles.

Given  $\log 3$ ,  $L \cos 78^\circ.54'.10'' = 9.2843730$ ,

$L \cos 78^\circ.54'.20'' = 9.2842656$ .

*Ans.*  $71^\circ.5'.44''.88$ ,  $48^\circ.54'.15''.12$ .

### MISCELLANEOUS EXAMPLES.

1. Prove that the ratios of the angles denoted by a French and English degree, minute, second, are expressed by

$$\frac{3.3}{2.5}, \frac{3.3^2}{2.5^2}, \frac{3.3^3}{2.5^3}, \text{ respectively.}$$

2. Shew that  $m' = m + \frac{23}{27}m$ , and  $n'' = 3n' + \frac{7}{81}n'$ .

3. Shew that the area of a triangle is equal to

$$\sin \frac{\alpha}{2} \sin \frac{\beta}{2} \sin \frac{\gamma}{2} \left( \frac{a^2}{\sin \alpha} + \frac{b^2}{\sin \beta} + \frac{c^2}{\sin \gamma} \right).$$

4. One regular figure has twice as many sides as another, and an angle of the first is one-third as large again as an angle of the second; find the interior angles of each. *Ans.*  $144^\circ, 108^\circ$ .

5. Solve the equations :

$$\sin 3\phi \sin 4\phi - \sin 2\theta \sin \phi = \frac{1}{2} \cos 4\phi,$$

$$4 \sin (\phi - \theta) \cos (\phi + \theta) = \cos 6\phi \operatorname{cosec} \phi.$$

$$\text{Ans. } \phi = 60^\circ, \theta = \frac{1}{2} \sin^{-1} \frac{1}{2\sqrt{3}}.$$

6. If regular pentagons be inscribed and circumscribed about a circle, then perimeter of the former : perimeter of the latter  $= \sqrt{5} + 1 : 4$ .

7. If  $\cos \theta = \cos \alpha \cos \beta + \sin \alpha \sin \beta \cos A$ ,

and  $\sin^2 \phi = \sin \alpha \sin \beta \cos^2 \frac{A}{2},$

then  $\sin^2 \frac{\theta}{2} = \sin \left( \frac{\alpha + \beta}{2} + \phi \right) \sin \left( \frac{\alpha + \beta}{2} - \phi \right).$

8. If  $\tan x = \cos a \tan y$ , then

$$\tan (y - x) = \frac{\tan^2 \frac{a}{2} \sin 2y}{1 + \tan^2 \frac{a}{2} \cos 2y}.$$

9. Eliminate  $\theta$  between the equations

$$m = \operatorname{cosec} \theta - \sin \theta, \quad n = \sec \theta - \cos \theta.$$

$$\text{Ans. } (mn)^{\frac{2}{3}} (m^{\frac{2}{3}} + n^{\frac{2}{3}}) = 1.$$

10. If  $\tan \frac{\alpha}{2} = \tan^3 \frac{\beta}{2}$ , and  $\tan \beta = 2 \tan \phi$ ,

then

$$\phi = \frac{\alpha + \beta}{2}.$$

11. If  $a \cos \theta + b \cos (\theta + \alpha)$  be put in the form

$$A \cos (\theta + B),$$

find the values of  $A$  and  $B$ .

$$\text{Ans. } A = \sqrt{(a^2 + b^2 + 2ab \cos \alpha)}, \quad B = \tan^{-1} \frac{b \sin \alpha}{a + b \cos \alpha}.$$

12. Divide a given angle  $\alpha$  into two parts, whose sines are in the ratio

$$m : n. \quad \text{Ans. } \sin^{-1} \frac{m \sin \alpha}{\sqrt{m^2 + n^2 + 2mn \cos \alpha}}, \quad \sin^{-1} \frac{n \sin \alpha}{\sqrt{m^2 + n^2 + 2mn \cos \alpha}}.$$

13. If  $\cos (\phi - \alpha)$ ,  $\cos \phi$ ,  $\cos (\phi + \alpha)$  are in H.P., then

$$\cos \phi = \sqrt{2} \cos \frac{\alpha}{2}.$$

14. The elevation of a tower standing on a horizontal plain is observed;  $a$  feet nearer it is found to be  $45^\circ$ ;  $b$  feet nearer still it is the complement of what it was at the first station; shew that the height of the tower is  $\frac{ab}{a-b}$  feet.

15. The distance between the centres of the inscribed circle and of the escribed circle touching the side  $BC$  is

$$a \sec \frac{\alpha}{2}.$$

16. Prove the following expression for the area of a triangle,

$$\frac{1}{2} \left( \frac{a^2 \cos \beta \cos \gamma}{\sin \alpha} + \frac{b^2 \cos \alpha \cos \gamma}{\sin \beta} + \frac{c^2 \cos \alpha \cos \beta}{\sin \gamma} \right).$$

17. A number of spheres are put in a hollow cone one above another, and each touching the one above, the one below, and the sides of the cone; shew that the radii of the spheres form a geometrical progression of which the common ratio is

$$\tan^2 \left( 45^\circ + \frac{\alpha}{4} \right),$$

$\alpha$  being the vertical angle of the cone.

18. From one of the angles of a rectangle a perpendicular is drawn to its diagonal, and from the point of intersection lines are drawn perpendicular to the sides which contain the opposite angle; shew that if  $p, p'$  be the perpendiculars last drawn, and  $d$  the diagonal of the rectangle,

$$p^2 + p'^2 = d^2.$$

19. If two circles whose radii are  $a, b$  touch each other externally, and if  $\theta$  be the angle between their common tangents, shew that

$$\sin \theta = \frac{4(a-b)\sqrt{ab}}{(a+b)^2}.$$

$$\begin{aligned} 20. \quad & \text{If } (a+b) \tan(\theta - \phi) = (a-b) \tan(\theta + \phi), \\ & a \cos 2\phi + b \cos 2\theta = c, \end{aligned}$$

$$\text{then} \quad b^2 - a^2 - c^2 + 2ac \cos 2\phi = 0.$$

$$21. \quad \text{If } \cot \beta - \cot \theta = \cot(\alpha + \theta) + \cot(\alpha - \beta),$$

$$\text{then} \quad \frac{\sin(\alpha - \beta)}{\sin \beta} = \frac{\sin(\alpha + \theta)}{\sin \theta}.$$

$$22. \quad \text{If } \tan \frac{\alpha}{2} = 2 - \sqrt{3}, \text{ find } \sin \alpha.$$

23. The elevation of a tower at a place  $A$  due south of it is  $30^\circ$ ; and at a place  $B$ , due west of  $A$ , and at the distance  $a$  from it, the elevation is  $18^\circ$ : shew that the height of the tower is

$$\frac{a}{\sqrt{(2\sqrt{5} + 2)}}.$$

$$24. \quad \text{If } \sin \alpha + \sin(\theta - \alpha) + \sin(2\theta + \alpha) = \sin(\theta + \alpha) + \sin(2\theta - \alpha), \text{ then}$$

$$\theta = \cos^{-1} \frac{\pm \sqrt{5} + 1}{4}.$$

$$25. \quad \text{If } \frac{m \tan(\alpha - x)}{\cos^2 x} = \frac{n \tan x}{\cos^2(\alpha - x)},$$

$$\text{then} \quad \tan(\alpha - 2x) = \frac{n - m}{n + m} \tan \alpha.$$

26. If  $a, a'$  be homologous sides of similar triangles inscribed in and described about a given circle, then

$$a = 4a' \sin \frac{\alpha}{2} \sin \frac{\beta}{2} \sin \frac{\gamma}{2}.$$

27. In the ambiguous case in the solution of triangles, if  $\alpha, \beta, \gamma, \alpha', \beta', \gamma'$  be the angles of the triangle in the two solutions,

$$\frac{\sin \gamma}{\sin \beta} + \frac{\sin \gamma'}{\sin \beta'} = 2 \cos \alpha.$$

28. The elevation of two clouds to a person in the same line with them is  $\alpha$ . When standing on the shadow of one of them its elevation is  $2\alpha$ , and the other is vertically over him. Shew that the heights of the clouds are as  $2 \cos^2 \alpha : 1$ .

29. At each of three stations in the same horizontal plane, and at given distances from each other, the elevation of a tower is observed to be  $\alpha$ ; find the height of the tower.

30. In any triangle

$$a \cos 2\beta + 2b \cos \alpha \cos \beta + a \cos 2\gamma + 2c \cos \alpha \cos \gamma = 0.$$

31. If  $\sqrt{(a^2 - x^2)} \cos \beta + a \sin \alpha = x \sin \beta$ , then

$$x = \mp a \cos (\alpha \pm \beta).$$

32. The elevation of a tower on a horizontal plane is observed: on advancing  $a$  feet nearer its elevation is found to be the complement of the former; on again advancing its elevation is found to be double of its first elevation: shew that the last station is  $\frac{a}{2}$  feet from the foot of the tower.

33. If the line bisecting the right angle  $\gamma$  of a triangle divide the hypotenuse into parts  $x, y$ , then

$$x : y = 1 - t : 1 + t, \text{ where } t = \tan \frac{\alpha - \beta}{2}.$$

34. If

$$\sin \theta + \sin 2\theta = m,$$

$$\cos \theta + \cos 2\theta = n,$$

then  $(m^2 + n^2)^2 - 3(m^2 + n^2) - 2n = 0$ .

35.  $(\sin \alpha + \sec \alpha)^2 + (\cos \alpha + \operatorname{cosec} \alpha)^2 = (1 + \sec \alpha \operatorname{cosec} \alpha)^2$ .

36.  $\tan^{-1} \left\{ \frac{1}{2} (\cot^2 \theta - \tan^2 \theta) \right\} = 2 \tan^{-1} (\cos 2\theta)$ .

37. A person observes that a tower and spire on a hill subtend the same angle  $\beta$ ; on ascending the hill  $a$  feet, he finds the tower subtends an angle  $\gamma$  and the spire again subtends an angle  $\beta$ ; shew that the height of the spire is

$$\frac{a \sin \beta}{\sin (\gamma - \beta)}.$$

38. If the diameter  $2R$  of a semicircle be divided into any two parts, and on these parts semicircles be described whose radii are  $r_1, r_2$ ; then the radius of the circle which touches all three semicircles is

$$\frac{Rr_1r_2}{R^2 - r_1r_2}.$$

39. Shew that  $\frac{\sin 2\alpha + \sin 4\alpha + \sin 6\alpha}{4 \cos^2 \alpha \cos^2 2\alpha} = \tan \alpha + \tan 2\alpha$ .

40. A person on the top of a mountain observes the depression of an object on the plane below him: he then turns through an angle and observes the depression of another object on the same plane. On descending the mountain he finds the distance between the objects is equal to the height of the mountain. Shew that the height of the mountain is also  $d$ .

41. If  $\alpha, \beta, \gamma$  be in arithmetical progression,

$$\frac{\sin(\alpha + \beta)}{\sin(\alpha - \beta)} - \frac{\sin(\beta + \gamma)}{\sin(\beta - \gamma)} = 2 \cos 2\beta.$$

42. A tower stands in a field in the form of an equilateral triangle. It subtends angles  $\tan^{-1} \frac{4}{\sqrt{3}}, \tan^{-1} \frac{4}{\sqrt{7}}, \tan^{-1} \frac{4}{\sqrt{7}}$  at the three corners: the distance from the tower to the nearest corner is equal to a side of the field.

43. If  $\tan^{-1} \frac{1}{x-1} - \tan^{-1} \frac{1}{x+1} = \frac{\pi}{12}$ ,

then  $x = \sqrt{3} + 1$ .

44. If  $\tan(\theta + \phi) = m \tan(\theta - \phi)$ ,

and  $a \cos^2 \phi = b \cos^2 \theta$ ,

then  $\cos^2 \theta = \frac{(m+1)^2 b - (m-1)^2 a}{(m+1)^2 b^2 - (m-1)^2 a^2} a$ .

45. If  $x, y, z$  be the lines joining the feet of the perpendiculars from the vertices of a triangle upon the opposite sides,

then  $\frac{x}{a^2} + \frac{y}{b^2} + \frac{z}{c^2} = \frac{a^2 + b^2 + c^2}{2abc}$ .

46. If  $\cos(\alpha - \beta) \sin(\gamma - \delta) = \cos(\alpha + \beta) \sin(\gamma + \delta)$ ,

then  $\cot \delta = \cot \alpha \cot \beta \cot \gamma$ .

47. Shew that

$$\sin \alpha \cos(\beta - \gamma) - \sin \beta \cos(\gamma - \alpha) = \sin(\alpha - \beta) \cos \gamma.$$

48. Shew that

$$\frac{\cos \alpha + \sin \gamma - \sin \beta}{\cos \beta + \sin \gamma - \sin \alpha} = \frac{1 + \tan \frac{\alpha}{2}}{1 + \tan \frac{\beta}{2}}, \text{ if } \alpha + \beta + \gamma = 90^\circ.$$

49. Shew that

$$\frac{2 \operatorname{cosec} 2\theta - \sec \theta}{2 \operatorname{cosec} 2\theta + \sec \theta} = \cot^2 \left( 45^\circ + \frac{\theta}{2} \right).$$

# MISCELLANEOUS EXAMPLES.

50. If  $\alpha + \beta + \gamma = 0$ ,

$$\cos \alpha + \cos \beta + \cos \gamma + 1 = 4 \cos \frac{\alpha}{2} \cos \frac{\beta}{2} \cos \frac{\gamma}{2}.$$

51. If the area of a triangle be constant, the sum of the cotangents of the angles varies as the sum of the squares of the sides.

52. A column on a pedestal 20 feet high subtends an angle of  $45^\circ$  to a person on the ground; on approaching 20 feet it again subtends an angle of  $45^\circ$ . The height of the column is 100 feet.

53. Shew that

$$\cos^2 \sigma + \cos^2 (\sigma - \theta) + \cos^2 (\sigma - \phi) + \cos^2 (\sigma - \psi) = 2 + 2 \cos \theta \cos \phi \cos \psi,$$

where

$$\sigma = \frac{\theta + \phi + \psi}{2}.$$

54. If  $\tan \theta = \cos 2\alpha$ ,

then

$$\sin 2\theta = \frac{1 - \tan^4 \alpha}{1 + \tan^4 \alpha}.$$

55. Shew that

$$2 \tan \left\{ \cos^{-1} \frac{1}{\sqrt{1+x^2}} - \cos^{-1} \frac{x}{\sqrt{1+x^2}} \right\} = x - \frac{1}{x}.$$

56. Shew that in a triangle

$$(r_a - r)(r_b - r)(r_c - r) = 4Rr^3.$$

57. If  $a'b'c'$  be the perpendiculars from the centre of the circumscribed circle upon the sides  $a, b, c$  respectively, then

$$\frac{a}{a'} + \frac{b}{b'} + \frac{c}{c'} = \frac{1}{4} \cdot \frac{abc}{a'b'c'}.$$

58. Shew that  $\cos^{-1} \frac{1}{2} - \sin^{-1} \frac{1}{5} = \tan^{-1} \left( \frac{9\sqrt{3} - 8\sqrt{2}}{5} \right)$ .

59. If  $\alpha = \tan^{-1} \frac{1}{7}$ ,  $\beta = \tan^{-1} \frac{1}{3}$ , then

$$\cos 2\alpha = \sin 4\beta.$$

60. If  $2 \tan x + \tan (\alpha - x) = \tan (\beta + x)$ , then

$$\tan x = \frac{\sin (\alpha - \beta)}{2 \sin \alpha \sin \beta}.$$



61. From the top of a tower the depressions  $\alpha$ ,  $\beta$  of two objects in the same horizontal plane with the foot of the tower are observed, and also the angle  $\omega$  which they subtend; and the distance  $a$  between them is known; the height of the tower is

$$\frac{a \sin \alpha \sin \beta}{\sqrt{(\sin^2 \alpha + \sin^2 \beta - 2 \sin \alpha \sin \beta \cos \omega)}}.$$

62. Shew that

$$\sin \alpha + \sin \beta + \sin \gamma - \sin (\alpha + \beta + \gamma) = 4 \sin \frac{\alpha + \beta}{2} \sin \frac{\alpha + \gamma}{2} \sin \frac{\beta + \gamma}{2}.$$

63. If

$$\sin \theta + \sin \phi = m,$$

and

$$\cos \theta + \cos \phi = n,$$

then

$$\cos (\theta + \phi) = \frac{n^2 - m^2}{n^2 + m^2}.$$

64. Shew that  $\tan^{-1} \frac{x \cos \phi}{1 - x \sin \phi} - \tan^{-1} \frac{x - \sin \phi}{\cos \phi} = \phi$ .

65. If the tangents of the half angles of a triangle be in arithmetical progression, shew that the cosines of the whole angles are so also.

66. Shew that

$$\cos^{-1} \frac{1 - m^2}{1 + m^2} - \cos^{-1} \frac{1 - n^2}{1 + n^2} = \tan^{-1} \frac{2(m - n)(1 + mn)}{(1 + mn)^2 - (m - n)^2}.$$

67. If the straight line which bisects the angle  $BAC$  of a triangle divides the side  $BC$  in the ratio  $m : n$ , then

$$\tan \beta = \frac{n \sin \alpha}{m - n \cos \alpha}.$$

68. Prove that the product of the perpendiculars from the angles on the opposite sides equals

$$\frac{\{(a + b + c)r\}^2}{abc}.$$

69. If lines be drawn from the angles of a triangle to the centre of the inscribed circle, and the points where these lines meet the circle are joined so as to form a new triangle, shew that its sides are in the ratio

$$\sin \frac{\alpha}{4} + \cos \frac{\alpha}{4} : \sin \frac{\beta}{4} + \cos \frac{\beta}{4} : \sin \frac{\gamma}{4} + \cos \frac{\gamma}{4}.$$

70. Eliminate  $\theta$  and  $\phi$  from the equations

$$a \sin^2 \theta + b \cos^2 \theta = \alpha,$$

$$b \sin^2 \phi + a \cos^2 \phi = \beta,$$

$$a \tan \theta = b \tan \phi.$$

$$\text{Ans. } \frac{1}{\alpha} + \frac{1}{\beta} = \frac{1}{a} + \frac{1}{b}.$$

71. Shew that the value of  $a \cos 2\theta + b \sin 2\theta$ , when  $\tan \theta = \frac{b}{a}$ , is  $a$ .

72.  $\sin \alpha \sin \beta \sin (\alpha - \beta) + \sin \beta \sin \gamma \sin (\beta - \gamma) + \sin \gamma \sin \alpha \sin (\gamma - \alpha)$   
 $= \frac{1}{4} \{ \sin 2(\alpha - \beta) + \sin 2(\beta - \gamma) + \sin 2(\gamma - \alpha) \}.$

73. A circle is inscribed in an equilateral triangle (side  $a$ ), and another circle touches this circle and two of the sides, shew that its radius is  $\frac{a}{6\sqrt{3}}$ .

74.  $AB$  is a tower at the foot of a hill whose inclination is  $\theta$ ;  $C, D$  are two stations directly up the hill from  $B$  such that  $BC = CD$ , and  $\angle ACD = \alpha$ ,  $\angle ADC = \beta$ . Shew that  $\theta$  is found from the equation

$$\cot \theta = \frac{\sin \alpha \sin \beta}{2 \cos \alpha \sin \beta + \cos \beta \sin \alpha}.$$

75. If  $\sin \alpha$  be the arithmetic, and  $\sin \beta$  the geometric mean between  $\sin \gamma$  and  $\cos \gamma$ , prove that

$$\cos 2\alpha = \frac{1}{2} \cos 2\beta = \cos^2 (45^\circ + \gamma).$$

76. If  $\alpha, \beta, \gamma$  are each less than  $90^\circ$ , then if  $\sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma = 1$ , any two of them are together less than  $90^\circ$ , and if  $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$ , any two of them are together greater than  $90^\circ$ .

77. If  $ABC$  be a triangle, prove from the definitions that

$$\sin C = \sin A \cos B + \cos A \sin B.$$

78. A balloon considered as a vertical object of given height floats at a constant height above the earth, and subtends angles  $\alpha, \beta$  at a place when the elevations of its lowest points are  $A, B$  respectively; shew that

$$\tan (A + \alpha) \cot A = \tan (B + \beta) \cot B.$$

79. Eliminate  $\alpha$  and  $\beta$  from the equations

$$\sin \alpha \cos \beta \sin \theta + \cos \alpha \cos \theta = A,$$

$$\sin \alpha \cos \beta \cos \theta - \cos \alpha \sin \theta = B,$$

$$\sin \alpha \sin \beta \cot \theta = c.$$

$$\text{Ans. } A^2 + B^2 = 1 - c^2 \tan^2 \theta.$$

80. Find the relation between  $\alpha, b, c$  that an angle of  $\alpha$  degrees may be expressed by  $b$ , when the unit of measurement is  $\frac{\omega}{c}$ ,  $\omega$  being the unit of circular measurement.

$$\text{Ans. } \pi \alpha c = 180 b.$$

81. Eliminate
- $\theta$
- and
- $\phi$
- from the equations

$$\cos \theta \sin \alpha + \cos \phi \cos \alpha = c,$$

$$\frac{\cos (\theta + \phi)}{\cos \alpha} = \frac{\cos (\theta - \phi)}{\sin \alpha},$$

$$\frac{\cos \theta}{\sin \alpha} = \frac{\cos \phi}{\cos \alpha}.$$

$$\text{Ans. } 2(1 + \sin 2\alpha)(c^2 - 1) = c^4 \sin^3 2\alpha.$$

82. If  $A, B$  represent a certain angle in two systems whose units are the angles subtended at the centre of a circle by  $\frac{1}{m}$  th,  $\frac{1}{n}$  th parts of the circumference respectively, find the relation between  $A, B$ , and express the angle in degrees.

$$\text{Ans. } An = Bm, \text{ and angle} = \frac{A}{m} \cdot 360^\circ.$$

83. If the sides of an acute-angled triangle of given perimeter be in A.P., the common difference of the sides cannot exceed, and the radius of the inscribed circle cannot be less than  $\frac{1}{12}$  th of the perimeter.

84.  $A, B$  are two objects in the same horizontal plane,  $P$  a point at which the angle  $\alpha$  subtended by  $AB$  is observed: from  $P$  two persons walk in directions at right angles to  $PA, PB$  respectively to points  $Q, R$ , at each of which the angle subtended by  $AB$  is  $\alpha$ : if  $PQ = a, PR = b$ , find the distance  $AB$ ,  $\alpha$  being acute.

$$\text{Ans. } \sqrt{a^2 + b^2 + 2ab \cos \alpha}.$$

85. Eliminate
- $a, b, c$
- from the equations

$$a = b \cos C + c \cos B,$$

$$b = a \cos C + c \cos A,$$

$$c = a \cos B + b \cos A.$$

$$\text{Ans. } \cos A + \cos (B \pm C) = 0.$$

86. Eliminate
- $\theta$
- from the equations

$$\operatorname{cosec}^2 \theta = m \tan \theta,$$

$$\sec^2 \theta = n \cot \theta.$$

$$\text{Ans. } \sqrt{m} + \sqrt{n} = \sqrt{m^2 n^3}.$$

87. The numerical measures of the angles of a quadrilateral when referred to units  $1^\circ, 2^\circ, 3^\circ, 4^\circ$  respectively are in A.P., and the difference between the second and fourth is  $90^\circ$ ; find the angles.

$$\text{Ans. } 30^\circ, 66^\circ, 108^\circ, 156^\circ.$$

88. A person walking up a slope towards the summit of a hill observes the elevations of the summit at  $A, B, C$  to be  $\alpha, 2\alpha, 3\alpha$  respectively; and also finds that  $AB : BC = 1 : n$ . Shew that the inclination of the slope is

$$\tan^{-1} \frac{n \sin 3\alpha - \sin \alpha}{n \cos 3\alpha - \cos \alpha}.$$

89. If  $ABCD$  be a quadrilateral figure capable of having a circle inscribed in it, and also described about it, prove that the radius of the inscribed circle is equal to

$$\frac{AB + BC + CD + DA}{2 (\operatorname{cosec} A + \operatorname{cosec} B + \operatorname{cosec} C + \operatorname{cosec} D)}.$$

90. If  $A'B'C'$  be the centres of the escribed circles of a triangle  $ABC$ , and  $O$  the centre of the inscribed circle, shew that the radii of the circles circumscribing  $A'B'C'$ ,  $A'OB'$ ,  $A'OC'$ ,  $B'OC'$ , are each equal to twice the radius of the circle circumscribing the triangle  $ABC$ .

91. If in last question  $a'b'c'$  be the sides of the triangle  $A'B'C'$ , shew that

$$\frac{a'^2}{r_a + r_o} = \frac{b'^2}{r_b + r_o} = \frac{c'^2}{r_c + r_o} = \frac{(r_a + r_b)(r_a + r_c)(r_b + r_c)}{r_a r_b + r_a r_c + r_b r_c} = \frac{a'b'c'}{2\sqrt{\sigma(\sigma-a')(\sigma-b')(\sigma-c')}},$$

where  $2\sigma = a' + b' + c'$ .

$$92. \text{ If } \frac{\sqrt{\sin(\theta + 2\lambda)}}{\cos(2\theta + \lambda)} = \frac{\sqrt{(\beta^2 - \alpha^2) \sin \theta + 2\alpha\beta \cos \theta}}{\cos \theta (\beta \cos \theta - \alpha \sin \theta) - \sin \theta (\beta \sin \theta + \alpha \cos \theta)},$$

then  $\tan \lambda = \frac{\alpha}{\beta}$ .

93. In a triangle

$$\sin \alpha \sin(\alpha - \beta) \sin(\alpha - \gamma) + \sin \beta \sin(\beta - \alpha) \sin(\beta - \gamma) + \sin \gamma \sin(\gamma - \alpha) \sin(\gamma - \beta) = \sin \alpha \sin \beta \sin \gamma - \sin 2\alpha \sin 2\beta \sin 2\gamma.$$

94. Find  $\theta$  and  $\phi$  from the equations

$$p \sin^4 \theta - q \sin^4 \phi = p,$$

$$p \cos^4 \theta - q \cos^4 \phi = q.$$

$$\text{Ans. } \cos \theta = \sqrt[4]{\frac{q^2}{p(q-p)}}, \quad \cos \phi = \sqrt[4]{\frac{p}{q-p}}.$$

95. If  $\alpha, \beta, \gamma$  be unequal and each less than  $\pm 2\pi$ , and if

$$\cos \alpha + \cos \beta + \cos \gamma = \sin \alpha + \sin \beta + \sin \gamma = 0,$$

then

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = \sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma = \frac{3}{2}.$$

$$96. \text{ If } \frac{\cos \frac{\beta}{2} \cos \frac{\gamma}{2}}{\cos \frac{\beta - \gamma}{2}}, \frac{\cos \frac{\alpha}{2} \cos \frac{\gamma}{2}}{\cos \frac{\alpha - \gamma}{2}}, \frac{\cos \frac{\alpha}{2} \cos \frac{\beta}{2}}{\cos \frac{\alpha - \beta}{2}} \text{ are in A.P., so also are}$$

$\cos 2\alpha, \cos 2\beta, \cos 2\gamma$ ,  $\alpha, \beta, \gamma$  being the angles of a triangle.

97. If in a triangle  $ABC$  lines be drawn through the angular points inclined at an angle  $\alpha$  to the three sides respectively, and  $A'B'C'$  be the triangle formed by their intersections, shew that area of  $A'B'C'$  : area of  $ABC = \sin^2(\phi - \alpha) : \sin^2 \phi$ , where  $\cot \phi = \cot A + \cot B + \cot C$ .

98. If  $(1 + \sin \theta)(1 + \sin \phi)(1 + \sin \psi) = \cos^3 \theta \cos \phi \cos \psi$ ,  
 shew that  $\sin \theta + \sin \phi + \sin \psi + \sin \theta \sin \phi \sin \psi = 0$ ,  
 and  $\sec^2 \theta + \sec^2 \phi + \sec^2 \psi - 2 \sec \theta \sec \phi \sec \psi = 1$ .

99. The four common tangents drawn to two circles (radii  $r, r'$ ) form a quadrilateral. If  $2\alpha, 2\alpha'$  be the inclinations of the pairs of tangents which are similarly situated, shew that the area of the quadrilateral is  $rr'(\cot \alpha - \cot \alpha')$ .

100. Shew from a figure that  
 $\tan 80^\circ = \sqrt{3 + 4 \cos 10^\circ}$ .

101. If a circle be drawn touching the sides  $AB, AC$  of a triangle  $ABC$  produced, and the circumscribed circle, shew that its radius is equal to

$$4R \sin \frac{\alpha}{2} \cos \frac{\beta}{2} \cos \frac{\gamma}{2} \sec^2 \frac{\alpha}{2}.$$

102. Eliminate  $\theta$  from the equations

$$\frac{x}{a \sin \theta} + \frac{y}{b \cos \theta} = 1, \quad \frac{x}{a \sin^3 \theta} + \frac{y}{b \cos^3 \theta} = 0.$$

*Ans.*  $\left\{ \left( \frac{x}{a} \right)^{\frac{2}{3}} - \left( \frac{y}{b} \right)^{\frac{2}{3}} \right\} \left\{ \left( \frac{x}{a} \right)^{\frac{2}{3}} + \left( \frac{y}{b} \right)^{\frac{2}{3}} \right\}^{\frac{1}{2}} = 1.$

103. If in a triangle  $ABC$ ,  $PD$  be drawn from a point  $P$  in  $AC$  inclined to  $AC$  at an angle  $\theta$ , and meeting  $AB$  in  $D$ , so as to bisect the triangle, then

$$\cot \theta = \frac{2AP^2}{bc} \operatorname{cosec} A - \cot A.$$

104. If  $ABCD$  be a quadrilateral inscribed in a circle, and  $O, P$  the middle points of  $AB, CD$ , and if

$$POA = \theta, OPC = \phi, AB = 2a, CD = 2b, PO = c,$$

then

$$\frac{\tan \theta}{\tan \phi} = \frac{c^2 - a^2 + b^2}{c^2 + a^2 - b^2}.$$

105. If  $\frac{\tan(x+y)}{a+b} = \frac{\tan(y+z)}{b+c} = \frac{\tan(x+z)}{a+c},$

and  $\frac{\tan(x+y)}{a'+b'} = \frac{\tan(y+z)}{b'+c'} = \frac{\tan(x+z)}{a'+c'},$

then

$$\frac{b^2 - a^2}{b^2 - c^2} = \frac{b'^2 - a'^2}{b'^2 - c'^2}.$$

106. Eliminate  $\theta$  from the equations

$$\frac{x}{a} = \cos \theta + \cos 2\theta, \quad \frac{y}{b} = \sin \theta + \sin 2\theta.$$

*Ans.*  $\left(1 - \frac{x^2}{a^2} - \frac{y^2}{b^2}\right)^3 = \left(1 + \frac{x}{a}\right)^3 + \frac{y^3}{b^3}.$

107.  $A, B, C$  is a triangle;  $A', B', C'$  the middle points of the arcs  $BC, AC, AB$  of the circumscribing circle. If chord  $B'C'$  intersects  $AB, AC$  in  $L, M$ , shew that

$$AL = AM = \frac{a \sin \frac{\beta}{2} \sin \frac{\gamma}{2}}{\sin a \cos \frac{\alpha}{2}}.$$

108.  $OP$  is a vertical pole in a circular inclosure  $ADB C$ ,  $AB$  a diameter of the inclosure, and  $CD$  a chord through  $O$ . If  $3\alpha, \alpha$  be the elevations of  $P$  at  $A, B$  and  $2\theta, \theta$  at  $D, C$  respectively, shew that

$$\cos 2\theta = \cos \alpha \cos 3\alpha \sec 2\alpha.$$

109. Three lines parallel to the sides of a triangle pass through a point and cut off portions of the triangle whose areas are to the area of the triangle as  $m : 1, n : 1, p : 1$  respectively: shew that

$$\sqrt{m} + \sqrt{n} + \sqrt{p} = 2.$$

110. The area of the triangle formed by joining the centres of the escribed circles is

$$\frac{1}{2} S \operatorname{cosec} \frac{\alpha}{2} \operatorname{cosec} \frac{\beta}{2} \operatorname{cosec} \frac{\gamma}{2}.$$

111. Shew that

$$\tan \alpha (\tan 2\alpha)^{\frac{1}{2}} (\tan 4\alpha)^{\frac{1}{4}} = \frac{4 \sin^2 \alpha}{(2 \sin 8\alpha)^{\frac{1}{4}}}.$$

112. Find all the values of  $x$  which satisfy the equation

$$\operatorname{vers}^{-1}(1+x) - \operatorname{vers}^{-1}(1-x) = \tan^{-1} 2 \sqrt{1-x^2}.$$

$$\text{Ans. } \pm 1, \frac{1}{2}.$$

113. In the three edges of a cube which meet at one angle, three points  $A, B, C$  are taken at distances  $a, b, c$  from that angle respectively. The area of the triangle  $ABC$  is

$$\frac{1}{2} \sqrt{(a^2 b^2 + a^2 c^2 + b^2 c^2)}.$$

114. Determine  $x$  from the equation  $\tan(\alpha+x) = m \tan x$ , and explain the result when  $m=1$ .

115. If  $AD, AE$  be drawn bisecting the interior and exterior angles at  $A$  of a triangle  $ABC$ , and  $r_1, r_2, r_3, r_4$  be the radii of the circles circumscribing the triangles  $ABD, ACD, ABE, ACE$ , respectively, then  $r_1 r_4 = r_2 r_3$ .

116. Having given the area, base, and vertical angle of a triangle, find the other angles.

117. Having given two sides of a triangle, and the difference of the angles opposite them, determine the angles.

118. If the angle  $\alpha$  be divided into two parts so that their versed sines may be in the ratio  $m : n$ , one part is

$$2 \cot^{-1} \left\{ \cot \frac{\alpha}{2} + \sqrt{\frac{n}{m}} \operatorname{cosec} \frac{\alpha}{2} \right\}.$$

119. If  $r, r'$  be radii of the circumscribed and inscribed circles of a regular polygon of  $n$  sides, shew that

$$r + r' = a \cot \frac{\pi}{2n},$$

where  $2a$  is one of the sides.

120. Find the number of acres in a field whose sides are 400, 300, 300 yards respectively, and one angle adjacent to the largest side is a right angle.

$$\text{Ans. } \frac{125}{242} (5\sqrt{11} + 24) \text{ acres.}$$

121. In a triangle

$$\sin^2 \frac{\alpha}{2} + \sin^2 \frac{\beta}{2} + \sin^2 \frac{\gamma}{2} + 2 \sin \frac{\alpha}{2} \sin \frac{\beta}{2} \sin \frac{\gamma}{2} = 1.$$

122. Shew that  $\frac{\sin \alpha \pm \sin n\alpha + \sin (2n-1)\alpha}{\cos \alpha \pm \cos n\alpha + \cos (2n-1)\alpha} = \tan n\alpha$ .

123. If the sides of a triangle are in arithmetical progression, the perpendicular on the mean side from the opposite angle, and the radius of the circumscribed circle touching the mean side, are each three times the radius of the inscribed circle.

124. If  $a \cos^2 \theta + b \sin^2 \theta = m \cos^2 \phi$ ,

$$a \sin^2 \theta + b \cos^2 \theta = n \sin^2 \phi,$$

$$m \tan^2 \theta = n \tan^2 \phi,$$

then

$$(a+b)(m+n) = 2mn.$$

125. Solve  $8 \sin^{-1} x \sin^{-1} 2x = 3 (\sin^{-1} 2x)^2 - 3 (\sin^{-1} x)^2$ .

$$\text{Ans. } x = \pm \frac{1}{2} \text{ or } \pm \frac{\sqrt{15}}{8}.$$

126. Solve  $8 \cos \theta = \frac{\sqrt{3}}{\sin \theta} + \frac{1}{\cos \theta}$ .

$$\text{Ans. } \theta = \frac{3\pi}{2} + \frac{\pi}{12} \text{ or } n\pi + \frac{\pi}{12}.$$

127. The distance between the centres of two wheels is  $a$ , and the sum of their radii is  $c$ ; shew that the length of a string which wraps round them and crosses between them is

$$2\sqrt{(a^2 - c^2)} + c\left(\pi + 2\sin^{-1}\frac{c}{a}\right).$$

128. An object  $a$  feet high at the top of a tower subtends a given angle  $\alpha$  at a distance  $d$  from the foot of the tower. The tower's height is

$$-\frac{a}{2} \pm \frac{\sqrt{a^2 + 4d(a \cot \alpha - d)}}{2}.$$

Explain the negative result.

129. In a triangle, if  $a, b, c$  be in arithmetical progression, then

$$\sin\left(\alpha + \frac{\beta}{2}\right) = 2 \sin \frac{\beta}{2}.$$

130. Shew that  $\cot \gamma = \frac{b}{c} \operatorname{cosec} \alpha - \cot \alpha$ , and adapt the result to logarithmic computation.

131. Shew that  $\tan \theta \tan (60 + \theta) \tan (60 - \theta) = \tan 3\theta$ .

132. If  $\frac{\sin(x + a)}{\cos(x - a)} = \frac{1 - m}{1 + m}$ , then

$$\tan(45^\circ - x) = m \cot(45^\circ - a).$$

133. The hypotenuse  $AB$  of a right-angled triangle  $ACB$  is divided in  $D$  so that  $AD : BD = CB : CA$ . Shew that

$$\tan \angle ACD = \frac{a^2}{b^2}, \quad CD = \frac{\sqrt{(a^4 + b^4)}}{a + b}.$$

134. Eliminate  $\theta$  from the equations

$$(a + b) \tan(\theta - \phi) = (a - b) \tan(\theta + \phi),$$

$$a \cos 2\phi + b \cos 2\theta = c.$$

$$\text{Ans. } b^2 = a^2 + c^2 - 2ac \cos 2\phi.$$

135. If the lines drawn from the angles of a triangle perpendicular to the opposite sides meet in a point whose distances from the three angles are  $p, q, r$ , respectively, then the area of the triangle is

$$\frac{1}{4}(pa + qb + rc).$$

Shew also that  $a^2p \operatorname{cosec} \alpha + b^2q \operatorname{cosec} \beta + c^2r \operatorname{cosec} \gamma = 2abc$ .



136. The area of the triangle formed by joining the centres of ~~the~~ escribed circles is  $\frac{abc\sqrt{s}}{2\sqrt{\{(s-a)(s-b)(s-c)\}}}$ .

137. Shew that

$$\sec^2(\alpha + 45^\circ) - \sec^2(\alpha - 45^\circ) = 4 \tan 2\alpha \sec 2\alpha.$$

138. If  $\cos(\sigma - 2\alpha) + \cos(\sigma - 2\beta) = \cos(\sigma - 2\gamma) + \cos(\sigma - 2\delta)$ , where

$$\sigma = \alpha + \beta + \gamma + \delta,$$

then  $\tan \alpha \tan \beta = \tan \gamma \tan \delta$ .

139. One angle of a triangle is  $60^\circ$ , and the sides including it are in ~~the~~ ratio 5 : 3. The other angles are

$$\tan^{-1} \frac{3\sqrt{3}}{7} \text{ and } \tan^{-1} 5\sqrt{3}.$$

140. If the sides of a triangle  $ABC$  be divided into parts at  $D, E, F$ , which have to one another the same ratio  $n : 1$ , shew that if the points ~~is~~  $D, E, F$  be joined, the triangles  $ADF, BDE, CEF$  are all equal, and that

$$\triangle DEF : \triangle ABC = n^2 - n + 1 : (n + 1)^2.$$

141. If  $\cos \alpha = \cos \beta \cos \phi = \cos \beta' \cos \phi'$ ,

$$\text{and } \sin \alpha = 2 \sin \frac{\phi}{2} \sin \frac{\phi'}{2},$$

shew that  $\tan \frac{\alpha}{2} = \tan \frac{\beta}{2} \tan \frac{\beta'}{2}$ .

142. A person walking along a straight road observes that the greatest ~~at~~ angle which two objects make with each other is  $\alpha$ : from the point where ~~the~~ this happens he walks  $a$  yards, and the objects there appear in the same ~~the~~ straight line making an angle  $\beta$  with the road: the distance between the ~~the~~ objects is

$$\frac{2a \sin \alpha \sin \beta}{\cos \alpha + \cos \beta}.$$

143. In any quadrilateral

$$\frac{\tan \alpha + \tan \beta + \tan \gamma + \tan \delta}{\cot \alpha + \cot \beta + \cot \gamma + \cot \delta} = \tan \alpha \tan \beta \tan \gamma \tan \delta.$$

144. Shew that

$$2 \tan^{-1} \left[ \tan \frac{\alpha}{2} \sqrt{\left\{ \tan \left( \frac{\pi}{4} - \beta \right) \right\}} \right] = \cos^{-1} \left( \frac{\cos \alpha + \tan \beta}{1 + \cos \alpha \tan \beta} \right).$$

145. A circle is described about a triangle  $ABC$ , and a new triangle is formed by joining the points of bisection of the arcs subtended by the sides of  $ABC$ ; shew that the angles of this triangle are

$$90 - \frac{\alpha}{2}, \quad 90 - \frac{\beta}{2}, \quad 90 - \frac{\gamma}{2};$$

and the sides are

$$\frac{a}{2 \sin \frac{\alpha}{2}}, \quad \frac{b}{2 \sin \frac{\beta}{2}}, \quad \frac{c}{2 \sin \frac{\gamma}{2}};$$

and its area : area of triangle  $ABC = 1 : 8 \sin \frac{\alpha}{2} \sin \frac{\beta}{2} \sin \frac{\gamma}{2}$ .

146. Three circles whose radii are  $a, b, c$  touch each other externally; prove that the tangents at the points of contact meet in a point whose distance from any one of them is

$$\sqrt{\left(\frac{abc}{a+b+c}\right)}.$$

147. If  $a = \operatorname{cosec} \theta \tan^3 \theta (\operatorname{cosec}^2 \theta + 1) + \sec \theta$ ,

and  $b = \tan^3 \theta (\operatorname{cosec}^2 \theta + 1) - \tan \theta$ ,

then  $a^3 - b^3 = 2^3$ .

148. A person wishing to determine the length of an inaccessible wall places himself due south of one end, and due west of the other, at such distances that the angles which the wall subtends at the two positions are each equal to  $\alpha$ . If  $a$  be the distance between the two positions, the length of the wall is  $a \tan \alpha$ .

149. A vessel observed another vessel  $\alpha^\circ$  from the north sailing in a direction parallel to its own. After an hour's sailing its bearing was  $\beta^\circ$ , and after another hour  $\gamma^\circ$ . In what direction were the vessels sailing?

$$\text{Ans. } \frac{\alpha + \gamma}{2} + \tan^{-1} \left\{ \tan^2 \frac{\gamma - \alpha}{2} \cot \left( \beta - \frac{\alpha + \gamma}{2} \right) \right\},$$

$$\text{or otherwise, } \tan^{-1} \frac{\tan \beta (\tan \alpha + \tan \gamma) - 2 \tan \alpha \tan \gamma}{2 \tan \beta - (\tan \alpha + \tan \gamma)}.$$

150. A circle is inscribed in a triangle and tangents drawn parallel to each side and intercepted by the other two. If  $l, m, n$  be the lengths of the tangents opposite the sides  $a, b, c$  respectively, then

$$\frac{l}{a} + \frac{m}{b} + \frac{n}{c} = 1.$$

151. A circle is inscribed in a triangle; similar triangles are cut off by tangents to the circle, and in these triangles are inscribed other circles; shew that the radius of the first is equal to the sum of the radii of the others.

152. Two squares may be inscribed in a right-angled triangle, one having two sides lying on the sides of the triangle, and the other having one side on the hypotenuse; shew that the latter is always the least.

153. The distance between the centres of the inscribed and circumscribed circles is  $\sqrt{R^2 - 2Rr}$ .

154. In the triangle  $ABC$ ,  $B'C'$  is drawn through  $A$  parallel to  $BC$ , and  $A'B'$  through  $C$  perpendicular to  $AC$ , and  $A'C'$  through  $B$  perpendicular to  $AB$ ; shew that the area of the triangle  $A'B'C'$  is

$$\frac{a^2}{2} \cdot \frac{\cos^2(\beta - \gamma)}{\cos \beta \cos \gamma \sin \alpha}.$$

155. A person walks  $x$  yards from  $A$  to  $E$  along the side  $AB$  of the triangle  $ABC$ , and observes the angle  $AEC = \theta$ . Again he walks  $y$  yards along  $BA$  from  $B$  to  $F$  and observes  $CFB = \theta$ . Given  $AB = c$ , find the other sides of the triangle.

$$\begin{aligned} \text{Ans. } & \sqrt{\left\{ \left( \frac{x+y-c}{2} \sec \theta \right)^2 + (c-y)x \right\}}, \\ & \text{and } \sqrt{\left\{ \left( \frac{x+y-c}{2} \sec \theta \right)^2 + (c-x)y \right\}}. \end{aligned}$$

156. In the hypotenuse  $AB$  of a right-angled isosceles triangle  $ACB$ ,  $D$  is taken so that  $CD$  is equal to  $\frac{5}{7}CB$ . Shew that  $AB$  is divided in  $D$  in the ratio 3 : 4.

$$\begin{aligned} 157. \text{ If } & c \sin \theta - a \sin(\theta + \phi) = 0, \\ & a \sin \phi - b \sin \theta = 0, \\ & \cos \theta - \cos \phi = 2m, \end{aligned}$$

$$\text{then } (a-b)\{(a+b)^2 - c^2\} + 4abmc = 0.$$

158. From the summits of two rocks  $A$ ,  $B$  at sea, at a given distance  $d$  apart, the dips  $\alpha$ ,  $\beta$  of the horizon are observed, and it is remarked that the summit of  $B$  is in a horizontal line through the summit of  $A$ ; shew that the radius of the Earth is

$$\frac{d}{\cos^{-1}(\sec \alpha \cos \beta)}.$$

159. Of the three squares which can be described in a triangle so that one side of the square coincides with part of one side of the triangle, the greatest is that which is on the least side.

160. If the perpendiculars from the angular points of triangle  $ABC$  meet the sides in  $D, E, F$ , and  $R, R_1$  be the radii of the circles described about triangles  $ABC, DEF$ , and  $r_1$  the circle inscribed in  $DEF$ , prove that

$$R_1 = \frac{1}{2}R \text{ and } r_1 = 2R \cos \alpha \cos \beta \cos \gamma.$$

161. In a triangle

$$a^2 : b^2 : c^2 = \cot \beta + \cot \gamma : \cot \gamma + \cot \alpha : \cot \alpha + \cot \beta.$$

162. A person walking along a straight road observes the greatest elevation of a tower to be  $\alpha$ . From another straight road he observes the greatest elevation of the tower to be  $\beta$ . The distances of the points of observation from the intersection of the two roads are  $a, b$  respectively: prove that the height of the tower is

$$\left( \frac{a^2 - b^2}{\cot^2 \beta - \cot^2 \alpha} \right)^{\frac{1}{2}}.$$

163. If the inscribed circle touch the sides of the triangle  $ABC$  in points  $A'B'C'$ , and  $R_1, R_2, R_3$  be the radii of the circles circumscribed about  $\triangle B'C', \triangle A'C', \triangle A'B'$  respectively, then  $2R_1 R_2 R_3 = r^2 R$ .

164. If in a triangle  $ABC$  circles are described with centres  $B, C$  to touch the opposite sides and intersect in  $A'$ , then

$$\cos BA'C = \frac{\cos \alpha \cos (\beta - \gamma)}{2 \sin \beta \sin \gamma}.$$

165. Prove that

$$\tan^{-1} \frac{m+n}{m-n} - \tan^{-1} \frac{m-n}{m+n} = \frac{1}{2} \sin^{-1} \frac{4mn(m^2 - n^2)}{(m^2 + n^2)^2}.$$

166. Eliminate  $\theta$  from the equations

$$a \sin 3\theta - b \cos 3\theta = c \sin 2\theta \sqrt{(\cos 2\theta)},$$

$$a \cos 3\theta + b \sin 3\theta = c \cos 2\theta \sqrt{(\cos 2\theta)}.$$

$$\text{Ans. } (a^2 + b^2)^2 = c^2 (a^2 - b^2).$$

167. If  $(x^2 + y^2) \sin^2 \alpha = (y \sin \beta - x \cos \beta)^2$ , then  $x = y \tan (\beta \pm \alpha)$ .

168. If  $x \cos \theta = b, x \cos (2\alpha - \theta) = c$ ,

$$\text{then } x = \frac{1}{2} \{ (b+c)^2 \sec^2 \alpha + (b-c)^2 \operatorname{cosec}^2 \alpha \}^{\frac{1}{2}}.$$

169. If  $\sin \theta = \sin \phi + \sin \psi$ ,  $\cos \phi = \cos \psi + \cos \theta$ ,  $\cot \psi = \cot \theta + \cot \phi$ , find  $\theta, \phi, \psi$ .  
*Ans.*  $\theta = 150^\circ$ ,  $\phi = 90^\circ$ ,  $\psi = -30^\circ$ .

170. If  $O$  be the centre of the circle inscribed in the triangle  $ABC$ ;  $D, E, F$  the points of contact, and  $\rho_1, \rho_2, \rho_3$  the radii of circles inscribed in  $AE OF, BDOF, CDOE$ , then

$$r^2 (\rho_1 + \rho_2 + \rho_3) - 2r (\rho_1 \rho_2 + \rho_1 \rho_3 + \rho_2 \rho_3) + 2\rho_1 \rho_2 \rho_3 = 0.$$

171. Prove that the tangents of the angles of depression of one balloon observed from another at equal intervals of time are in A.P., if the two balloons are under the influence of the same horizontal currents of air, and are moving in straight lines.

172. Solve the equation

$$\cot \theta + \cot \left( \theta + \frac{\pi}{3} \right) + \cot \left( \theta + \frac{2\pi}{3} \right) = 3 \cot 3\alpha.$$

$$\text{Ans. } \theta = \frac{n\pi}{3} + \alpha.$$

173. Shew that

$$\operatorname{cosec}^2 \alpha + \operatorname{cosec}^2 \left( \alpha + \frac{\pi}{3} \right) + \operatorname{cosec}^2 \left( \alpha + \frac{2\pi}{3} \right) = 9 \operatorname{cosec}^2 3\alpha.$$

174.  $AB$  is a diameter of a circle produced to any point  $P$ , and  $AC, AD$  are chords of the circle. If  $BC, BD, PC, PD$  are joined, shew that

$$\tan PCB : \tan PDB = \cot BAC : \cot BAD.$$

175. In a triangle, shew that

$$8 \sin \frac{\alpha}{2} \sin \frac{\beta}{2} \sin \frac{\gamma}{2} < 1,$$

unless  $\alpha = \beta = \gamma$ . Hence shew that if  $R = 2r$ , the triangle is equilateral.

176. If  $D, E, F$  are the feet of the perpendiculars from the angles  $A, B, C$  of a triangle on the opposite sides, and  $D', E', F'$  the points of contact of the inscribed circle, shew that

$$DD' \sin \alpha + EE' \sin \beta + FF' \sin \gamma = 0,$$

if  $DD', EE', FF'$  have the same signs as  $b - c, c - a, a - b$  respectively.

177. In a triangle  $ABC$ , if  $D, E, F$  be the points of contact of the inscribed circle, and  $AD, BE, CF$  meet the circle in  $D', E', F'$ , shew that

$$\frac{AE^2}{AD^2} + \frac{BF^2}{BE^2} + \frac{CD^2}{CF^2} = 3 + \frac{4}{s} \left\{ \frac{r_a^2}{a} + \frac{r_b^2}{b} + \frac{r_c^2}{c} \right\}.$$

178. If  $\alpha + \beta + \gamma + \delta = 860^\circ$ , and  $\tan \alpha, \tan \beta, \tan \gamma, \tan \delta$  be in G.P., then  $\tan \alpha \tan \delta = \tan \beta \tan \gamma = 1$ , and if  $\tan \alpha = a$ , find the other angles.

$$\text{Ans. } \tan \beta = \sqrt[3]{a}, \tan \gamma = \sqrt[3]{a^{-1}}, \tan \delta = a^{-1}.$$

179. If 
$$\frac{6 \sin \theta}{\cos (\phi + \theta)} = \frac{3 \sin 2\theta}{\cos (\phi + 2\theta)} = \frac{2 \sin 3\theta}{\cos (\phi + 3\theta)},$$

then  $\theta = 0$ .

180. In a triangle

$$\tan 2\alpha + \tan 2\beta + \tan 2\gamma = -\frac{\sin 4\alpha + \sin 4\beta + \sin 4\gamma}{1 + \cos 4\alpha + \cos 4\beta + \cos 4\gamma}.$$

181. In a triangle

$$1 + \cos C < 2 \left( \cos \frac{A-B}{2} + \sin \frac{A+B}{2} \right).$$

182.  $ABCD$  is a quadrilateral figure inscribed in a circle whose sides are in *c.p.* (common ratio  $r$ ): shew that

$$\tan ABC : \tan BCD = r^2 - 1 : r^2 + 1.$$

183. If  $AD$ ,  $BC$ ,  $CF$  bisect the angles of a triangle  $ABC$ , shew that triangle  $DEF$  : triangle  $ABC = 2abc : (a+b)(a+c)(b+c)$ .

184. If lines be drawn from the angular points  $A$ ,  $B$ ,  $C$  of a triangle through  $O$  the centre of the inscribed circle to meet the circumference of the circumscribed circle in  $P$ ,  $Q$ ,  $R$  respectively; shew that the product of the radii of the circles circumscribed about  $BOP$ ,  $ROA$ ,  $COQ$  equals the product of those about  $BOR$ ,  $AOQ$ ,  $COP$ , and also equals

$$\frac{a^2 b^2 c^2}{r^2 (a+b+c)^4}.$$

185. If  $p_1$ ,  $p_2$ ,  $p_3$  be the perpendiculars from the angles on the opposite sides of a triangle, and  $q_1$ ,  $q_2$ ,  $q_3$  the perpendiculars from the centre of the circumscribed circle upon the same sides respectively, and  $r$  the radius of the inscribed circle, then

$$\frac{r - q_1}{p_1} + \frac{r - q_2}{p_2} + \frac{r - q_3}{p_3} = 0.$$

186. A triangle is described about a given circle, and three different similar triangles are escribed about it. Prove that each side of the first triangle is equal to the sum of the corresponding sides of the escribed triangles.

187. If  $O$  be the centre of the escribed circle touching the side  $BC$  of the triangle  $ABC$ , and  $BO$ ,  $CO$  be produced to meet  $AC$ ,  $AB$  produced in  $F$ ,  $E$ , then

$$\frac{OE}{OF} = \frac{\sin \frac{\gamma - \alpha}{2}}{\sin \frac{\beta - \alpha}{2}}.$$

188. If  $p, q, t$  be the distances between the centres of the escribed circles of a triangle, then

$$\frac{r_a r_b r_c}{pqt} = \frac{1}{8} \sin \alpha \sin \beta \sin \gamma.$$

Shew also that area of triangle is equal to

$$r_a r_b \sqrt{\frac{t^2}{(r_a + r_b)^2} - 1}.$$

189. Prove that  $(1 + \sec \alpha)(1 + \sec 2\alpha)(1 + \sec 4\alpha) \dots$  to  $n$  factors is equal to

$$\frac{\tan 2^{n-1} \alpha}{\tan \frac{\alpha}{2}}.$$

190. If  $A, A'$  are the peaks of two mountains, and  $BC$  a straight horizontal road; shew that the nearer of the peaks will just conceal the other at some point of the road, if  $\frac{\sin \alpha}{\sin \beta} = \frac{\sin \alpha'}{\sin \beta'}$ , where  $\alpha$  is the altitude of  $A$ , as seen from any point  $B$  of the road,  $\beta$  the angle  $ABC$ , and  $\alpha', \beta'$  similar angles for the peak  $A'$  as seen from any point  $B'$  of the road.

191.  $ADB$  is a semicircle;  $AB$  ( $2a$ ) its diameter;  $CD$  any radius dividing the semicircle into two sectors in each of which is inscribed a circle: these circles touch  $AB$  in  $M$  and  $N$ . Prove that if  $MN = b$ , and  $r, r'$  be the radii of the two circles,

$$(a - b)^2 = (a - 2r)(a - 2r') = \left(\frac{rr'}{a}\right)^2.$$

192. If

$$a^2 = y^2 + z^2 + 2yz \cos \theta,$$

$$b^2 = z^2 + x^2 + 2zx \cos \phi,$$

$$c^2 = x^2 + y^2 + 2xy \cos \chi,$$

and  $\theta + \phi + \chi = \pi$ , prove that

$$2yz \sin \theta + 2zx \sin \phi + 2xy \sin \chi = \sqrt{(a+b+c)(a+b-c)(a-b+c)(b+c-a)}.$$

193. If the sides of a triangle  $ABC$  are divided in points  $A', B', C'$  so that

$$BA' : A'C = CB' : B'A = AC' : C'B = m : n; \text{ where } m + n = 1,$$

and if  $AA', BB', CC'$  be joined and a triangle formed whose sides are equal to these three lines, then its area equals  $(1 - mn)$  times the area of  $ABC$ .

194. A man walks directly towards an object  $A$  until two other objects  $B, C$  subtend the greatest angle in his path, when he observes that  $A$  and  $B$  appear under the same angle ( $\alpha$ ) as  $B$  and  $C$ . He then walks directly towards  $C$  until  $A$  and  $B$  subtend the greatest angle in his new path, when he once more observes that  $A$  and  $B$  appear under the same angle ( $\beta$ ) as  $B$  and  $C$ . If the length of his walk between the two observations be  $d$ , shew that  $2\beta = 90^\circ + \alpha$ , and find the distances  $AB, BC, AC$ .

$$\text{Ans. } AB = d \sin \alpha \sec \beta, \quad BC = d \tan \beta,$$

$$AC = d \sqrt{1 + 4 \sin \alpha + 12 \sin^2 \alpha + 8 \sin^4 \alpha}.$$

195. The sun's rays supposed parallel, cast a shadow throughout the day on a horizontal table of a sphere resting on the table. The sun's altitude is  $\alpha$  when it is S.S.E., and  $\beta$  when it is W.S.W.: prove that if  $D$  be the distance between the two points of the shadows farthest from, and  $d$  the distance of the two points nearest to, the point of contact of the sphere,

$$d = D \tan \frac{\alpha}{2} \tan \frac{\beta}{2}.$$

196. Solve the equation

$$\cos \theta + \frac{1}{2} \operatorname{cosec}^2 \frac{\theta}{2} - \frac{2 \sin^2 \frac{\theta}{2}}{\cos \theta} = \cos \alpha + \frac{1}{2} \operatorname{cosec}^2 \frac{\alpha}{2} - \frac{2 \sin^2 \alpha}{\cos \alpha}.$$

$$\text{Ans. } \sin \frac{\alpha + \theta}{2} = 0, \quad \sin \frac{\alpha - \theta}{2} = 0, \quad \cos \theta = \frac{1}{1 - \cos \alpha} \text{ or } \frac{\cos \alpha - 1}{\cos \alpha}.$$

197. Two circles (radii  $R, r$ ) touch each other internally. A tangent to the smaller circle is a chord of the larger, and is inclined at an angle  $\theta$  to their common diameter, shew that its length is

$$2\sqrt{(R-r)(1+\sin \theta)\{R(1-\sin \theta)+r(1+\sin \theta)\}}.$$

198.  $O$  is the centre of the inscribed circle in the triangle  $ABC$ ;  $P, Q, R$  are the centres of the circles inscribed in  $BOC, COA, AOB$  respectively: prove that the area of the triangle  $PQR$ : the area of  $ABC$  ::

$$\sin \frac{A}{4} \sin \frac{B}{4} \sin \frac{C}{4} \left( \tan \frac{A}{4} + \tan \frac{B}{4} + \tan \frac{C}{4} + 3 \right) : \sqrt{2} \cos \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2}.$$

199.  $O$  and  $E$  are the centres of the circles inscribed in and circumscribed about the triangle  $ABC$ , and  $D$  is the intersection of the perpendiculars from  $A, B, C$  on the opposite sides; prove that the area of the triangle

$$DEO : \text{area of } ABC :: \sin \frac{C-B}{2} \sin \frac{A-C}{2} \sin \frac{B-A}{2} : \sin A \sin B \sin C.$$

200. The inscribed circle of a triangle passes through the centre of the circumscribing circle, and also through the point of intersection of the perpendiculars from the angles on the opposite sides, prove that the angles of the triangle are

$$\frac{\pi}{3}, \quad \frac{\pi}{3} + \cos^{-1} \left( \sqrt{2} - \frac{1}{2} \right), \quad \frac{\pi}{3} - \cos^{-1} \left( \sqrt{2} - \frac{1}{2} \right).$$

201. The tops of three mountains  $B, C, D$  are in the same plane with the station  $A$ .  $D$  is invisible from  $A$ , but the angles of elevation of  $B$  and  $C$  are observed to be  $\theta$  and  $\phi$ , and the angle  $BAC$  to be  $\alpha$ . From  $B$  the angles  $\angle ABD, CBD$  are observed to be  $\beta$  and  $\alpha$  respectively, and from  $C$  the angle  $\angle CD$  is found to be  $\beta$ ; prove that  $D$  is at a height above  $A$  equal to

$$\frac{\alpha \sin \beta (2 \cos \alpha \sin \phi - \sin \theta)}{\sin (2\alpha + \beta)},$$

where  $\angle B = \alpha$ .



202. In a triangle if

$$y \sin^2 \alpha + x \sin^2 \beta = z \sin^2 \beta + y \sin^2 \gamma = x \sin^2 \gamma + z \sin^2 \alpha,$$

then

$$x : y : z = \sin 2\alpha : \sin 2\beta : \sin 2\gamma.$$

203. In a triangle

$$\sin\left(\gamma + \frac{\alpha}{2}\right) + \sin\left(\beta + \frac{\gamma}{2}\right) + \sin\left(\alpha + \frac{\beta}{2}\right) + 1 = 4 \cos \frac{\alpha - \beta}{4} \cos \frac{\alpha - \gamma}{4} \cos \frac{\beta - \gamma}{4}.$$

204. In a triangle

$$\sin^2 \frac{\alpha}{2} + \sin^2 \frac{\beta}{2} + \sin^2 \frac{\gamma}{2}$$

cannot be less than  $\frac{3}{4}$ .

205. If

$$x + y \cos \gamma + z \cos \beta = \cos(\sigma - \alpha),$$

$$y + z \cos \alpha + x \cos \gamma = \cos(\sigma - \beta),$$

$$z + x \cos \beta + y \cos \alpha = \cos(\sigma - \gamma),$$

then

$$\frac{x}{\sin \alpha} = \frac{y}{\sin \beta} = \frac{z}{\sin \gamma} = \frac{1}{2 \sin \sigma}$$

where  $2\sigma = \alpha + \beta + \gamma$ .

206. If

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1,$$

then  $(\sin 2\alpha + \sin 2\beta + \sin 2\gamma)^2 + 4\{\sin^2(\beta - \gamma) + \sin^2(\alpha - \beta) + \sin^2(\alpha - \gamma)\} = 8$ .

207. Prove that the area of any triangle is a mean proportional between the areas of two triangles formed, the one by joining the points of contact of any one of the four circles touching the sides, and the other by joining the centres of the other three.

208. A plane polygon, whose sides are  $a_1, a_2, a_3, \dots$  and area  $A$ , is divided into triangles by joining its angular points with a point at which the sides subtend angles  $\theta_1, \theta_2, \dots$ . If the centres of the circumscribing circles of these triangles be joined in order, the area of the polygon thus formed is

$$\frac{1}{2} A - \frac{1}{8} (a_1^2 \cot \theta_1 + a_2^2 \cot \theta_2 + \dots).$$

209. The angles subtended by one tower at the base and summit of another tower of height  $a$  on the same horizontal plane, are respectively  $\alpha$  and  $\beta$ . Prove that the height of the first tower is given by the equation

$$x^2 \cos \alpha \sin(\beta - \alpha) - ax \sin^2 \alpha \sin \beta + a^2 \sin^2 \alpha \sin \beta = 0.$$

210. Two equal circles roll on the circumference of another circle, the one inside and the other outside: if when they have completed the

### MISCELLANEOUS EXAMPLES.

same number of circuits, they have made complete revolutions, number as 7 : 1, compare the radii of the circles.

When the exterior circle has performed half a circuit, what interior one be so as to have revolved as much?

It has performed  $3\frac{1}{2}$  circuits.

211. If the sides  $a, b, c$  of a triangle be increased in the ratios  $l : 1, m : 1, n : 1$ , and the angle  $c$  be thereby halved and the sum of the other two doubled, then

$$(la^2 + mb^2)(l + m) = c^2(lm + n^2).$$

212. From the extremity  $B$  of the radius  $AB$  of a circle a straight line  $BC$  is placed in the circle =  $\frac{1}{2}$  radius, from  $C$  another straight line is drawn through the middle point of  $AB$  to meet the circumference in  $F$ . Shew that triangle  $BFA$  is double the triangle  $BCA$ . Also if  $FAB = \phi$ ,  $CAB = \theta$ , then

$$\operatorname{cosec} \phi + \cot \theta + 3 \cot (\phi + \theta) = 0.$$

213. A rod is moved towards the eye of an observer so that its middle point and upper extremity are always viewed in the same directions: if at two positions when its upper and lower halves subtend angles at the eye which are as 1 : 1, 1 : 2, the distances of its middle point from the eye are  $a\sqrt{5}, a$ : find the length of the rod. Ans.  $2a\sqrt{5}$ .

214. At the extremity of an arc ( $AB$ ) of a circle (radius  $r$ ) subtending an angle  $\theta$ , a tangent  $BD$  is drawn in the direction of angular movement, whose length is  $r \tan \theta$ . If  $AC$  is the diameter through  $A$ , and  $DC$  be joined making angle  $DCA = \phi$ , then if when  $\theta$  is doubled  $\phi$  becomes  $\phi'$ , shew that

$$\frac{\tan \phi}{\tan \phi'} = \frac{\cos 3\theta}{\cos \theta + \cos 2\theta} \cdot \frac{\cos \theta}{\cos 2\theta}.$$

215. A trefoil consisting of three equal semicircles is inscribed symmetrically within another such trefoil: shew that their areas are in the same ratio as the radii of the circles inscribed in the smaller and circumscribing the greater trefoil.

216. A person wishing to calculate the radius of a railway curve, measures a distance  $a$  from one of the telegraph posts at right angles to the curve at that point: he then observes that the  $p$ th and  $q$ th posts from the one, from which he made his measurement, are in a straight line with him: the distance between two consecutive posts, measured along the curve, is  $a$ , shew that the radius is formed from the equation

$$\cos (p+q) \frac{a}{2r} = \frac{r}{r+a} \cos \frac{(q-p)a}{2r}.$$

## EXAMINATION PAPERS.

### PREVIOUS EXAMINATION. *December, 1866.*

1. WHAT is the circular measure of an angle? The ratio of the circumference of a circle to its diameter is 3.14159; express in degrees (to four places of decimals) the angle whose circular measure is unity.

2. Define the sine, cosine, and cotangent of an angle. Trace the changes in sign and magnitude of the cosine as the angle increases from  $0^\circ$  to  $360^\circ$ .

3. Investigate a general expression for all angles which have a given tangent.

4. Express the cosine of an angle in terms of the tangent only, and of the cosecant only.

Solve the equation,  $4 \sec^2 \theta - 3 \tan^2 \theta = 1$ .

5. Shew that  $\sin(A+B) = \sin A \cos B + \cos A \sin B$ , where  $A$  and  $B$  are two angles whose sum is less than  $90^\circ$ .

Given  $\sin 45^\circ = \frac{1}{\sqrt{2}}$ ,  $\sin 30^\circ = \frac{1}{2}$ , find  $\sin 15^\circ$ .

6. Shew that in every triangle any side is equal to the sum of the products obtained by multiplying each of the other sides into the cosine of the angle between it and the first side.

### *April, 1867.*

1. DESCRIBE the different ways of measuring angles.

Write down to five places of decimals the ratio which the circumference of a circle bears to its diameter.

2. An arc of a circle whose radius is 7 inches, subtends an angle of  $15^\circ.39'.7''$ : what angle will an arc of the same length subtend in a circle whose radius is 2 inches?

3. There is an angle whose tangent is twice its sine: find the length of the arc subtending it in terms of the radius.

4. Prove the formula

$$\cos(A-B) = \cos A \cos B + \sin A \sin B.$$

5. Shew that the difference of cosines of any two angles is equal to twice the product of the sine of half the sum of the angles, and the sine of half the difference.

6. If  $ABC$  be a triangle, prove that

$$\cos^2 A + \cos^2 B + \cos^2 C + 2 \cos A \cos B \cos C = 1.$$

7. Solve the equation

$$2 \sin^2 3\theta + \sin^2 6\theta = 2.$$

8. The town *C* is halfway between the towns *D* and *E*: and the towns *C*, *E* and *F* are equidistant from each other. Compare the distance of *D* from *F* with its distance from *E*.

December, 1868.

1. DEFINE an angle. How are angles measured? Compare the measures of any angles referred to the three usual unit angles. What is the circular measure of an English minute?

Express  $\frac{4}{5}$  of two right angles in each of the French and circular systems.

If the unit of measurement be  $5^\circ$ , what is the measure of  $22\frac{1}{2}^\circ$ ?

2. Define the tangent of an angle; draw the figure of an angle whose tangent is 2. What is the cosecant of this angle? Can more than one such angle be drawn? If the tangent be negative, what inference may be drawn as to the magnitude of the angle?

3. What is a supplement? What are the supplements of  $25^\circ$ ? of  $205^\circ$ ? of  $-\frac{3\pi}{4}$ ? State and prove the relations subsisting between the secants of  $A$ ,  $(90^\circ + A)$ , and of  $\theta$ ,  $\frac{\pi}{2} - \theta$ .

4. Prove Geometrically that

$$\cos 24^\circ = \cos 60^\circ \cos 36^\circ + \sin 60^\circ \sin 36^\circ.$$

Given  $\cos 36^\circ = \frac{\sqrt{5}+1}{4}$ ,  $\cos 60^\circ = \frac{1}{2}$ ; find  $\cos 24^\circ$  to three places of decimals.

5. Establish the following formulæ:

$$(1) \sec^2 x - \tan^2 x = 1.$$

$$(2) \cos 4x = \cos^2 2x - \sin^2 2x.$$

$$(3) \sin 2y + \sin 2z = 2 \sin (y+z) \cos (y-z).$$

6. Obtain an expression for the area of a triangle.

What is the area of a triangle whose base is 4 ft., and altitude  $1\frac{1}{2}$  yds.? What of a triangle whose sides are 5, 6, 5 inches respectively?

7. A target is 6 feet in height and 4 feet in breadth; find the tangents of the angles which its 4 edges subtend at a point 100 yards straight in front of its top left-hand corner.

March, 1869.

1. An angle referred to different units has measures in the ratio 8 to 5, one unit is  $2^\circ$ , what is the other? Express each unit in terms of the other.

2. Trace the changes in sign of  $\cos A$  as  $A$  changes from  $45^\circ$  to  $315^\circ$ . Illustrate by a figure.

3. Shew that  $\sin A$  is numerically greatest when  $\cos A$  is numerically least, and vice versa. When is  $\cos^4 A - \sin^4 A$  greatest?—and when least?

4. What relation have the cosine, cosecant, and cotangent of an angle to the sine, secant, tangent respectively?

Find the difference between the supplement of the complement of an angle, and the complement of its supplement.

5. Prove that  $\tan 45^\circ = 1$ , and  $\tan 60^\circ = \sqrt{3}$ . Hence find  $\tan 15^\circ$  to 3 places of decimals.

6. Prove that

$$(1) \sec^2 \alpha + \operatorname{cosec}^2 \alpha = \sec^2 \alpha \operatorname{cosec}^2 \alpha.$$

$$(2) \cot^2 \alpha - \tan^2 \alpha = \frac{4 \cos 2\alpha}{\sin^2 2\alpha}.$$

7. In any triangle obtain an expression for the cosine of each angle in terms of the sides, e.g. let the sides be 5, 7, 8. Find also the radius of the circle which circumscribes this triangle.

8. From  $G$  the centre of gravity of an equilateral triangle  $GB$  is drawn at right angles to the plane of the triangle and equal to a side; determine the angle between the lines joining  $B$  to any two corners of the triangle.

December, 1871.

1. What are the three different methods employed in measuring angles and what are their respective units?

Find an expression for an angle of a regular hexagon referred to an angle of a regular pentagon as unit.

2. Define  $\sin A$ ,  $\cot A$ ,  $\sec A$ : and express any two in terms of the third.

If  $\tan A = \frac{1}{\sqrt{8}}$ , find the value of  $\sin A$ .

3. Find an expression for  $\cos(A - B)$  in terms of the sines and cosines of  $A$  and  $B$ , when  $A$  and  $B$  are each less than  $90^\circ$ , and  $A$  greater than  $B$ .

Shew that the sum of the cosines of two angles is equal to twice the cosine of half the sum of the angles multiplied by the cosine of half the difference.

4. Find the value of  $\sin 30^\circ$ , and of  $\cos 45^\circ$ .

5. Prove the formulæ:

$$(1) \tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}.$$

$$(2) 2 \sin \frac{A}{2} = \pm \sqrt{1 + \sin A} \pm \sqrt{1 - \sin A}.$$

Given  $\sin 30^\circ$ , find  $\sin 15^\circ$ .

6. Find a general expression for all angles that have a given cosine.

If  $\sin \theta + \cos \theta = \sqrt{2}$ , find the value of  $\theta$ .

7. In any triangle, shew that the sides are proportional to the sines of the opposite angles.

Hence deduce an expression for the cosine of an angle of a triangle in terms of the sides.

The sides of a triangle are 3 in., 5 in., 7 in. Find the greatest angle.

S. JOHN'S COLLEGE. May, 1868.

1. DEFINE the Trigonometrical Ratios. Shew that

$$\sec^2 A = 1 + \tan^2 A.$$

Find the least value of  $a^2 \sec^2 x + b^2 \cos^2 x$ , where  $a$  and  $b$  are constant quantities.

2. Investigate an expression for all the angles which have a given cosine. Write down the general value of
- $x$
- which satisfies the equation

$$\sin^2 x = \sin^2 a.$$

3. Shew that

$$\sin(A+B) = \sin A \cos B + \cos A \sin B,$$

and

$$\cos(A+B) = \cos A \cos B - \sin A \sin B.$$

If  $A+B+C=180^\circ$ , shew that

$$\begin{aligned} \sin^2 A \sin 2A + \sin^2 B \sin 2B + \sin^2 C \sin 2C - \sin 2A \sin^2 B - \sin 2B \sin^2 A \\ = \sin 2A \sin 2B \sin 2C. \end{aligned}$$

4. If
- $\theta$
- be the circular measure of a positive angle less than a right angle
- $\theta$
- is greater than
- $\sin \theta$
- and less than
- $\tan \theta$
- .

Find the limit when  $\theta$  is indefinitely diminished

$$\text{of } \frac{\sin a\theta}{\sin b\theta}, \text{ and } \frac{\text{vers } a\theta}{\text{vers } b\theta}.$$

7. Investigate an expression for the area of a triangle in terms of the sides.

The sides  $a, b, c$  of a triangle are in arithmetical progression: shew that the area is to that of an equilateral triangle having the same perimeter as

$$\sqrt{1 - \frac{4(a-b)^2}{b^2}}$$

is to unity.

8. Shew that in any triangle,

$$a \sin \frac{1}{2}(B-C) = (b-c) \cos \frac{A}{2},$$

and

$$a \cos \frac{1}{2}(B-C) = (b+c) \sin \frac{A}{2}.$$

If  $b=14, c=11, A=60^\circ$ , find  $B$  and  $C$ , having given

$$\log 2, \log 3, \text{ and } \log \tan 11^\circ. 44'. 29''. 5.$$

9. Explain the notation used for inverse trigonometrical functions.

Find the value of  $\cos 4(\tan^{-1} a)$  in terms of  $a$ .

Solve the equation

$$\tan^{-1} x + \frac{1}{2} \sec^{-1} 5x = \frac{\pi}{4}.$$

May, 1869.

1. EXPLAIN what is meant by the circular measure of an angle. Why is it necessary to shew that the angle subtended at the centre of a circle by an arc equal to the radius is an invariable angle? Shew how to connect the circular measure of an angle with the measure of the same angle in degrees.

The difference of two angles is  $1^\circ$ ; the circular measure of their sum is 1: find the angles.

2. Define the Trigonometrical Ratios. Shew that  $\sin^2 A + \cos^2 A = 1$ . Find the value of  $\theta$  from the equation

$$\sin^2 \theta - 2 \cos \theta + \frac{1}{4} = 0.$$

Investigate the conditions which must hold in order that the equation

$$\sin^2 \theta + b \sin \theta + c = 0$$

may have *two* admissible roots, or may have *one*.

3. Investigate the expressions for  $\sin(A+B)$  and  $\cos(A+B)$  in terms of the sines and cosines of  $A$  and  $B$ .

Solve the following equations:

$$(1) \quad \sin 3\theta = 8 \sin^3 \theta.$$

$$(2) \quad \sin 5\theta = 16 \sin^5 \theta.$$

4. Find the sine and cosine of an angle of  $18^\circ$ .

Find the values of  $\theta$  between 0 and  $\frac{\pi}{2}$  which satisfy the equation

$$\sin^2 \theta + \cos^2 2\theta = \frac{3}{4}.$$

5. Define a logarithm. Explain the advantage of the common system of logarithms.

Find approximately the value of  $x$  from the equation

$$\left(\frac{10}{3}\right)^{x+2} = 9^{2x-1};$$

having given  $\log 3 = .4771213$ .

7. Express the cosine of an angle of a triangle in terms of the sides.

A person on the slope of a hill observes the angles of elevation  $\alpha$  and  $\beta$  of two objects on the hill; and also the angle  $\gamma$  which they subtend at his position: if  $\theta$  be the inclination of the hill to the horizon, shew that

$$\sin^2 \theta \sin^2 \gamma = \sin^2 \alpha + \sin^2 \beta - 2 \sin \alpha \sin \beta \cos \gamma.$$

8. Shew how to solve a triangle, having given two sides and the included angle.

If  $a = 210$ ,  $b = 110$ ,  $C = 84^\circ. 42'. 30''$ , find the other angles, having given

$$\log 2 = .3010300,$$

$$L \cot 17^\circ. 21'. 15'' = 10.5051500.$$

9 Find the area of a circle, and of a sector of a circle.

## CAMBRIDGE LOCAL EXAMINATIONS.

## JUNIOR STUDENTS, 1880.

1. The sine of an angle is  $\frac{1}{2}$ ; find its secant and tangent.

Draw figures to represent the four least positive angles, of which the cosine is double the sine.

2. Prove that  $\tan A = \cot (90^\circ - A)$ .

Express  $\sin 2300^\circ$ ,  $\tan 800^\circ$  in terms of trigonometrical ratios of angles less than  $45^\circ$ .

3. Prove geometrically or otherwise that

$$(i) \quad \sin (A + B) + \sin (A - B) = 2 \sin A \cos B.$$

$$(ii) \quad \tan (C + D) = \frac{\tan C + \tan D}{1 - \tan C \tan D}.$$

$$(iii) \quad \sin 2E = 2 \sin E \cos E.$$

4. Prove

$$(i) \quad (1 - 2 \cos^2 A) (\tan A + \cot A) = (\sin A - \cos A) (\sec A + \operatorname{cosec} A).$$

$$(ii) \quad \sin 3A = \sin A (2 \cos 2A - 1) \tan (60^\circ + A) \tan (60^\circ - A).$$

5. Solve the equation  $\tan 5\theta = \tan \theta$ .

6. Explain the use of logarithms in facilitating the processes of multiplication and division.

Find the following logarithms:  $\log_2 1024$ ,  $\log_{\frac{1}{2}} 125$ ,  $\log_{\frac{1}{3}} 4$ ; and the characteristics of the following:  $\log_3 250$ ,  $\log_{\frac{1}{3}} 87$ ,  $\log_{10} .001$ .

7. Prove that in any triangle of which  $A$ ,  $B$ ,  $C$  are the angles, and  $a$ ,  $b$ ,  $c$  the sides,

$$c = a \cos B + b \cos A.$$

Express the distance of the middle point of a side of a triangle from the perpendicular upon that side from the opposite angle in terms of the sides of the triangle.

8. Find the number of cubic inches of metal in a hollow right circular iron pipe, height 10 ft. 4 in., outer radius 9 in. and thickness  $1\frac{1}{2}$  in.; obtain your result accurate to a cubic inch, assuming  $\pi = 3.1416$ .

## JUNIOR STUDENTS, 1882.

1. A CERTAIN small angle was found by a Frenchman to measure 27.2 French seconds: an Englishman found it to measure 8.82 English seconds. Do the two measures agree? If not, find their difference in English seconds to three decimal places.



2. State carefully what you mean by the sine and the cosine of an obtuse angle.

Write down the values of  $\sin 90^\circ$ ,  $\sin 270^\circ$ ,  $\cos 180^\circ$ ,  $\cos 360^\circ$ ,  $\tan 450^\circ$ .

3. What is the *base* of a system of logarithms? Prove that  $\log(\sqrt[n]{x}) = \frac{1}{n} \log x$ .

Find the seventh root of .004, having given that

$$\log 2 = .30103 \text{ and } \log 45439 = 4.65743.$$

4. Prove the following identities:

$$(i) \quad \sin^2 A + \sin^2 B = 1 - \cos(A - B) \cos(A + B);$$

$$(ii) \quad \cos A + \cos(120^\circ - A) + \cos(120^\circ + A) = 0.$$

5. In any triangle the tangent of half the difference of two angles is to the tangent of half their sum as the difference of the sides respectively opposite to the angles is to the sum of these sides. Prove this.

Find the angles  $B$  and  $C$  in the triangle  $ABC$  from these data:

$$A = 60^\circ, b = 17, c = 7, \log 2 = .30103, \log 3 = .47712,$$

$$L \tan 35^\circ 49' = 9.85833, L \tan 35^\circ 49' 10'' = 9.85838.$$

6. A lighthouse facing N. sends out a fan-shaped beam extending from N.E. to N.W. A steamer sailing due W. first sees the light when 5 miles away from the lighthouse and continues to see it for  $30\sqrt{2}$  minutes. What is the speed of the steamer?

7. Find the area of the entire surface of a right cone whose height is 4 feet and diameter of base also 4 feet. Give the answer in square feet correct to two places of decimals.

#### SENIOR STUDENTS, 1880.

1. DEFINE the unit of circular measure.

A line turning about a point makes three revolutions and a sixth, what are the measures of the angle it has turned through in circular measure and in degrees?

2. Define the sine and cosine of an angle.

Prove that  $\sin^2 A + \cos^2 A = 1$ , and that  $\sin(90^\circ - A) = \cos A$ , and hence deduce the value of  $\sin 45^\circ$ .

3. Prove that

$$(1) \quad \cos(A + B) = \cos A \cos B - \sin A \sin B,$$

$$(2) \quad 4 \cos(A + B + 45^\circ) \cos(A + B - 45^\circ) \cos(A - B) \\ = \cos(A + 3B) + \cos(3A + B).$$

4. Solve the equation

$$2 \sin^2 \theta + \sin 2\theta + \cos 2\theta = 0.$$

5. Express the sine of half an angle in terms of the sine of the angle, and thence find the value of  $\sin 67^\circ. 30'$ .

6. Investigate formulæ for solving a triangle when two sides and the included angle are given.

**Ex.** Having given  $A = 60^\circ$ ,  $b = 14$  ft.,  $c = 11$  ft.,

$$\log 4.32 = .6354837, \quad L \tan 11^\circ. 44' = 9.3174299,$$

$$L \tan 11^\circ. 45' = 9.3180640,$$

solve the triangle.

7. Find the radius of the circle circumscribing a given triangle.

From the angular points of a triangle  $ABC$  perpendiculars are drawn to the opposite sides meeting them in  $D, E, F$ . Find the sides and angles of the triangle  $DEF$ , and thence the radius of the circle circumscribing it.

# SENIOR STUDENTS, 1881.

1. ESTABLISH a method for finding the number of degrees in an angle whose circular measure is given.

If the sum of the numbers expressing the measures of an angle in degrees and in circular measure be 90, determine the angle.

2. Prove that  $\sec^2 A = 1 + \tan^2 A$ , and that

$$\sin (90^\circ + A) = \cos A.$$

If  $\tan A = \frac{a}{b}$ , find the value of  $\cos A$  and of  $\tan (45^\circ + A)$ .

3. Prove that

$$(1) \quad \cos A + \cos B = 2 \cos \frac{A+B}{2} \cos \frac{A-B}{2},$$

$$(2) \quad \cos (A - 2B) - \cos (A - B) + \cos A - \cos (A + B) + \cos (A + 2B) \\ = \cos A \frac{\cos \frac{5}{2}B}{\cos \frac{1}{2}B} = 4 \cos A (\cos B - \cos 36^\circ) (\cos B + \cos 72^\circ).$$

4. Solve the equation

$$\cos 2\theta + 2 \sin^2 2\theta = 1.$$

5. Shew how to find  $\cos A$  in terms of  $\sin 2A$ .

Hence find the values of  $\cos 15^\circ$ ,  $\cos 75^\circ$ ,  $\cos 195^\circ$ ,  $\cos 255^\circ$ .

6. Prove that in any triangle

$$\tan \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{s(s-a)}}.$$

The sides of a triangle are 4, 5, 6 feet respectively, determine the least angle, having given

$$\log 7 = .845098$$

$$L \tan 20^\circ.42' = 9.577341$$

$$L \tan 20^\circ.48' = 9.577723.$$

#### SENIOR STUDENTS, 1882.

1. DEFINE the unit of circular measure, and explain what is meant by the circular measure of an angle. Assuming that  $\pi = 3.1416$ , find the number of degrees in this unit.

2. Define the trigonometrical ratios of an angle. Compare them with the corresponding ratios of its supplement.

If  $\cos A = \frac{m^2 - n^2}{m^2 + n^2}$ , find  $\sin A$  and  $\tan A$ .

3. Prove that

$$(i) \quad \cos(A - B) = \cos A \cos B + \sin A \sin B.$$

$$(ii) \quad (\sin 2A - \sin 2B) \tan(A + B) = 2(\sin^2 A - \sin^2 B).$$

4. Find an expression for all the angles which have a given tangent.

Find all the values of  $\theta$  which satisfy the equation:

$$(1 - \tan \theta)(1 + \sin 2\theta) = 1 + \tan \theta.$$

5. In any triangle the sides are proportional to the sines of the opposite angles.

If points  $D, E, F$  be taken in the sides of a triangle  $ABC$  so that

$$\frac{BD}{BC} = \frac{CE}{CA} = \frac{AF}{AB} = \frac{p}{q},$$

prove that

$$\frac{\cot DAC + \cot EBA + \cot FCB}{\cot A + \cot B + \cot C} = \frac{q+p}{q-p}.$$

6. Prove that in any triangle

$$\tan \frac{1}{2}(A - B) = \frac{a-b}{a+b} \cot \frac{C}{2}.$$

If  $a = 27$ ,  $b = 23$ ,  $C = 40^\circ.30'$ ; find  $A$  and  $B$  having given

$$L \cot 22^\circ.15' = 10.3881591, \quad L \tan 11^\circ.3' = 9.2906713,$$

$$\log 2 = .3010300, \quad L \tan 11^\circ.4' = 9.2913424.$$

ENTRANCE EXAMINATION, SANDHURST. 1881.

1. WHAT is meant by the unit of circular measure?

Prove the formula  $\theta = \frac{\text{arc}}{\text{radius}}$ .

Find the length of that part of a circular railway-curve which subtends an angle of  $22\frac{1}{2}^\circ$  to a radius of a mile. ( $\pi = 3.1416$ .)

2. Prove from a figure that

$$\cos(A - B) = \cos A \cos B + \sin A \sin B,$$

when  $A$  lies between  $315^\circ$  and  $360^\circ$ , and  $A - B$  between  $180^\circ$  and  $225^\circ$ .

3. Prove that

$$2 \sin \frac{A}{2} = \pm \sqrt{1 + \sin A} \pm \sqrt{1 - \sin A}.$$

Shew, *a priori*, the reason of the four different values of  $\sin \frac{A}{2}$  found from  $\sin A$ .

4. Prove that

$$(1) \sec^4 \theta + \tan^4 \theta = 1 + 2 \sec^2 \theta \tan^2 \theta.$$

$$(2) \sin^{-1} \left( \frac{x - a + b}{2b} \right)^{\frac{1}{2}} = \frac{1}{2} \cos^{-1} \frac{a - x}{b}.$$

$$(3) \tan 7^\circ.30' = (\cot 30^\circ - \operatorname{cosec} 45^\circ) (\sec 45^\circ - 1).$$

5. Shew that in any triangle  $ABC$

$$(b + c) \cos A + (c + a) \cos B + (a + b) \cos C = a + b + c,$$

and if  $r$ ,  $R$ ,  $r_a$ ,  $r_b$ ,  $r_c$  are the radii of the circles inscribed in, circumscribed about, and escribed to the triangle  $ABC$ ,

$$\frac{1}{r_a - r} + \frac{1}{r_b + r_c} = \frac{4R}{a^2}.$$

6. The three sides  $BC$ ,  $CA$ ,  $AB$  of a triangle are as 4 : 5 : 6. Find the angle  $B$ .

$$\log 2 = .3010300,$$

$$L \cos 27^\circ.53' = 9.9464040, \quad L \cos 27^\circ.54' = 9.9463371.$$

7. The angular elevation of a steeple at a place due south of it is  $45^\circ$ , and at another place due west of the former station the elevation is  $15^\circ$ : shew that the height of the steeple is  $\frac{a}{2} (3^{\frac{1}{2}} - 3^{-\frac{1}{2}})$ ,  $a$  being the distance between the places.

## ENTRANCE EXAMINATION, SANDHURST. 1882.

1. DISTINGUISH between the circular measure of an angle, and its measure in degrees; and prove that to turn circular measure into seconds we must multiply by 206265, and to turn seconds into circular measure we must multiply by .000004848.

2. Compare the Trigonometrical Ratios of any angle and its supplement: and determine the Trigonometrical Ratios of  $495^\circ$ .

Find all the angles between  $0^\circ$  and  $500^\circ$  which satisfy the equation

$$\sin^2 \theta = \frac{3}{4}.$$

3. Find geometrically an expression for the cosine of the difference of two angles in terms of the trigonometrical ratios of those angles.

Find all the values of  $x$ , which satisfy the equation

$$\cos x \cos 3x = \cos 2x \cos 6x.$$

Solve the equation

$$\tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{4} + \tan^{-1} \frac{1}{5} + \tan^{-1} x = \frac{\pi}{4}.$$

4. If  $r, r_1, r_2, r_3$  be the radii of the inscribed and escribed circles of a triangle, and  $x, y, z$  the perpendiculars from the angles on the opposite sides, prove

$$(1) \quad xyz = \frac{(a+b+c)^3}{abc} r^3.$$

$$(2) \quad \sin \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{bc}} = \frac{r}{\sqrt{(r_2-r)(r_3-r)}}.$$

5. In a triangle, prove that

$$\tan \left( \frac{A}{2} + B \right) = \frac{c+b}{c-b} \tan \frac{A}{2};$$

and if  $3c=7b$ , and  $A=6^\circ. 37'. 24''$ , find the other angles.

$$\text{Given } \log 2 = .30103, \quad L \tan 8^\circ. 18'. 42'' = 8.7624069,$$

$$L \tan 8^\circ. 13'. 50'' = 9.1603083, \text{ diff. for } 10'' = 1486.$$

6. In a plane triangle, prove that

$$1. \quad \tan A \tan B \tan C = \tan A + \tan B + \tan C,$$

$$2. \quad a \sin A + b \sin B + c \sin C = 2 (a \cos A + b \cos B + c \cos C),$$

where  $a, b, c$  are the sides, and  $\alpha, \beta, \gamma$  the perpendiculars let fall on them from the opposite angles respectively.

7. **P**rove that the area of a triangle

$$= \frac{a^2}{4} \sin 2B + \frac{b^2}{4} \sin 2A;$$

and if  $R, r$  are the radii of the circumscribing and inscribed circles

$$Rr = \frac{abc}{4(a+b+c)}.$$

8. Given  $\log 1\frac{1}{2} = .0791812$ , and  $\log 2\frac{2}{3} = .3802112$ , find the value of

$$\sqrt[5]{(3.6)^3} \times \sqrt[4]{\frac{1}{25}} \div \sqrt[3]{8\frac{2}{3}},$$

the mantissæ for 46929 and 46930 being 6714413 and 6714506.

9. A measured line is drawn from a point on a horizontal plane in a direction at right angles to the line joining that point to the base of a tower standing on the plane. The angles of elevation of the tower from the two ends of the measured line are  $30^\circ$  and  $18^\circ$ . Find the height of the tower in terms of the length of the measured line.

#### ENTRANCE EXAMINATION, WOOLWICH. June, 1882.

1. **P**ROVE that the angle subtended at the centre of a circle by an arc **equal** in length to its radius is an invariable angle.

One angle of a triangle is  $45^\circ$ , and the circular measure of another is  $\frac{\pi}{3}$ . **F**ind the third, both in degrees and in circular measure.

2. Define the secant of an angle, and shew how your definition applies to angles between  $180^\circ$  and  $270^\circ$ .

If  $\sec A = -2$ , what two values between  $0^\circ$  and  $360^\circ$  may  $A$  have?

3. Obtain a formula embracing all the angles which have a given **tangent**.

Determine all the values of  $\theta$  which satisfy the equation :

$$\sqrt{3} \tan^2 \theta + 1 = (1 + \sqrt{3}) \tan \theta.$$

4. Find an expression for  $\tan 3A$  in terms of  $\tan A$ . Shew also that

$$\tan 3A \tan 2A \tan A = \tan 3A - \tan 2A - \tan A.$$

5. Prove that

$$\sin 18^\circ = \frac{\sqrt{5}-1}{4};$$

and that

$$\sin^2 30^\circ = \sin 18^\circ \sin 54^\circ.$$

Shew that in any circle the chord of an arc of  $108^\circ$  is equal to the sum of the chords of arcs of  $36^\circ$  and  $60^\circ$ .

6. Demonstrate the identities

$$(1) \frac{(\operatorname{cosec} A + \sec A)^2}{\operatorname{cosec}^2 A + \sec^2 A} = 1 + \sin 2A,$$

$$(2) \sin 3A = 4 \sin A \sin (60 + A) \sin (60 - A),$$

$$(3) 4 (\cot^{-1} 3 + \operatorname{cosec}^{-1} \sqrt{5}) = \pi.$$

7. What are the advantages gained by the use of logarithms calculated to the base 10?

If  $\log_{10} 2 = .30103$ , find the logarithms of 5,  $\frac{1}{125}$ ,  $4\sqrt{.005}$  to the base 10.

8. Prove that in any triangle

$$(1) 2bc \cos A = b^2 + c^2 - a^2.$$

$$(2) \frac{1 + \cos(A - B) \cos C}{1 + \cos(A - C) \cos B} = \frac{a^2 + b^2}{a^2 + c^2}.$$

9. If  $r_1$  be the radius of a circle touching the side  $a$  of a triangle, and the other two sides produced, shew that

$$r_1 \cos \frac{A}{2} = a \cos \frac{B}{2} \cos \frac{C}{2}.$$

If  $a$  be the side of a regular polygon of  $n$  sides, and  $R$ ,  $r$  the radii respectively of its circumscribed and inscribed circles, prove that

$$R + r = \frac{a}{2} \cot \frac{\pi}{2n}.$$

10. Two sides of a triangle, which are respectively 250 and 200 yards long, contain an angle of  $54^\circ. 36'. 24''$ .

Find the two other angles, having given

$$L \cot 27^\circ. 18' = 10.2872338, \text{ diff. for } 1' = 8100,$$

$$L \tan 12^\circ. 8'. 50'' = 9.3329292 : \log 3 = .4771213.$$

11. The eye of a soldier in a straight trench of uniform depth is 2 feet above a level plain on which he sees two men standing in the same straight line as the trench; the parts of their bodies above the level of his eye subtending at it the angles  $\tan^{-1} .00416$ , and  $\tan^{-1} .004$ . On walking 200 feet towards them in the trench, he notices that the height of one exactly hides that of the other; and on approaching 596 feet 8 inches closer still he finds that the portion of the height of the nearer above the level of his eye subtends at it  $45^\circ$ . Find the heights of the men.

#### PREVIOUS EXAMINATION. June, 1880.

1. DISTINGUISH between Euclid's definition of an angle and the trigonometrical definition.

What angle does the minute-hand of a clock describe between half-past four and a quarter past six?

2. Define the cosine of an angle, and trace the change in the value of the cosine as the angle increases from  $180^\circ$  to  $360^\circ$ .

3. Express the sine and the cosine of an angle in terms of the tangent.

The angle  $A$  is greater than  $180^\circ$  but less than  $270^\circ$ , and  $\tan A = \frac{1}{2}$ : find  $\sin A$ .

4. Find  $\cos 30^\circ$  and  $\cot 135^\circ$ .

5. Prove geometrically  $\cos(A - B) = \cos A \cos B + \sin A \sin B$ ,  $A$  and  $B$  being both positive angles less than  $90^\circ$ .

Shew that

$$(1) \quad \tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B},$$

$$(2) \quad \sin 2A = \frac{2 \cot A}{1 + \cot^2 A}.$$

6. Shew that if  $A + B + C = 90^\circ$ ,

$$\sin 2A + \sin 2B + \sin 2C = 4 \cos A \cos B \cos C.$$

7. Find an expression for all the values of  $\theta$  for which  $\sin \theta + \sin 2\theta = 0$ .

8. Express the cosine of an angle of a triangle in terms of the sides.

If in a triangle  $b \cos A = a \cos B$ , shew that the triangle is isosceles.

9. If two sides of a triangle be given, and the angle opposite to one of them; shew how to find the other side and the other angles.

Ex. The sides are 1 foot and  $\sqrt{2}$  feet respectively, and the angle opposite to the shorter side is  $30^\circ$ .

# PREVIOUS EXAMINATION. June, 1882.

1. DEFINE the unit of circular measure. Calling this angle the radian, find how many radians there are in a right angle.

If the circumference of a circle be divided into five parts in arithmetical progression, the greatest part being six times the least, express in radians the angle each subtends at the centre.

2. Define the sine of an angle, wording your definition so as to include angles of any magnitude.

Prove that

$$\sin(90^\circ + A) = \cos A,$$

and

$$\cos(90^\circ + A) = -\sin A,$$

and by means of these deduce the formulæ

$$\sin(180^\circ + A) = -\sin A, \quad \cos(180^\circ + A) = -\cos A.$$



3. Prove the formulæ

$$(1) \cot^2 A = \operatorname{cosec}^2 A - 1,$$

$$(2) \cot^4 A + \cot^2 A = \operatorname{cosec}^4 A - \operatorname{cosec}^2 A.$$

Verify (2) when  $A = 30^\circ$ .

4. Shew that all angles satisfying the equation  $\cos \theta = \cos \alpha$  are included in the formula  $\theta = 2n\pi \pm \alpha$ .

Solve completely the equation  $2 \cos^3 \theta + \sin^3 \theta - 1 = 0$ .

5. Prove that  $\sin(A+B) = \sin A \cos B + \cos A \sin B$ , and deduce the expression for  $\cos(A+B)$ .

Shew that  $\sin A \cos(B+C) - \sin B \cos(A+C) = \sin(A-B) \cos C$ .

6. Prove that

$$\cos \frac{A}{2} + \sin \frac{A}{2} = \pm \sqrt{1 + \sin A},$$

$$\cos \frac{A}{2} - \sin \frac{A}{2} = \pm \sqrt{1 - \sin A},$$

and determine the signs of the ambiguities when  $A$  lies between  $450^\circ$  and  $630^\circ$ .

7. Establish the identities

$$(1) 1 + \cos A + \sin A = \sqrt{2(1 + \cos A)(1 + \sin A)},$$

$$(2) \operatorname{cosec} 2A = \frac{\operatorname{cosec}^3 A}{2\sqrt{\operatorname{cosec}^2 A - 1}},$$

$$(3) \sin \frac{2\pi}{7} + \sin \frac{4\pi}{7} - \sin \frac{6\pi}{7} = 4 \sin \frac{\pi}{7} \sin \frac{3\pi}{7} \sin \frac{5\pi}{7}.$$

8. Shew that in any triangle (with the usual notation)

$$a^2 = b^2 + c^2 - 2bc \cos A.$$

If  $a=7$ ,  $b=8$ ,  $c=9$ , shew that the length of the line joining the angular point  $B$  to the middle point of the opposite side is 7.

9. Solve completely the triangle whose sides are given in the preceding question, given  $\log 2$ ,  $\log 6$ ,

$$L \tan 24^\circ.5' = 9.6502809, \text{ tabular difference for } 60'' = .0003390,$$

$$L \tan 36^\circ.41' = 9.8721123, \text{ tabular difference for } 60'' = .0002637.$$

#### MATHEMATICAL TRIPOS. January 3, 1876.

1. DEFINE the cotangent and the cosecant of an angle, and trace the changes in the values of these functions as the angle increases from two to four right angles.

What values of  $\theta$  satisfy the equation

$$\operatorname{cosec} 4\alpha - \operatorname{cosec} 4\theta = \cot 4\alpha - \cot 4\theta?$$

2. Prove the formulæ

$$(1) \cos(A+B) = \cos A \cos B - \sin A \sin B,$$

$$(2) \cos B - \cos A = 2 \sin \frac{A+B}{2} \sin \frac{A-B}{2}.$$

Prove that if  $A+B+C=180^\circ$ , and if

$\sin(2n+1)A \sin(B-C) + \sin(2n+1)B \sin(C-A) + \sin(2n+1)C \sin(A-B) = 0$ ,  
 $n$  being an integer, then

$$\begin{aligned} \sin(n-1)A \sin(n+1)(B-C) + \sin(n-1)B \sin(n+1)(C-A) \\ + \sin(n-1)C \sin(n+1)(A-B) = 0. \end{aligned}$$

3. Shew how to express the tangent of half an angle of a triangle in terms of the sides.

The sides are observed to be  $a=5$ ,  $b=4$ ,  $c=6$ , but it is known that there is a small error in the measurement of  $c$ : examine which angle can be determined with the greatest accuracy.

4. Shew how to find the height of a mountain by means of observations made at two given places in a horizontal plane.

On a plane which is inclined to the horizon at an angle of  $45^\circ$  is described a circle of known radius, and a post is placed at the highest point of the circle perpendicular to the plane; at one end of the horizontal diameter of the circle the tangent of the angular elevation of the post is  $\sqrt{2}$ . Find the length of the post.

5. If  $O, O_1, O_2, O_3$  be the centres of the inscribed and escribed circles of a triangle, and  $r, r_1, r_2, r_3$  the radii of those circles, and  $R$  the radius of the circumscribing circle; shew that

$$4R = \frac{r}{\sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}} = \frac{r_1}{\sin \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2}},$$

and that the areas of the triangles  $O_1O_2O_3, O_2O_3O, O_3OO_1, OO_1O_2$  are to one another inversely as  $r, r_1, r_2, r_3$ .

6. From a point  $P$  perpendiculars  $PL, PM, PN$  are drawn to the sides of a triangle  $ABC$ : shew that twice the area of the triangle  $LMN$  is equal to

$$(R^2 - d^2) \sin A \sin B \sin C,$$

where  $R$  is the radius of the circle circumscribing the triangle  $ABC$ , and  $d$  the distance of its centre from  $P$ .

## MATHEMATICAL TRIPOS. 1869.

1. PROVE the formulæ:

(1)  $\tan A = \tan (n \cdot 180^\circ + A)$ , when  $n$  is a positive or negative integer.

(2)  $\sin (A - B) = \sin A \cos B - \sin B \cos A$ , where  $180^\circ$ ,  $A$ ,  $90^\circ$ ,  $B$ ,  $0^\circ$  are in descending order of magnitude.

(3)  $2 \sin \frac{A}{2} = \sqrt{1 + \sin A} + \sqrt{1 - \sin A}$ , where  $A$  is greater than  $90^\circ$  and less than  $270^\circ$ .

2. If  $\beta$ ,  $\gamma$  be two different values of  $\theta$  which satisfy the equation

$$\frac{1}{a} \cos \theta + \frac{1}{b} \sin \theta = \frac{1}{c},$$

then will  $a \cos \frac{\beta + \gamma}{2} = b \sin \frac{\beta + \gamma}{2} = c \cos \frac{\beta - \gamma}{2}$ .

Having given

$$a^2 \cos \alpha \cos \beta + a (\sin \alpha + \sin \beta) + 1 = 0,$$

$$a^2 \cos \alpha \cos \gamma + a (\sin \alpha + \sin \gamma) + 1 = 0,$$

prove that  $a^2 \cos \beta \cos \gamma + a (\sin \beta + \sin \gamma) + 1 = 0$ ,

and that  $\cos \alpha + \cos \beta + \cos \gamma = \cos (\alpha + \beta + \gamma)$ ,

$\beta$ ,  $\gamma$  being unequal and less than  $\pi$ .

3. Investigate the expression for one side of a triangle in terms of the other sides and the angle included by them, and put it into a form proper for logarithmic computation.

Equilateral triangles  $DBC$ ,  $D'BC$  are described on the side  $BC$  of a triangle  $ABC$ ; prove that  $AD^2 + AD'^2 = a^2 + b^2 + c^2$ , and express  $\cos DAD'$  in terms of the sides  $a$ ,  $b$ ,  $c$ . Hence shew that if equilateral triangles be in like manner constructed on the other two sides, and the angles  $DAD'$ ,  $EBE'$ ,  $FCF'$  be denoted by  $\alpha$ ,  $\beta$ ,  $\gamma$ ,

$$\cos \alpha + \cos \beta + \cos \gamma = 0 = \cos 2\alpha + \cos 2\beta + \cos 2\gamma.$$

4. If  $A$ ,  $B$ ,  $C$  be angles of a triangle, and  $x$ ,  $y$ ,  $z$  any real quantities satisfying the equation

$$\frac{y \sin C - z \sin B}{x - y \cos C - z \cos B} = \frac{z \sin A - x \sin C}{y - z \cos A - x \cos C},$$

then will

$$\frac{x}{\sin A} = \frac{y}{\sin B} = \frac{z}{\sin C}.$$

5. Given one side of a triangle and its angles, determine its area and the radii of the four circles which can be described touching the sides.

If  $a$ ,  $b$ ,  $c$ ,  $d$  be the sides of a convex quadrilateral in which a circle can be inscribed, its area will be  $\sqrt{abcd} \sin \omega$ ,  $2\omega$  being the sum of two opposite angles.

6. A gun is fired from a fort  $A$  and the intervals between seeing the flash and hearing the report at two stations  $B$ ,  $C$  are  $t$ ,  $t'$  respectively;  $D$  is

a point in the same straight line with  $BC$  at a known distance  $a$  from  $A$  : prove that if  $BD=b$ , and  $CD=c$ , the velocity of sound is

$$\left\{ \frac{(b-c)(a^2-bc)}{bt^2-ct^2} \right\}^{\frac{1}{2}}.$$

Examine the case where  $a^2=bc$ .

### MATHEMATICAL TRIPOS. 1870.

1. DEFINE a logarithm. Shew that in the common system of logarithms the integral part of any logarithm need not be tabulated. If a table of logarithms be calculated to any base, prove that they may be transformed to any other base by multiplication by a constant factor.

If  $a, b, c$  be in Geometrical Progression, and  $\log_a a, \log_a b, \log_a c$  in Arithmetical Progression, the common difference of this progression is  $\frac{3}{2}$ .

2. Assuming that the formulæ

$$(1) \quad \sin(A+B) = \sin A \cos B + \cos A \sin B,$$

$$(2) \quad \cos(A+B) = \cos A \cos B - \sin A \sin B,$$

are true for angles less than a right angle, shew that they are also true for positive angles of any magnitude.

Prove the formulæ :

$$(1) \quad \cos 2A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A,$$

$$(2) \quad \cos^2 A + \sin^2 A \cos 2B = \cos^2 B + \sin^2 B \cos 2A,$$

$$(3) \quad \sin^2 A - \cos^2 A \cos 2B = \sin^2 B - \cos^2 B \cos 2A.$$

3. If  $A+B+C$  be a multiple of  $180^\circ$ , then will

$$\cot B \cot C + \cot C \cot A + \cot A \cot B = 1.$$

If  $A+B+C = (2m+1)\pi$  or  $2m\pi + \frac{\pi}{2}$ , then will

$$\begin{aligned} & (\sin A + \cos A)(\sin B + \cos B)(\sin C + \cos C) \\ &= 2 \sin A \sin B \sin C + 2 \cos A \cos B \cos C + 1, \end{aligned}$$

but if  $A+B+C = 2m\pi$  or  $2m\pi - \frac{\pi}{2}$ , then will

$$\begin{aligned} & (\sin A + \cos A)(\sin B + \cos B)(\sin C + \cos C) \\ &= 2 \sin A \sin B \sin C + 2 \cos A \cos B \cos C - 1. \end{aligned}$$

4. Shew how to solve a triangle in which are given the sides  $a, b$  and the angle  $A$  : and determine in what cases there are two solutions, one, or none.

If there be two triangles having the given parts, and if  $c_1, c_2$  be their third sides, the distance between the centres of their circumscribing circles

$$= \frac{c_1 - c_2}{2 \sin A}.$$

5. If  $a, b, c$  be the sides of a triangle, and  $2s = a+b+c$ , then the area of the triangle  $= \sqrt{s(s-a)(s-b)(s-c)}$ .

If  $S_1, S_2, S_3, S_4$  be the areas of the four triangles whose sides are  $b, c, d;$   
 $c, d, a;$   $d, a, b;$   $a, b, c$  respectively, prove that

$$\frac{S_1^2 - S_2^2}{a^2 - b^2} + \frac{S_2^2 - S_4^2}{c^2 - d^2} = \frac{S_2^2 - S_3^2}{b^2 - c^2} + \frac{S_4^2 - S_1^2}{d^2 - a^2}.$$

6. Find the radius of the circle which touches one side of a triangle and the other two produced.

If  $r, r'$  be the radii of two circles, and if  $P, P'$  be the perimeters,  $A, A'$  the areas of the triangles whose sides touch them, prove that

$$(1) \quad PP'(r - r') - rP^2 = 4r^2r',$$

$$(2) \quad A^2(r - r')^2 - A'^2(r + r')^2 = 4r^2r'^2,$$

$r, P, A$ , being greater respectively than  $r', P', A'$ .

#### MATHEMATICAL TRIPOS. 1871.

1. DEFINE a logarithm, and prove, if  $a, b, N$  be any three numbers,

$$\log_a b \log_b a = 1, \text{ and } \log_a N = \frac{\log_b N}{\log_b a},$$

prove also that

$$\frac{\log_a \log_a N}{\sqrt{\log_a b}} - \frac{\log_b \log_b N}{\sqrt{\log_b a}} = \frac{\log_a \log_a b}{\sqrt{\log_a b}} = -\frac{\log_b \log_b a}{\sqrt{\log_b a}}.$$

2. Find a general expression representing all the angles which have a given sine.

If  $\sin(\theta + \alpha) = \sin(\phi + \alpha) = \sin \beta$ ,  
 and  $a \sin(\theta + \phi) + b \sin(\theta - \phi) = c$ ,

prove that, either

$$a \sin(2\alpha \pm 2\beta) = -c, \text{ or } a \sin 2\alpha \pm b \sin 2\beta = c.$$

3. Prove the formula

$$\sin(A - B) = \sin A \cos B - \cos A \sin B,$$

where  $A$  is greater than  $B$ , and each of them less than  $90^\circ$ . Also assuming this formula, prove

$$\sin(A + B) = \sin A \cos B + \cos A \sin B.$$

If  $\sqrt{2} \cos A = \cos B + \cos^3 B$ ,  
 and  $\sqrt{2} \sin A = \sin B - \sin^3 B$ ,

prove that  $\pm \sin(B - A) = \cos 2B = \frac{1}{3}$ .

If  $4 \cos(x - y) \cos(y - z) \cos(z - x) = 1$ ,  
 prove that

$$1 + 12 \cos 2(x - y) \cos 2(y - z) \cos 2(z - x) = 4 \cos 3(x - y) \cos 3(y - z) \cos 3(z - x).$$

4. Investigate a formula connecting any angle of a triangle with the sides of the triangle.

$O$  is any point in the interior of an equilateral triangle  $ABC$ , prove that

$$\cos(\angle BOC - 60^\circ) = \frac{BO^2 + CO^2 - AO^2}{2BO \cdot CO}.$$

5. Find the radius of the circle which is described about a given triangle.

If  $G$  be the centre of gravity of a triangle, and  $R_1, R_2, R_3$  be the radii of the circles circumscribing  $BGC, CGA, AGB$  respectively: prove that

$$\frac{a^2(b^2 - c^2)}{R_1^3} + \frac{b^2(c^2 - a^2)}{R_2^3} + \frac{c^2(a^2 - b^2)}{R_3^3} = 0.$$

6. Find the lengths of the diagonals of a quadrilateral inscribed in a circle in terms of its sides.

If  $ABCD$  be the quadrilateral,  $\Delta$  its area, and if  $CC'$  be a chord parallel to  $BD$ , find the length of  $CC'$  in terms of the sides, and prove that the radius of the circle is

$$\frac{AC \cdot AC \cdot BD}{4\Delta}.$$

*Note.* The centre of gravity of a triangle is the intersection of the lines drawn from the angles to bisect the opposite sides.

The following tables comprise all the natural sines, &c. and logarithms required for the solution of the examples in this book; but the student is recommended where practicable to use complete tables, as he will thus become more familiar with their use. He will however only be able to arrive at an approximate solution, without the use of proportional parts. In many of the earlier examples the natural sines, &c. have only been employed to four decimal places. The student must remember that when he wants to find a cosine, he must look for the sine of the complement, and similarly for the cotangents and cosecants.

TABLE I.  
NATURAL SINES, &c.

Sines.		Sines.	
5°. 58'	·1039499	67°. 23'	·9230984
7°. 46'	·1351392	68°. 13'	·9285933
11°. 32'	·1999380	73°. 32'	·9589848
12°	·2079117	Tangents.	
16°. 26'	·2828995		
19°. 28'	·3332584	10°. 40'	·1883495
21°. 47'	·3710977	26°	·4877326
25°	·4226183	26°. 34'	·5000352
32°	·5299193	33°. 41'	·6664969
32°. 53'	·5429302	37°	·7535541
34°. 24'	·5649670	50°. 46'	1·2246658
36°. 53'	·6001876	52°. 50'	1·3190441
38°. 13'	·6186370	61°	1·8040478
39°	·6293204	67°. 23'	2·4003774
40°. 39'	·6514366	Secants.	
43°. 26'	·6875101		
45°. 35'	·7142691	26°	1·1126019
48°. 24'	·7477981	52°. 50'	1·6552575
51°	·7771460	66°. 48'	2·5384453
61°. 3'	·8750425	68°. 12'	2·6927480
65°	·9063078	73°. 45'	3·5736108

# PLANE TRIGONOMETRY.

## TABLE II.

### LOGARITHMS OF NUMBERS.

Numbers.	Logarithms.	Numbers.	Logarithms.
1.068185	·0286466	2.41	·3820170
1.1	·0413927	3	·4771213
1.143716	·0583182	3.09017	·4899824
1.22239	·0872111	3.652281	·5625642
1.3	·1139434	3.727593	·5714286
1.39009	·1430448	4.013116	·6034817
1.4458	·1601082	4.03826	·6061942
1.590643	·2015726	4.23665	·6270227
1.62335	·2104122	4.251787	·6285714
1.69408	·2289333	4.45813	·6491525
2	·3010300	7	·8450980
2.07	·3159703	7.39148	·8687314
2.3	·3617278	7.90672	·8979965

## TABLE III.

### TABULAR LOGARITHMS OF THE TRIGONOMETRICAL RATIOS.

Sines.	Logarithms.	Tangents.	
6°. 22'. 45"	9.0457574	3°. 43'. 39". 03	8.8139084
17°. 11'. 12". 11	9.4705539	10°. 53'. 36"	9.2843181
18°	9.4899824	11°. 16'. 10"	9.2994355
23°. 42'. 43"	9.6043801	11°. 44'. 29". 5	9.3177416
29°. 5'. 21". 52	9.6867903	34°. 6'. 24"	9.8307301
41°. 8'. 54". 6	9.8182344	57°. 22'. 26". 8	10.1937091
41°. 42'. 43"	9.8230636	57°. 54'	10.2025255
48°. 51'. 5". 94	9.8767989	63°. 47'. 35". 59	10.3078513
53°. 27'. 20"	9.9049296	71°. 33'. 54"	10.4771213
62°. 8'. 51"	9.9463950		
64°. 25'. 49"	9.9552358		
72°	9.9782063		

*By the same Author.*

ARITHMETIC FOR SCHOOLS,

With numerous Examples, and Examination Papers. Price 3s.

*The Examples can also be had separately.*

ELEMENTARY EXAMPLES. Price 8d.

HIGHER EXAMPLES, and PAPERS. Price 1s. 6d.

RIVINGTONS AND CO.





*September, 1883.*

A  
CATALOGUE  
OF  
**EDUCATIONAL BOOKS**

PUBLISHED BY

**MACMILLAN AND CO.,**  
**BEDFORD STREET, STRAND, LONDON.**

# INDEX.

	PAGE
<b>CLASSICAL—</b>	
ELEMENTARY CLASSICS . . . . .	3
CLASSICAL SERIES . . . . .	6
CLASSICAL LIBRARY (1) Texts, (2) Translations . . . . .	11
GRAMMAR, COMPOSITION, AND PHILOLOGY . . . . .	15
ANTIQUITIES, ANCIENT HISTORY, AND PHILOSOPHY . . . . .	20
<b>MATHEMATICS—</b>	
ARITHMETIC . . . . .	22
ALGEBRA . . . . .	24
EUCLID AND ELEMENTARY GEOMETRY . . . . .	24
MENSURATION . . . . .	25
HIGHER MATHEMATICS . . . . .	26
<b>SCIENCE—</b>	
NATURAL PHILOSOPHY . . . . .	33
ASTRONOMY . . . . .	37
CHEMISTRY . . . . .	38
BIOLOGY . . . . .	39
MEDICINE . . . . .	43
ANTHROPOLOGY . . . . .	44
PHYSICAL GEOGRAPHY AND GEOLOGY . . . . .	44
AGRICULTURE . . . . .	45
POLITICAL ECONOMY . . . . .	45
MENTAL AND MORAL PHILOSOPHY . . . . .	46
<b>HISTORY AND GEOGRAPHY . . . . .</b>	47
<b>MODERN LANGUAGES AND LITERATURE—</b>	
ENGLISH GRAMMAR AND LITERATURE . . . . .	51
GLOBE READERS . . . . .	52
GLOBE READINGS . . . . .	52
LITERATURE PRIMERS . . . . .	53
READING BOOKS . . . . .	54
COPY BOOKS . . . . .	54
<b>FRENCH . . . . .</b>	58
PROGRESSIVE FRENCH COURSE . . . . .	59
PROGRESSIVE FRENCH READERS . . . . .	59
<b>FOREIGN SCHOOL CLASSICS—</b>	
FRENCH . . . . .	59
GERMAN . . . . .	61
<b>GERMAN—</b>	
PROGRESSIVE GERMAN COURSE . . . . .	60
<b>MODERN GREEK . . . . .</b>	61
<b>ITALIAN . . . . .</b>	62
<b>DOMESTIC ECONOMY . . . . .</b>	62
<b>ART AND KINDRED SUBJECTS . . . . .</b>	63
<b>WORKS ON TEACHING . . . . .</b>	63
<b>DIVINITY . . . . .</b>	64

*September, 1883.*

# A CATALOGUE OF EDUCATIONAL BOOKS

PUBLISHED BY

**MACMILLAN & CO.,**  
BEDFORD STREET, STRAND, LONDON.

## **CLASSICS.** **ELEMENTARY CLASSICS.**

18mo, eighteenpence each.

**THIS SERIES FALLS INTO TWO CLASSES—**

(1) **First Reading Books for Beginners**, provided not only with **Introductions and Notes**, but with **Vocabularies**.

(2) **Stepping-stones to the study of particular authors**, intended for more advanced students who are beginning to read such authors as Terence, Plato, the Attic Dramatists, and the harder parts of Cicero, Horace, Virgil, and Thucydides.

These are provided with **Introductions and Notes**, but **no Vocabulary**. The Publishers have been led to provide the more strictly **Elementary Books** with **Vocabularies** by the representations of many teachers, who hold that **beginners** do not understand the use of a Dictionary, and of **others** who, in the case of the very large class of **middle-class schools** where the cost of books is a serious consideration, advocate the **Vocabulary system** on grounds of economy. It is hoped that the two parts of the Series, fitting into one another, may together fulfil all the requirements of **Elementary and Preparatory Schools**, and the **Lower Forms of Public Schools**.

The following Elementary Books, with Introductions, Notes, and Vocabularies, are either ready or in preparation :—

**Cæsar.**—THE GALLIC WAR. BOOK I. Edited by A. S. WALPOLE, M.A. [Ready.]

THE GALLIC WAR. BOOKS II. AND III. Edited by W. G. RUTHERFORD, M.A., Head-Master of Westminster School. [Ready.]

THE GALLIC WAR. SCENES FROM BOOKS V. AND VI. Edited by C. COLBECK, M.A., Assistant-Master at Harrow = formerly Fellow of Trinity College, Cambridge. [Ready. VOCABULARY in preparation.]

**Cicero.**—DE SENECTUTE. Edited by E. S. SHUCKBURGH, M.A., late Fellow of Emmanuel College, Cambridge ; Assistant-Master at Eton College. [In preparation.]

DE AMICITIA. By the same Editor. [In preparation.]

**Eutropii Historia Romana.**—SELECTIONS. Edited by WILLIAM WELCH, Assistant-Master at Surrey County School, Cranleigh. [In the press.]

**Greek Testament.**—SELECTIONS FROM THE GOSPELS. Edited by Rev. G. F. MACLEAR, D.D., Warden of St. Augustine's College, Canterbury. [In preparation.]

**Homer.**—ILIAD. BOOK I. Edited by Rev. JOHN BOND, M.A., and A. S. WALPOLE, M.A. [In preparation.]

ILIAD. BOOK XVIII. THE ARMS OF ACHILLES. Edited by S. R. JAMES, M.A., Assistant-Master at Eton College. [Ready. VOCABULARY in preparation.]

ODYSSEY. BOOK I. Edited by Rev. JOHN BOND, M.A., and A. S. WALPOLE, M.A. [Ready.]

**Horace.**—ODES. BOOKS I.—III. Edited by T. E. PAGE, M.A., late Fellow of St. John's College, Cambridge ; Assistant-Master at the Charterhouse. Each 1s. 6d.

[Ready. VOCABULARIES in preparation.]

BOOK IV. By the same Editor. [In the press.]

**Livy.**—BOOK I. Edited by H. M. STEPHENSON, M.A., Head Master of St. Peter's School, York. [Ready.]

**THE HANNIBALIAN WAR.** Being part of the XXI. AND XXII. BOOKS OF LIVY, adapted for the use of beginners, by G. C. MACAULAY, M.A., Assistant-Master at Rugby; formerly Fellow of Trinity College, Cambridge. [Ready.]

**Ovid.**—SELECTIONS. Edited by E. S. SHUCKBURGH, M.A. [Ready. VOCABULARY in preparation.]

**Phædrus.**—SELECT FABLES. Edited by A. S. WALPOLE, M.A. [In the press.]

**Thucydides.**—THE RISE OF THE ATHENIAN EMPIRE. BOOK I. cc. LXXXIX. — CXVIII. AND CXXVIII. — CXXXVIII. Edited by F. H. COLSON, B.A., Fellow of St. John's College, Cambridge; Assistant-Master at Clifton College. [In preparation.]

**Virgil.**—ÆNEID. BOOK I. Edited by A. S. WALPOLE, M.A. [Ready.]

ÆNEID. BOOK V. Edited by Rev. A. CALVERT, M.A., Fellow of St. John's College, Cambridge. [Ready.]

SELECTIONS. Edited by E. S. SHUCKBURGH, M.A. [Ready.]

**Xenophon.**—ANABASIS. BOOK I. Edited by A. S. WALPOLE, M.A. [Ready.]

The following more advanced Books, with Introductions and Notes, but no Vocabulary, are either ready, or in preparation:—

**Cicero.**—SELECT LETTERS. Edited by Rev. G. E. JEANS, M.A., Fellow of Hertford College, Oxford, and Assistant-Master at Haileybury College. [Ready.]

**Euripides.**—HECUBA. Edited by Rev. JOHN BOND, M.A. and A. S. WALPOLE, M.A. [Ready.]

**Herodotus.**—SELECTIONS FROM BOOKS VII. AND VIII. THE EXPEDITION OF XERXES. Edited by A. H. COOKE, M.A., Fellow and Lecturer of King's College, Cambridge. [Ready.]

**Horace.**—SELECTIONS FROM THE SATIRES AND EPISTLES. Edited by Rev. W. J. V. BAKER, M.A., Fellow of St. John's College, Cambridge; Assistant-Master at Marlborough. [Ready.]

**Livy.**—THE LAST TWO KINGS OF MACEDON. SCENES FROM THE LAST DECADE OF LIVY. Selected and Edited by F. H. RAWLINS, M.A., Fellow of King's College, Cambridge; and Assistant-Master at Eton College. [In preparation.]

**Plato.**—EUTHYPHRO AND MENEXENUS. Edited by C. E. GRAVES, M.A., Classical Lecturer and late Fellow of St. John's College, Cambridge. [Ready.]

**Terence.**—SCENES FROM THE ANDRIA. Edited by F. W. CORNISH, M.A., Assistant-Master at Eton College. [In the press.]

**The Greek Elegiac Poets.**—FROM CALLINUS TO CALLIMACHUS. Selected and Edited by Rev. HERBERT KYNASTON, M.A., Principal of Cheltenham College, and formerly Fellow of St. John's College, Cambridge. [Ready.]

**Thucydides.**—BOOK IV. CHS. I.—XLI. THE CAPTURE OF SPHACTERIA. Edited by C. E. GRAVES, M.A. [Ready.]

**Virgil.**—GEORGICS. BOOK II. Edited by Rev. J. H. SKRINE, M.A., late Fellow of Merton College, Oxford; Assistant-Master at Uppingham. [Ready.]

\* \* Other Volumes to follow.

---

### CLASSICAL SERIES FOR COLLEGES AND SCHOOLS.

Fcap. 8vo.

Being select portions of Greek and Latin authors, edited with Introductions and Notes, for the use of Middle and Upper forms of Schools, or of candidates for Public Examinations at the Universities and elsewhere.

**Æschines.**—IN CTESIPHONTEM. Edited by Rev. T. GWATKIN, M.A., late Fellow of St. John's College, Cambridge. [In the press.]

**Æschylus.**—PERSÆ. Edited by A. O. PRICKARD, M.A.,  
Fellow and Tutor of New College, Oxford. With Map. 3s. 6d.

**Catullus.**—SELECT POEMS. Edited by F. P. SIMPSON, B.A.,  
late Scholar of Balliol College, Oxford. New and Revised  
Edition. 5s.

**Cicero.**—THE ACADEMICA. Edited by JAMES REID, M.L.,  
Fellow of Caius College, Cambridge. 4s. 6d.

THE CATILINE ORATIONS. From the German of KARL  
HALM. Edited, with Additions, by A. S. WILKINS, M.A., Pro-  
fessor of Latin at the Owens College, Manchester. New Edition.  
3s. 6d.

PRO LEGE MANILIA. Edited after HALM by Professor A. S.  
WILKINS, M.A. 3s. 6d.

THE SECOND PHILIPPIC ORATION. From the German  
of KARL HALM. Edited, with Corrections and Additions,  
by JOHN E. B. MAYOR, Professor of Latin in the University of  
Cambridge, and Fellow of St. John's College. New Edition,  
revised. 5s.

PRO ROSCIO AMERINO. Edited, after HALM, by E. H. DON-  
KIN, M.A., late Scholar of Lincoln College, Oxford; Assistant-  
Master at Sherborne School. 4s. 6d.

PRO P. SESTIO. Edited by Rev. H. A. HOLDEN, M.A., LL.D.,  
late Fellow of Trinity College, Cambridge; and Classical Exam-  
iner to the University of London. 5s.

**Demosthenes.**—DE CORONA. Edited by B. DRAKE, M.A.,  
late Fellow of King's College, Cambridge. New and revised  
Edition. 4s. 6d.

ADVERSUS LEPTINEM. Edited by Rev. J. R. KING, M.A.,  
Fellow and Tutor of Oriel College, Oxford. 4s. 6d.

THE FIRST PHILIPPIC. Edited, after C. REHDANTZ, by Rev.  
T. GWATKIN, M.A., late fellow of St. John's College, Cambridge.  
2s. 6d.

**Euripides.**—BACCHAE. Edited by E. S. SHUCKBURGH, M.A.,  
Assistant-Master at Eton College. [*In preparation.*]

HIPPOLYTUS. Edited by J. P. MAHAFFY, M.A., Fellow and  
Professor of Ancient History in Trinity College, Dublin, and J. B.  
BURY, Scholar of Trinity College, Dublin. 3s. 6d.



- Euripides.**—MEDEA. Edited by A. W. VERRALL, M.A., Fellow and Lecturer of Trinity College, Cambridge. 3s. 6d.
- IPHIGENIA IN TAURIS. Edited by E. B. ENGLAND, M.A., Lecturer at the Owens College, Manchester. *[In the press.]*
- Herodotus.**—BOOKS V. AND VI. Edited by Rev. A. H. COOKE, M.A., Fellow of King's College, Cambridge. *[In preparation.]*
- BOOKS VII. AND VIII. THE INVASION OF GREECE BY XERXES. Edited by THOMAS CASE, M.A., formerly Fellow of Brasenose College, Oxford. *[In preparation.]*
- Homer.**—ILIAD. BOOKS I., IX., XI., XVI.—XXIV. THE STORY OF ACHILLES. Edited by the late J. H. PRATT, M.A., and WALTER LEAF, M.A., Fellows of Trinity College, Cambridge. 6s.
- ODYSSEY. BOOKS IX.—XII. THE NARRATIVE OF ODYSSEUS. Edited by Prof. JOHN E. B. MAYOR. Part I. 3s.
- ODYSSEY. BOOKS XXI.—XXIV. THE TRIUMPH OF ODYSSEUS. Edited by S. G. HAMILTON, B.A., Fellow of Hertford College, Oxford. 3s. 6d.
- Horace.**—THE ODES AND CARMEN SECLARE. Edited by T. E. PAGE, M.A., formerly Fellow of St. John's College, Cambridge; Assistant-Master at Charterhouse. 6s.
- THE ODES. BOOK IV. and CARMEN SECLARE. Edited by T. E. PAGE, M.A. 2s.
- THE SATIRES. Edited by ARTHUR PALMER, M.A., Fellow of Trinity College, Dublin; Professor of Latin in the University of Dublin. 6s.
- THE EPISTLES AND ARS POETICA. Edited by Professor A. S. WILKINS, M.A. *[In the press.]*
- Juvenal.** Edited, for the Use of Schools, by E. G. HARDY, M.A., Head-Master of Grantham Grammar School; late Fellow of Jesus College, Oxford. *[In the press.]*
- SELECT SATIRES. Edited by Professor JOHN E. B. MAYOR. X. AND XI. 3s. 6d. XII.—XVI. 4s. 6d.
- Livy.**—BOOKS II. AND III. Edited by Rev. H. M. STEPHENSON, M.A., Head-Master of St. Peter's School, York. 5s.
- BOOKS XXI. AND XXII. Edited by the Rev. W. W. CAPES, M.A., Reader in Ancient History at Oxford. With Maps. 5s.
- BOOKS XXIII. AND XXIV. Edited by G. C. MACAULAY, M.A., Assistant-Master at Rugby. *[In preparation.]*

- Lucretius.** BOOKS I.—III. Edited by J. H. WARBURTON LEE, M.A., late Scholar of Corpus Christi College, Oxford, and Assistant-Master at Rossall. [*In preparation.*]
- Lysias.**—SELECT ORATIONS. Edited by E. S. SHUCKBURGH, M.A., Assistant-Master at Eton College. 6s.
- Martial.** — SELECT EPIGRAMS. Edited by Rev. H. M. STEPHENSON, M.A. 6s.
- Ovid.**—FASTI. Edited by G. H. HALLAM, M.A., Fellow of St. John's College, Cambridge, and Assistant-Master at Harrow. With Maps. 5s.
- HEROIDUM EPISTULÆ XIII. Edited by E. S. SHUCKBURGH, M.A. 4s. 6d.
- METAMORPHOSES. BOOKS XIII. AND XIV. Edited by C. SIMMONS, M.A. [*In the press.*]
- Plato.**—MENO. Edited by E. S. THOMPSON, M.A., Fellow of Christ's College, Cambridge. [*In preparation.*]
- APOLOGY AND CRITO. Edited by F. J. H. JENKINSON, M.A., Fellow of Trinity College, Cambridge. [*In preparation.*]
- THE REPUBLIC. BOOKS I.—V. Edited by T. H. WARREN, M.A., Fellow of Magdalen College, Oxford. [*In preparation.*]
- Plautus.**—MILES GLORIOSUS. Edited by R. V. TYRRELL, M.A., Fellow and Professor of Greek in Trinity College, Dublin. 5s.
- Pliny.**—LETTERS. BOOK III. Edited by Professor JOHN E. B. MAYOR. With Life of Pliny by G. H. RENDALL, M.A. 5s.
- Plutarch.**—LIFE OF THEMISTOKLES. Edited by Rev. H. A. HOLDEN, M.A., LL.D. 5s.
- Propertius.**—SELECT POEMS. Edited by Professor J. P. POSTGATE, M.A., Fellow of Trinity College, Cambridge. 6s.
- Sallust.**—CATILINE AND JUGURTHA. Edited by C. MERVILLE, D.D., Dean of Ely. New Edition, carefully revised and enlarged, 4s. 6d. Or separately, 2s. 6d. each.

**Sophocles.**—ANTIGONE. Edited by Rev. JOHN BOND, M.A., and A. S. WALPOLE, M.A. *[In preparation.]*

**Tacitus.**—AGRICOLA AND GERMANIA. Edited by A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A., Translators of Tacitus. New Edition, 3s. 6d. Or separately, 2s. each.

THE ANNALS. BOOK VI. By the same Editors. 2s. 6d.

THE HISTORY. BOOKS I. AND II. Edited by C. E. GRAVES, M.A. *[In preparation.]*

**Terence.**—HAUTON TIMORUMENOS. Edited by E. S. SHUCKBURGH, M.A. 3s. With Translation, 4s. 6d.

PHORMIO. Edited by Rev. JOHN BOND, M.A., and A. S. WALPOLE, B.A. 4s. 6d.

**Thucydides.** BOOKS III. IV. AND V. Edited by C. E. GRAVES, M.A., Classical Lecturer, and late Fellow of St. John's College, Cambridge. (To be published separately.)

*[Book IV. in the press.]*

BOOKS VI. AND VII. THE SICILIAN EXPEDITION. Edited by the Rev. PERCIVAL FROST, M.A., late Fellow of St. John's College, Cambridge. New Edition, revised and enlarged, with Map. 5s.

**Tibullus.**—SELECTIONS. Edited by Professor J. P. POSTGATE, M.A. *[In preparation.]*

**Virgil.**—ÆNEID. BOOKS II. AND III. THE NARRATIVE OF ÆNEAS. Edited by E. W. HOWSON, M.A., Fellow of King's College, Cambridge, and Assistant-Master at Harrow. 3s.

**Xenophon.**—HELLENICA, BOOKS I. AND II. Edited by H. HAILSTONE, B.A., late Scholar of Peterhouse, Cambridge. With Map. 4s. 6d.

CYROPÆDIA. BOOKS VII. AND VIII. Edited by ALFRED GOODWIN, M.A., Professor of Greek in University College, London. 5s.

MEMORABILIA SOCRATIS. Edited by A. R. CLUER, B.A. Balliol College, Oxford. 6s.

THE ANABASIS. BOOKS I.—IV. Edited by Professors W. W. GOODWIN and J. W. WHITE. Adapted to Goodwin's Greek Grammar. With a Map. 5s.

**Xenophon.**—ECONOMICUS. Edited by Rev. H. A. HOLDEN, M.A., LL.D. *[In the press.]*

**HIERO.** By the same Editor. *[In the press.]*

*\*\*\* Other Volumes will follow.*

---

CLASSICAL LIBRARY.

(1) **Texts**, Edited with **Introductions and Notes**, for the use of Advanced Students. (2) **Commentaries and Translations**.

**Æschylus.**—THE EUMENIDES. The Greek Text, with Introduction, English Notes, and Verse Translation. By BERNARD DRAKE, M.A., late Fellow of King's College, Cambridge. 8vo. 5s.

**AGAMEMNON, CHOEPHORCE, AND EUMENIDES.** Edited, with Introduction and Notes, by A. O. PRICKARD, M.A., Fellow and Tutor of New College, Oxford. 8vo. *[In preparation.]*

**Antoninus, Marcus Aurelius.**—BOOK IV. OF THE MEDITATIONS. The Text Revised, with Translation and Notes, by HASTINGS CROSSLEY, M.A., Professor of Greek in Queen's College, Belfast. 8vo. 6s.

**Aristotle.**—THE METAPHYSICS. BOOK I. Translated by a Cambridge Graduate. 8vo. 5s. *[Book II. in preparation.]*

**THE POLITICS.** Edited, after SUSEMIHL, by R. D. HICKS, M.A., Fellow of Trinity College, Cambridge. 8vo. *[In the press.]*

**THE POLITICS.** Translated by J. E. C. WELLDON, M.A., Fellow of King's College, Cambridge, and Master of Dulwich College. Crown 8vo. 1cs. 6d.

**AN INTRODUCTION TO ARISTOTLE'S RHETORIC.** With Analysis, Notes, and Appendices. By E. M. COPE, Fellow and Tutor of Trinity College, Cambridge. 8vo. 14s.

**THE RHETORIC.** Translated by J. E. C. WELLDON, M.A. *[In preparation.]*

**THE SOPHISTICI ELENCHI.** With Translation and Notes by E. POSTE, M.A., Fellow of Oriel College, Oxford. 8vo. 8s. 6d.



**Aristophanes.**—THE BIRDS. Translated into English Verse, with Introduction, Notes, and Appendices, by B. H. KENNEDY, D.D., Regius Professor of Greek in the University of Cambridge. Crown 8vo. 6s. Help Notes to the same, for the use of Students, 1s. 6d.

**Attic Orators.**—FROM ANTIPHON TO ISÆEOS. By R. C. JEBB, M.A., LL.D., Professor of Greek in the University of Glasgow. 2 vols. 8vo. 25s.

SELECTIONS FROM ANTIPHON, ANDOKIDES, LYSIAS, ISOKRATES, AND ISÆEOS. Edited, with Notes, by Professor JEBB. Being a companion volume to the preceding work. 8vo. 12s. 6d.

**Babrius.** Edited, with Introductory Dissertations, Critical Notes, Commentary and Lexicon. By W. GUNION RUTHERFORD, M.A., Head-Master of Westminster School. 8vo. 12s. 6d.

**Cicero.**—THE ACADEMICA. The Text revised and explained by J. S. REID, M.L., Fellow of Caius College, Cambridge. New Edition. With Translation. 8vo. [In the press.]

THE ACADEMICS. Translated by J. S. REID, M.L. 8vo. 5s. 6d.

SELECT LETTERS. After the Edition of ALBERT WATSON, M.A. Translated by G. E. JEANS, M.A., Fellow of Hertford College, Oxford, and Assistant-Master at Haileybury. 8vo. 10s. 6d.

(See also *Classical Series.*)

**Euripides.**—MEDEA. Edited, with Introduction and Notes, by A. W. VERRALL, M.A., Fellow and Lecturer of Trinity College, Cambridge. 8vo. 7s. 6d.

INTRODUCTION TO THE STUDY OF EURIPIDES. By Professor J. P. MAHAFFY. Fcap. 8vo. 1s. 6d. (*Classical Writers Series.*)

(See also *Classical Series.*)

**Herodotus.**—BOOKS I.—III. THE EMPIRES OF THE EAST. Edited, with Notes and Introductions, by Rev. A. H. SAYCE, M.A., LL.D., Fellow of Queen's College, Oxford, and Deputy-Professor of Comparative Philology. 8vo.

[Nearly ready.]

BOOKS IV.—IX. Edited by REGINALD W. MACAN, M.A., Lecturer in Ancient History at Brasenose College, Oxford. 8vo.

[In preparation.]

- Homer.**—THE ILIAD. Edited, with Introduction and Notes, by WALTER LEAF, M.A., Fellow of Trinity College, Cambridge, and the late J. H. PRATT, M.A. 8vo. [In preparation.]
- THE ILIAD. Translated into English Prose. By ANDREW LANG, M.A., WALTER LEAF, M.A., and ERNEST MYERS, M.A. Crown 8vo. 12s. 6d.
- THE ODYSSEY. Done into English by S. H. BUTCHER, M.A., Professor of Greek in the University of Edinburgh, and ANDREW LANG, M.A., late Fellow of Merton College, Oxford. Fourth Edition, revised and corrected. Crown 8vo. 10s. 6d.
- INTRODUCTION TO THE STUDY OF HOMER. By the Right Hon. W. E. GLADSTONE, M.P. 18mo. 1s. (*Literature Primers.*)
- HOMERIC DICTIONARY. For Use in Schools and Colleges. Translated from the German of Dr. G. AUTENRIETH, with Additions and Corrections, by R. P. KEEP, Ph.D. With numerous Illustrations. Crown 8vo. 6s.  
(See also *Classical Series.*)
- Horace.**—THE WORKS OF HORACE RENDERED INTO ENGLISH PROSE. With Introductions, Running Analysis, Notes, &c. By J. LONSDALE, M.A., and S. LEE, M.A. (*Globe Edition.*) 3s. 6d.  
(See also *Classical Series.*)
- Juvenal.**—THIRTEEN SATIRES OF JUVENAL. With a Commentary. By JOHN E. B. MAYOR, M.A., Professor of Latin in the University of Cambridge. Second Edition, enlarged. Crown 8vo. Vol. I. 7s. 6d. Vol. II. 10s. 6d.
- THIRTEEN SATIRES. Translated into English after the Text of J. E. B. MAYOR by HERBERT STRONG, M.A., Professor of Latin, and ALEXANDER LEEPER, M.A., Warden of Trinity College, in the University of Melbourne. Crown 8vo. 3s. 6d.  
(See also *Classical Series.*)
- Livy.** BOOKS XXI.—XXV. Translated by ALFRED JOHN CHURCH, M.A., of Lincoln College, Oxford, Professor of Latin, University College, London, and WILLIAM JACKSON BRODRIBB, M.A., late Fellow of St. John's College, Cambridge. Crown 8vo. 7s. 6d.
- INTRODUCTION TO THE STUDY OF LIVY. By Rev. W. W. CAPES, Reader in Ancient History at Oxford. Fcap. 8vo. 1s. 6d. (*Classical Writers Series.*)  
(See also *Classical Series.*)

**Blackie.**—GREEK AND ENGLISH DIALOGUES FOR USE IN SCHOOLS AND COLLEGES. By JOHN STUART BLACKIE, Emeritus Professor of Greek in the University of Edinburgh. New Edition. Fcap. 8vo. 2s. 6d.

**Ellis.**—PRACTICAL HINTS ON THE QUANTITATIVE PRONUNCIATION OF LATIN, for the use of Classical Teachers and Linguists. By A. J. ELLIS, B.A., F.R.S. Extra fcap. 8vo. 4s. 6d.

**England.**—EXERCISES ON LATIN SYNTAX AND IDIOM, ARRANGED WITH REFERENCE TO ROBY'S SCHOOL LATIN GRAMMAR. By E. B. ENGLAND, M.A., Assistant Lecturer at the Owens College, Manchester. Crown 8vo. 2s. 6d. Key for Teachers only, 2s. 6d.

**Goodwin.**—Works by W. W. GOODWIN, LL.D., Professor of Greek in Harvard University, U.S.A.

SYNTAX OF THE MOODS AND TENSES OF THE GREEK VERB. New Edition, revised. Crown 8vo. 6s. 6d.

A GREEK GRAMMAR. New Edition, revised. Crown 8vo. 6s.

"It is the best Greek Grammar of its size in the English language."—*ATHENÆUM*.

A GREEK GRAMMAR FOR SCHOOLS. Crown 8vo. 3s. 6d.

**Greenwood.**—THE ELEMENTS OF GREEK GRAMMAR, including Accidence, Irregular Verbs, and Principles of Derivation and Composition; adapted to the System of Crude Forms. By J. G. GREENWOOD, Principal of Owens College, Manchester. New Edition. Crown 8vo. 5s. 6d.

**Hodgson.**—MYTHOLOGY FOR LATIN VERSIFICATION. A brief Sketch of the Fables of the Ancients, prepared to be rendered into Latin Verse for Schools. By F. HODGSON, B.D., late Provost of Eton. New Edition, revised by F. C. HODGSON, M.A. 18mo. 3s.

**Jackson.**—FIRST STEPS TO GREEK PROSE COMPOSITION. By BLOMFIELD JACKSON, M.A., Assistant-Master in King's College School, London. New Edition, revised and enlarged. 18mo. 1s. 6d.

KEY TO FIRST STEPS (for Teachers only). 18mo. 3s. 6d.

**Jackson** (*continued*)—

**SECOND STEPS TO GREEK PROSE COMPOSITION**, with Miscellaneous Idioms, Aids to Accentuation, and Examination Papers in Greek Scholarship. 18mo. 2s. 6d.

**KEY TO SECOND STEPS** (for Teachers only). 18mo. 3s. 6d.

**Kynaston**.—**EXERCISES IN THE COMPOSITION OF GREEK IAMBIC VERSE** by Translations from English Dramatists. By Rev. H. KYNASTON, M.A., Principal of Cheltenham College. With Introduction, Vocabulary, &c. Extra fcap. 8vo. 4s. 6d.

**KEY TO THE SAME** (for Teachers only). Extra fcap. 8vo. 4s. 6d.

**Macmillan**.—**FIRST LATIN GRAMMAR**. By M. C. MACMILLAN, M.A., late Scholar of Christ's College, Cambridge; sometime Assistant-Master in St. Paul's School. New Edition, enlarged. 18mo. 1s. 6d. A **SHORT SYNTAX** is in preparation to follow the **ACCIDENCE**.

**Marshall**.—**A TABLE OF IRREGULAR GREEK VERBS**, classified according to the arrangement of Curtius's Greek Grammar. By J. M. MARSHALL, M.A., one of the Masters in Clifton College. 8vo, cloth. New Edition. 1s.

**Mayor (John E. B.)**.—**FIRST GREEK READER**. Edited after KARL HALM, with Corrections and large Additions by Professor JOHN E. B. MAYOR, M.A., Fellow of St. John's College, Cambridge. New Edition, revised. Fcap. 8vo. 4s. 6d.

**Mayor (Joseph B.)**.—**GREEK FOR BEGINNERS**. By the Rev. J. B. MAYOR, M.A., Professor of Classical Literature in King's College, London. Part I., with Vocabulary, 1s. 6d. Parts II. and III., with Vocabulary and Index, 3s. 6d. Complete in one Vol. fcap. 8vo. 4s. 6d.

**Nixon**.—**PARALLEL EXTRACTS** arranged for translation into English and Latin, with Notes on Idioms. By J. E. NIXON, M.A., Fellow and Classical Lecturer, King's College, Cambridge. Part I.—Historical and Epistolary. New edition, revised and enlarged. Crown 8vo. 3s. 6d.

**Postgate and Vince**.—**A DICTIONARY OF LATIN ETYMOLOGY**. By J. P. POSTGATE, M.A., and C. A. VINCE, M.A. [*In preparation.*]



**Potts (A. W.)**.—Works by ALEXANDER W. POTTS, M.A., LL.D., late Fellow of St. John's College, Cambridge; Head-Master of the Fettes College, Edinburgh.

**HINTS TOWARDS LATIN PROSE COMPOSITION.** New Edition. Extra fcap. 8vo. 3s.

**PASSAGES FOR TRANSLATION INTO LATIN PROSE.** Edited with Notes and References to the above. New Edition. Extra fcap. 8vo. 2s. 6d.

**LATIN VERSIONS OF PASSAGES FOR TRANSLATION INTO LATIN PROSE** (for Teachers only). 2s. 6d.

**Roby.**—A GRAMMAR OF THE LATIN LANGUAGE, from Plautus to Suetonius. By H. J. ROBY, M.A., late Fellow of St. John's College, Cambridge. In Two Parts. Third Edition. Part I. containing:—Book I. Sounds. Book II. Inflections. Book III. Word-formation. Appendices. Crown 8vo. 8s. 6d. Part II. Syntax, Prepositions, &c. Crown 8vo. 10s. 6d.

"Marked by the clear and practised insight of a master in his art. A book that would do honour to any country."—*ATHENÆUM*.

**SCHOOL LATIN GRAMMAR.** By the same Author. Crown 8vo. 5s.

**Rush.**—SYNTHETIC LATIN DELECTUS. A First Latin Construing Book arranged on the Principles of Grammatical Analysis. With Notes and Vocabulary. By E. RUSH, B.A. With Preface by the Rev. W. F. MOULTON, M.A., D.D. New and Enlarged Edition. Extra fcap. 8vo. 2s. 6d.

**Rust.**—FIRST STEPS TO LATIN PROSE COMPOSITION. By the Rev. G. RUST, M.A., of Pembroke College, Oxford, Master of the Lower School, King's College, London. New Edition. 18mo. 1s. 6d.

**Rutherford.**—Works by W. GUNION RUTHERFORD, M.A., Head-Master of Westminster School.

**A FIRST GREEK GRAMMAR.** New Edition, enlarged. Extra fcap. 8vo. 1s. 6d.

**THE NEW PHRYNICHUS;** being a Revised Text of the Ecloga of the Grammarian Phrynichus. With Introduction and Commentary. 8vo. 18s.

**Thring.**—Works by the Rev. E. THRING, M.A., Head-Master of Uppingham School.

**A LATIN GRADUAL.** A First Latin Construing Book for Beginners. New Edition, enlarged, with Coloured Sentence Maps. Fcap. 8vo. 2s. 6d.

**A MANUAL OF MOOD CONSTRUCTIONS.** Fcap. 8vo. 1s. 6d.

**White.**—FIRST LESSONS IN GREEK. Adapted to GOODWIN'S GREEK GRAMMAR, and designed as an introduction to the ANABASIS OF XENOPHON. By JOHN WILLIAMS WHITE, Ph.D., Assistant-Professor of Greek in Harvard University. Crown 8vo. 4s. 6d.

**Wright.**—Works by J. WRIGHT, M.A., late Head Master of Sutton Coldfield School.

**A HELP TO LATIN GRAMMAR ;** or, The Form and Use of Words in Latin, with Progressive Exercises. Crown 8vo. 4s. 6d.

**THE SEVEN KINGS OF ROME.** An Easy Narrative, abridged from the First Book of Livy by the omission of Difficult Passages ; being a First Latin Reading Book, with Grammatical Notes and Vocabulary. New and revised Edition. Fcap. 8vo. 3s. 6d.

**FIRST LATIN STEPS ; OR, AN INTRODUCTION BY A SERIES OF EXAMPLES TO THE STUDY OF THE LATIN LANGUAGE.** Crown 8vo. 3s.

**ATTIC PRIMER.** Arranged for the Use of Beginners. Extra fcap. 8vo. 2s. 6d.

**A COMPLETE LATIN COURSE,** comprising Rules with Examples, Exercises, both Latin and English, on each Rule and Vocabulary. Crown 8vo. 2s. 6d.

**Wright (H. C.)**—EXERCISES ON THE LATIN SYNTAX. By H. C. WRIGHT, B.A., Assistant-Master at Haileybury College. 18mo. [In preparation.]

## ANTIQUITIES, ANCIENT HISTORY, AND PHILOSOPHY.

**Arnold.**—Works by W. T. ARNOLD, B.A.

A HANDBOOK OF LATIN EPIGRAPHY. [*In preparation.*]

THE ROMAN SYSTEM OF PROVINCIAL ADMINISTRATION TO THE ACCESSION OF CONSTANTINE THE GREAT. Crown 8vo. 6s.

**Beesly.**—STORIES FROM THE HISTORY OF ROME.

By Mrs. BEESLEY. Fcap. 8vo. 2s. 6d.

**Classical Writers.**—Edited by JOHN RICHARD GREEN, M.A., LL.D. Fcap. 8vo. 1s. 6d. each.

EURIPIDES. By Professor MAHAFFY.

MILTON. By the Rev. STOPFORD A. BROOKE, M.A.

LIVY. By the Rev. W. W. CAPES, M.A.

VIRGIL. By Professor NETTLESHIP, M.A.

SOPHOCLES. By Professor L. CAMPBELL, M.A.

DEMOSTHENES. By Professor S. H. BUTCHER, M.A.

TACITUS. By Professor A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A.

**Freeman.**—HISTORY OF ROME. By EDWARD A. FREEMAN, D.C.L., LL.D., Hon. Fellow of Trinity College, Oxford. (*Historical Course for Schools.*) 18mo. [*In the press.*]

**Fyffe.**—A SCHOOL HISTORY OF GREECE. By C. A. FYFFE, M.A., late Fellow of University College, Oxford. Crown 8vo. [*In preparation.*]

**Geddes.**—THE PROBLEM OF THE HOMERIC POEMS. By W. D. GEDDES, Professor of Greek in the University of Aberdeen. 8vo. 14s.

**Gladstone.**—Works by the Rt. Hon. W. E. GLADSTONE, M.P.

THE TIME AND PLACE OF HOMER. Crown 8vo. 6s. 6d.

A PRIMER OF HOMER. 18mo. 1s.

**Goodwin.**—A TEXT-BOOK OF GREEK PHILOSOPHY, based on RITTER and PRELLER'S "Historia Philosophiae Graecae et Romanae." By ALFRED GOODWIN, M.A., Fellow of Balliol College, Oxford, and Professor of Greek in University College, London. 8vo. [*In preparation.*]

**Jackson.**—A MANUAL OF GREEK PHILOSOPHY. By HENRY JACKSON, M.A., Fellow and Prælector in Ancient Philosophy, Trinity College, Cambridge. [*In preparation.*]

**Jebb.**—Works by R. C. JEBB, M.A., Professor of Greek in the University of Glasgow.

THE ATTIC ORATORS FROM ANTIPHON TO ISAEOS. 2 vols. 8vo. 25s.

SELECTIONS FROM THE ATTIC ORATORS, ANTIPHON, ANDOKIDES, LYSIAS, ISOKRATES, AND ISÆOS. Edited, with Notes. Being a companion volume to the preceding work. 8vo. 12s. 6d.

A PRIMER OF GREEK LITERATURE. 18mo. 1s.

**Kiepert.**—MANUAL OF ANCIENT GEOGRAPHY, Translated from the German of Dr. HEINRICH KIEPERT. Crown 8vo. 5s.

**Mahaffy.**—Works by J. P. MAHAFFY, M.A., Professor of Ancient History in Trinity College, Dublin, and Hon. Fellow of Queen's College, Oxford.

SOCIAL LIFE IN GREECE; from Homer to Menander. Fourth Edition, revised and enlarged. Crown 8vo. 9s.

RAMBLES AND STUDIES IN GREECE. With Illustrations. Second Edition. With Map. Crown 8vo. 10s. 6d.

A PRIMER OF GREEK ANTIQUITIES. With Illustrations. 18mo. 1s.

EURIPIDES. 18mo. 1s. 6d. (*Classical Writers Series.*)

**Mayor (J. E. B.)**—BIBLIOGRAPHICAL CLUE TO LATIN LITERATURE. Edited after HÜBNER, with large Additions by Professor JOHN E. B. MAYOR. Crown 8vo. 10s. 6d.

**Ramsay.**—A SCHOOL HISTORY OF ROME. By G. G. RAMSAY, M.A., Professor of Humanity in the University of Glasgow. With Maps. Crown 8vo. [*In preparation.*]

**Wilkins.**—A PRIMER OF ROMAN ANTIQUITIES. By Professor WILKINS. Illustrated. 18mo. 1s.

## MATHEMATICS.

(1) Arithmetic, (2) Algebra, (3) Euclid and Elementary Geometry, (4) Mensuration, (5) Higher Mathematics.

### ARITHMETIC.

**Aldis.**—THE GIANT ARITHMOS. A most Elementary Arithmetic for Children. By MARY STEADMAN ALDIS. With Illustrations. Globe 8vo. 2s. 6d.

**Brook-Smith (J.).**—ARITHMETIC IN THEORY AND PRACTICE. By J. BROOK-SMITH, M.A., LL.B., St. John's College, Cambridge; Barrister-at-Law; one of the Masters of Cheltenham College. New Edition, revised. Crown 8vo. 4s. 6d.

**Candler.**—HELP TO ARITHMETIC. Designed for the use of Schools. By H. CANDLER, M.A., Mathematical Master of Uppingham School. Extra fcap. 8vo. 2s. 6d.

**Dalton.**—RULES AND EXAMPLES IN ARITHMETIC. By the Rev. T. DALTON, M.A., Assistant-Master of Eton College. New Edition. 18mo. 2s. 6d.

*[Answers to the Examples are appended.]*

**Pedley.**—EXERCISES IN ARITHMETIC for the Use of Schools. Containing more than 7,000 original Examples. By S. PEDLEY, late of Tamworth Grammar School. Crown 8vo. 5s.

**Smith.**—Works by the Rev. BARNARD SMITH, M.A., late Rector of Glaston, Rutland, and Fellow and Senior Bursar of S. Peter's College, Cambridge.

ARITHMETIC AND ALGEBRA, in their Principles and Application; with numerous systematically arranged Examples taken from the Cambridge Examination Papers, with especial reference to the Ordinary Examination for the B.A. Degree. New Edition, carefully Revised. Crown 8vo. 10s. 6d.

ARITHMETIC FOR SCHOOLS. New Edition. Crown 8vo. 4s. 6d.

A KEY TO THE ARITHMETIC FOR SCHOOLS. New Edition. Crown 8vo. 8s. 6d.

**Smith.**—Works by the Rev. BARNARD SMITH, M.A. (*continued*)—

EXERCISES IN ARITHMETIC. Crown 8vo, limp cloth, 2s.  
With Answers, 2s. 6d.

Answers separately, 6d.

SCHOOL CLASS-BOOK OF ARITHMETIC. 18mo, cloth. 3s.  
Or sold separately, in Three Parts, 1s. each.

KEYS TO SCHOOL CLASS-BOOK OF ARITHMETIC.  
Parts I., II., and III., 2s. 6d. each.

SHILLING BOOK OF ARITHMETIC FOR NATIONAL  
AND ELEMENTARY SCHOOLS. 18mo, cloth. Or sepa-  
rately, Part I. 2d. ; Part II. 3d. ; Part III. 7d. Answers. 6d.

THE SAME, with Answers complete. 18mo, cloth. 1s. 6d.

KEY TO SHILLING BOOK OF ARITHMETIC. 18mo. 4s. 6d.

EXAMINATION PAPERS IN ARITHMETIC. 18mo. 1s. 6d.  
The same, with Answers, 18mo, 2s. Answers, 6d.

KEY TO EXAMINATION PAPERS IN ARITHMETIC.  
18mo. 4s. 6d.

THE METRIC SYSTEM OF ARITHMETIC, ITS PRIN-  
CIPLES AND APPLICATIONS, with numerous Examples,  
written expressly for Standard V. in National Schools. New  
Edition. 18mo, cloth, sewed. 3d.

A CHART OF THE METRIC SYSTEM, on a Sheet, size 42 in.  
by 34 in. on Roller, mounted and varnished. New Edition.  
Price 3s. 6d.

Also a Small Chart on a Card, price 1d.

EASY LESSONS IN ARITHMETIC, combining Exercises in  
Reading, Writing, Spelling, and Dictation. Part I. for Standard  
I. in National Schools. Crown 8vo. 9d.

EXAMINATION CARDS IN ARITHMETIC. (Dedicated to  
Lord Sandon.) With Answers and Hints.

Standards I. and II. in box, 1s. Standards III., IV., and V., in  
boxes, 1s. each. Standard VI. in Two Parts, in boxes, 1s. each.

A and B papers, of nearly the same difficulty, are given so as to  
prevent copying, and the colours of the A and B papers differ in each  
Standard, and from those of every other Standard, so that a master  
or mistress can see at a glance whether the children have the proper  
papers.

## ALGEBRA.

**Dalton.**—RULES AND EXAMPLES IN ALGEBRA. By the Rev. T. DALTON, M.A., Assistant-Master of Eton College. Part I. New Edition. 18mo. 2s. Part II. 18mo. 2s. 6d.

**Jones and Cheyne.**—ALGEBRAICAL EXERCISES. Progressively Arranged. By the Rev. C. A. JONES, M.A., and C. H. CHEYNE, M.A., F.R.A.S., Mathematical Masters of Westminster School. New Edition. 18mo. 2s. 6d.

**Smith.**—ARITHMETIC AND ALGEBRA, in their Principles and Application; with numerous systematically arranged Examples taken from the Cambridge Examination Papers, with especial reference to the Ordinary Examination for the B.A. Degree. By the Rev. BARNARD SMITH, M.A., late Rector of Glaston, Rutland, and Fellow and Senior Bursar of St. Peter's College, Cambridge. New Edition, carefully Revised. Crown 8vo. 10s. 6d.

**Todhunter.**—Works by I. TODHUNTER, M.A., F.R.S., D.Sc., of St. John's College, Cambridge.

"Mr. Todhunter is chiefly known to Students of Mathematics as the author of a series of admirable mathematical text-books, which possess the rare qualities of being clear in style and absolutely free from mistakes, typographical or other."—SATURDAY REVIEW.

ALGEBRA FOR BEGINNERS. With numerous Examples. New Edition. 18mo. 2s. 6d.

KEY TO ALGEBRA FOR BEGINNERS. Crown 8vo. 6s. 6d.

ALGEBRA. For the Use of Colleges and Schools. New Edition. Crown 8vo. 7s. 6d.

KEY TO ALGEBRA FOR THE USE OF COLLEGES AND SCHOOLS. Crown 8vo. 10s. 6d.

## EUCLID & ELEMENTARY GEOMETRY.

**Constable.**—GEOMETRICAL EXERCISES FOR BEGINNERS. By SAMUEL CONSTABLE. Crown 8vo. 3s. 6d.

**Cuthbertson.**—EUCLIDIAN GEOMETRY. By FRANCIS CUTHBERTSON, M.A., LL.D., Head Mathematical Master of the City of London School. Extra fcap. 8vo. 4s. 6d.



**Dodgson.**—EUCLID. BOOKS I. AND II. Edited by CHARLES L. DODGSON, M.A., Student and late Mathematical Lecturer of Christ Church, Oxford. Second Edition, with words substituted for the Algebraical Symbols used in the First Edition. Crown 8vo. 2s.

\*.\* The text of this Edition has been ascertained, by counting the words, to be *less than five-sevenths* of that contained in the ordinary editions.

**Kitchener.**—A GEOMETRICAL NOTE-BOOK, containing Easy Problems in Geometrical Drawing preparatory to the Study of Geometry. For the use of Schools. By F. E. KITCHENER, M.A., Mathematical Master at Rugby. New Edition. 4to. 2s.

**Mault.**—NATURAL GEOMETRY: an Introduction to the Logical Study of Mathematics. For Schools and Technical Classes. With Explanatory Models, based upon the Tachymetrical works of Ed. Lagout. By A. MAULT. 18mo. 1s.  
Models to Illustrate the above, in Box, 12s. 6d.

**Syllabus of Plane Geometry** (corresponding to Euclid, Books I.—VI.). Prepared by the Association for the Improvement of Geometrical Teaching. New Edition. Crown 8vo. 1s.

**Todhunter.**—THE ELEMENTS OF EUCLID. For the Use of Colleges and Schools. By I. TODHUNTER, M.A., F.R.S., D.Sc., of St. John's College, Cambridge. New Edition. 18mo. 3s. 6d.

KEY TO EXERCISES IN EUCLID. Crown 8vo. 6s. 6d.

**Wilson (J. M.).**—ELEMENTARY GEOMETRY. BOOKS I.—V. Containing the Subjects of Euclid's first Six Books. Following the Syllabus of the Geometrical Association. By the Rev. J. M. WILSON, M.A., Head Master of Clifton College. New Edition. Extra fcap. 8vo. 4s. 6d.

## MENSURATION.

**Tebay.**—ELEMENTARY MENSURATION FOR SCHOOLS. With numerous examples. By SEPTIMUS TEBAY, B.A., Head Master of Queen Elizabeth's Grammar School, Rivington. Extra fcap. 8vo. 3s. 6d.

**Todhunter.**—MENSURATION FOR BEGINNERS. By I. TODHUNTER, M.A., F.R.S., D.Sc., of St. John's College, Cambridge. With numerous Examples. New Edition. 18mo. 2s. 6d.



## HIGHER MATHEMATICS.

**Airy.**—Works by Sir G. B. AIRY, K.C.B., formerly Astronomer-Royal:—

**ELEMENTARY TREATISE ON PARTIAL DIFFERENTIAL EQUATIONS.** Designed for the Use of Students in the Universities. With Diagrams. Second Edition. Crown 8vo. 5s. 6d.

**ON THE ALGEBRAICAL AND NUMERICAL THEORY OF ERRORS OF OBSERVATIONS AND THE COMBINATION OF OBSERVATIONS.** Second Edition, revised. Crown 8vo. 6s. 6d.

**Alexander (T.).**—**ELEMENTARY APPLIED MECHANICS.**

Being the simpler and more practical Cases of Stress and Strain wrought out individually from first principles by means of Elementary Mathematics. By T. ALEXANDER, C.E., Professor of Civil Engineering in the Imperial College of Engineering, Tokyo, Japan. Crown 8vo. Part I. 4s. 6d.

**Alexander and Thomson.**—**ELEMENTARY APPLIED**

**MECHANICS.** By THOMAS ALEXANDER, C.E., Professor of Engineering in the Imperial College of Engineering, Tokyo, Japan; and ARTHUR WATSON THOMSON, C.E., B.Sc., Professor of Engineering at the Royal College, Cirencester. Part II. **TRANSVERSE STRESS.** Crown 8vo. 10s. 6d.

**Bayma.**—**THE ELEMENTS OF MOLECULAR MECHANICS.**

By JOSEPH BAYMA, S.J., Professor of Philosophy, Stonyhurst College. Demy 8vo. 10s. 6d.

**Beasley.**—**AN ELEMENTARY TREATISE ON PLANE**

**TRIGONOMETRY.** With Examples. By R. D. BEASLEY, M.A. Eighth Edition, revised and enlarged. Crown 8vo. 3s. 6d.

**Blackburn (Hugh).**—**ELEMENTS OF PLANE TRIGO-**

**NOMETRY,** for the use of the Junior Class in Mathematics in the University of Glasgow. By HUGH BLACKBURN, M.A., late Professor of Mathematics in the University of Glasgow. Globe 8vo. 1s. 6d.

**Boole.**—Works by G. BOOLE, D.C.L., F.R.S., late Professor of Mathematics in the Queen's University, Ireland.

**A TREATISE ON DIFFERENTIAL EQUATIONS.** Third and Revised Edition. Edited by I. TODHUNTER. Crown 8vo. 14s.

- Boole.**—Works by G. BOOLE, D.C.L., &c. (*continued*)—  
 A TREATISE ON DIFFERENTIAL EQUATIONS. Supplementary Volume. Edited by I. TODHUNTER. Crown 8vo. 8s. 6d.  
 THE CALCULUS OF FINITE DIFFERENCES. Third Edition, revised by J. F. MOULTON. Crown 8vo. 10s. 6d.
- Cambridge Senate-House Problems and Riders, with Solutions:—**  
 1875—PROBLEMS AND RIDERS. By A. G. GREENHILL, M.A. Crown 8vo. 8s. 6d.  
 1878—SOLUTIONS OF SENATE-HOUSE PROBLEMS. By the Mathematical Moderators and Examiners. Edited by J. W. L. GLAISHER, M.A., Fellow of Trinity College, Cambridge. 12s.
- Cheyne.**—AN ELEMENTARY TREATISE ON THE PLANETARY THEORY. By C. H. H. CHEYNE, M.A., F.R.A.S. With a Collection of Problems. Second Edition. Crown 8vo. 6s. 6d.
- Christie.**—A COLLECTION OF ELEMENTARY TEST-QUESTIONS IN PURE AND MIXED MATHEMATICS; with Answers and Appendices on Synthetic Division, and on the Solution of Numerical Equations by Horner's Method. By JAMES R. CHRISTIE, F.R.S., Royal Military Academy, Woolwich. Crown 8vo. 8s. 6d.
- Clausius.**—MECHANICAL THEORY OF HEAT. By R. CLAUDIUS. Translated by WALTER R. BROWNE, M.A., late Fellow of Trinity College, Cambridge. Crown 8vo. 10s. 6d.
- Clifford.**—THE ELEMENTS OF DYNAMIC. An Introduction to the Study of Motion and Rest in Solid and Fluid Bodies. By W. K. CLIFFORD, F.R.S., late Professor of Applied Mathematics and Mechanics at University College, London. Part I.—KINEMATIC. Crown 8vo. 7s. 6d.
- Cotterill.**—A TREATISE ON APPLIED MECHANICS. By JAMES COTTERILL, M.A., F.R.S., Professor of Applied Mechanics at the Royal Naval College, Greenwich. With Illustrations. 8vo. [In the press.]
- Day.**—PROPERTIES OF CONIC SECTIONS PROVED GEOMETRICALLY. Part I. THE ELLIPSE. With Problems. By the Rev. H. G. DAY, M.A. 8vo. 3s. 6d.
- Day (R. E.)**—ELECTRIC LIGHT ARITHMETIC. By R. E. DAY, M.A., Evening Lecturer in Experimental Physics at King's College, London. Pott 8vo. 2s.

**Drew.**—GEOMETRICAL TREATISE ON CONIC SECTIONS. By W. H. DREW, M.A., St. John's College, Cambridge. New Edition, enlarged. Crown 8vo. 5s.

SOLUTIONS TO THE PROBLEMS IN DREW'S CONIC SECTIONS. Crown 8vo. 4s. 6d.

**Dyer.**—EXERCISES IN ANALYTICAL GEOMETRY. Compiled and arranged by J. M. DYER, M.A., Senior Mathematical Master in the Classical Department of Cheltenham College. With Illustrations. Crown 8vo. 4s. 6d.

**Edgar (J. H.) and Pritchard (G. S.).**—NOTE-BOOK ON PRACTICAL SOLID OR DESCRIPTIVE GEOMETRY. Containing Problems with help for Solutions. By J. H. EDGAR, M.A., Lecturer on Mechanical Drawing at the Royal School of Mines, and G. S. PRITCHARD. Fourth Edition, revised by ARTHUR MEEZE. Globe 8vo. 4s. 6d.

**Ferrers.**—Works by the Rev. N. M. FERRERS, M.A., Fellow and Master of Gonville and Caius College, Cambridge.

AN ELEMENTARY TREATISE ON TRILINEAR COORDINATES, the Method of Reciprocal Polars, and the Theory of Projectors. New Edition, revised. Crown 8vo. 6s. 6d.

AN ELEMENTARY TREATISE ON SPHERICAL HARMONICS, AND SUBJECTS CONNECTED WITH THEM. Crown 8vo. 7s. 6d.

**Frost.**—Works by PERCIVAL FROST, M.A., D.Sc., formerly Fellow of St. John's College, Cambridge; Mathematical Lecturer at King's College.

AN ELEMENTARY TREATISE ON CURVE TRACING. By PERCIVAL FROST, M.A. 8vo. 12s.

SOLID GEOMETRY. A New Edition, revised and enlarged, of the Treatise by FROST and WOLSTENHOLME. In 2 Vols. Vol. I. 8vo. 16s.

**Hemming.**—AN ELEMENTARY TREATISE ON THE DIFFERENTIAL AND INTEGRAL CALCULUS, for the Use of Colleges and Schools. By G. W. HEMMING, M.A., Fellow of St. John's College, Cambridge. Second Edition, with Corrections and Additions. 8vo. 9s.

**Jackson.**—GEOMETRICAL CONIC SECTIONS. An Elementary Treatise in which the Conic Sections are defined as the Plane Sections of a Cone, and treated by the Method of Projection. By J. STUART JACKSON, M.A., late Fellow of Gonville and Caius College, Cambridge. Crown 8vo. 4s. 6d.

- Jellet (John H.).**—A TREATISE ON THE THEORY OF FRICTION. By JOHN H. JELLET, B.D., Provost of Trinity College, Dublin; President of the Royal Irish Academy. 8vo. 8s. 6d.
- Kelland and Tait.**—INTRODUCTION TO QUATERNIONS, with numerous examples. By P. KELLAND, M.A., F.R.S., and P. G. TAIT, M.A., Professors in the Department of Mathematics in the University of Edinburgh. Second Edition. Crown 8vo. 7s. 6d.
- Kempe.**—HOW TO DRAW A STRAIGHT LINE: a Lecture on Linkages. By A. B. KEMPE. With Illustrations. Crown 8vo. 1s. 6d. (*Nature Series.*)
- Lock.**—ELEMENTARY TRIGONOMETRY. By Rev. J. B. LOCK, M.A., Fellow of Caius College, Cambridge; Assistant-Master at Eton. Globe 8vo. 4s. 6d.
- HIGHER TRIGONOMETRY.** By the same Author. Crown 8vo. [In the Press.]
- Lupton.**—ELEMENTARY CHEMICAL ARITHMETIC. With 1,100 Problems. By SYDNEY LUPTON, M.A., Assistant-Master in Harrow School. Globe 8vo. 5s.
- Merriman.**—ELEMENTS OF THE METHOD OF LEAST SQUARE. By MANSFIELD MERRIMAN, Ph.D., Professor of Civil and Mechanical Engineering, Lehigh University, Bethlehem, Penn. Crown 8vo. 7s. 6d.
- Morgan.**—A COLLECTION OF PROBLEMS AND EXAMPLES IN MATHEMATICS. With Answers. By H. A. MORGAN, M.A., Sadlerian and Mathematical Lecturer of Jesus College, Cambridge. Crown 8vo. 6s. 6d.
- Millar.**—ELEMENTS OF DESCRIPTIVE GEOMETRY. By J. B. MILLAR, C.E., Assistant Lecturer in Engineering in Owens College, Manchester. Crown 8vo. 6s.
- Muir.**—A TREATISE ON THE THEORY OF DETERMINANTS. With graduated sets of Examples. For use in Colleges and Schools. By THOS. MUIR, M.A., F.R.S.E., Mathematical Master in the High School of Glasgow. Crown 8vo. 7s. 6d.
- Parkinson.**—AN ELEMENTARY TREATISE ON MECHANICS. For the Use of the Junior Classes at the University and the Higher Classes in Schools. By S. PARKINSON, D.D., F.R.S., Tutor and Prælector of St. John's College, Cambridge. With a Collection of Examples. Sixth Edition, revised. Crown 8vo. 9s. 6d.

- Phear.**—ELEMENTARY HYDROSTATICS. With Numerous Examples. By J. B. PHEAR, M.A., Fellow and late Assistant Tutor of Clare College, Cambridge. New Edition. Crown 8vo. 5s. 6d.
- Pirie.**—LESSONS ON RIGID DYNAMICS. By the Rev. G. PIRIE, M.A., late Fellow and Tutor of Queen's College, Cambridge; Professor of Mathematics in the University of Aberdeen. Crown 8vo. 6s.
- Puckle.**—AN ELEMENTARY TREATISE ON CONIC SECTIONS AND ALGEBRAIC GEOMETRY. With Numerous Examples and Hints for their Solution; especially designed for the Use of Beginners. By G. H. PUCKLE, M.A. New Edition, revised and enlarged. Crown 8vo. 7s. 6d.
- Rawlinson.**—ELEMENTARY STATICS. By the Rev. GEORGE RAWLINSON, M.A. Edited by the Rev. EDWARD STURGES, M.A. Crown 8vo. 4s. 6d.
- Reynolds.**—MODERN METHODS IN ELEMENTARY GEOMETRY. By E. M. REYNOLDS, M.A., Mathematical Master in Clifton College. Crown 8vo. 3s. 6d.
- Reuleaux.**—THE KINEMATICS OF MACHINERY. Outlines of a Theory of Machines. By Professor F. REULEAUX. Translated and Edited by Professor A. B. W. KENNEDY, C.E. With 450 Illustrations. Medium 8vo. 21s.
- Robinson.**—TREATISE ON MARINE SURVEYING. Prepared for the use of younger Naval Officers. With Questions for Examinations and Exercises principally from the Papers of the Royal Naval College. With the results. By Rev. JOHN L. ROBINSON, Chaplain and Instructor in the Royal Naval College, Greenwich. With Illustrations. Crown 8vo. 7s. 6d.
- CONTENTS.—Symbols used in Charts and Surveying—The Construction and Use of Scales—Laying off Angles—Fixing Positions by Angles—Charts and Chart-Drawing—Instruments and Observing—Base Lines—Triangulation—Levelling—Tides and Tidal Observations—Soundings—Chronometers—Meridian Distances—Method of Plotting a Survey—Miscellaneous Exercises—Index.
- Routh.**—Works by EDWARD JOHN ROUTH, M.A., F.R.S., D.Sc., late Fellow and Assistant Tutor at St. Peter's College, Cambridge; Examiner in the University of London.
- A TREATISE ON THE DYNAMICS OF THE SYSTEM OF RIGID BODIES. With numerous Examples. Fourth and enlarged Edition. Two Vols. Vol. I.—Elementary Parts. 8vo. 14s. Vol. II.—The Higher Parts. 8vo. [In the press.]



**Routh**—Works by E. J. ROUTH, M.A., F.R.S., D.Sc. (*continued*)—  
STABILITY OF A GIVEN STATE OF MOTION, PARTICULARLY STEADY MOTION. Adams' Prize Essay for 1877. 8vo. 8s. 6d.

**Smith (C.)**—CONIC SECTIONS. By CHARLES SMITH, M.A.,  
Fellow and Tutor of Sidney Sussex College, Cambridge. Crown  
8vo. 7s. 6d.

**Snowball**—THE ELEMENTS OF PLANE AND SPHERICAL TRIGONOMETRY; with the Construction and Use of Tables of Logarithms. By J. C. SNOWBALL, M.A. New Edition. Crown 8vo. 7s. 6d.

**Tait and Steele**—A TREATISE ON DYNAMICS OF A PARTICLE. With numerous Examples. By Professor TAIT and Mr. STEELE. Fourth Edition, revised. Crown 8vo. 12s.

**Todhunter**—Works by I. TODHUNTER, M.A., F.R.S., D.Sc., of St. John's College, Cambridge.

“Mr. Todhunter is chiefly known to students of Mathematics as the author of a series of admirable mathematical text-books, which possess the rare qualities of being clear in style and absolutely free from mistakes, typographical and other.”—SATURDAY REVIEW.

TRIGONOMETRY FOR BEGINNERS. With numerous Examples. New Edition. 18mo. 2s. 6d.

KEY TO TRIGONOMETRY FOR BEGINNERS. Crown 8vo. 8s. 6d.

MECHANICS FOR BEGINNERS. With numerous Examples. New Edition. 18mo. 4s. 6d.

KEY TO MECHANICS FOR BEGINNERS. Crown 8vo. 6s. 6d.

AN ELEMENTARY TREATISE ON THE THEORY OF EQUATIONS. New Edition, revised. Crown 8vo. 7s. 6d.

PLANE TRIGONOMETRY. For Schools and Colleges. New Edition. Crown 8vo. 5s.

KEY TO PLANE TRIGONOMETRY. Crown 8vo. 10s. 6d.

A TREATISE ON SPHERICAL TRIGONOMETRY. New Edition, enlarged. Crown 8vo. 4s. 6d.

PLANE CO-ORDINATE GEOMETRY, as applied to the Straight Line and the Conic Sections. With numerous Examples. New Edition, revised and enlarged. Crown 8vo. 7s. 6d.

**Todhunter.**—Works by I. TODHUNTER, M.A., &c. (*continued*)—

A TREATISE ON THE DIFFERENTIAL CALCULUS. With numerous Examples. New Edition. Crown 8vo. 10s. 6d.

A TREATISE ON THE INTEGRAL CALCULUS AND ITS APPLICATIONS. With numerous Examples. New Edition, revised and enlarged. Crown 8vo. 10s. 6d.

EXAMPLES OF ANALYTICAL GEOMETRY OF THREE DIMENSIONS. New Edition, revised. Crown 8vo. 4s.

A TREATISE ON ANALYTICAL STATICS. With numerous Examples. New Edition, revised and enlarged. Crown 8vo. 10s. 6d.

A HISTORY OF THE MATHEMATICAL THEORY OF PROBABILITY, from the time of Pascal to that of Laplace. 8vo. 18s.

RESEARCHES IN THE CALCULUS OF VARIATIONS, principally on the Theory of Discontinuous Solutions: an Essay to which the Adams' Prize was awarded in the University of Cambridge in 1871. 8vo. 6s.

A HISTORY OF THE MATHEMATICAL THEORIES OF ATTRACTION, AND THE FIGURE OF THE EARTH, from the time of Newton to that of Laplace. 2 vols. 8vo. 24s.

AN ELEMENTARY TREATISE ON LAPLACE'S, LAME'S, AND BESSEL'S FUNCTIONS. Crown 8vo. 10s. 6d.

**Wilson (J. M.).**—SOLID GEOMETRY AND CONIC SECTIONS. With Appendices on Transversals and Harmonic Division. For the Use of Schools. By Rev. J. M. WILSON, M.A. Head Master of Clifton College. New Edition. Extra fcap. 8vo. 3s. 6d.

**Wilson.**—GRADUATED EXERCISES IN PLANE TRIGONOMETRY. Compiled and arranged by J. WILSON, M.A., and S. R. WILSON, B.A. Crown 8vo. 4s. 6d.

"The exercises seem beautifully graduated and adapted to lead a student on most gently and pleasantly."—E. J. ROUTH, F.R.S., St. Peter's College, Cambridge.

(See also *Elementary Geometry*.)

**Wilson (W. P.).**—A TREATISE ON DYNAMICS. By W. P. WILSON, M.A., Fellow of St. John's College, Cambridge, and Professor of Mathematics in Queen's College, Belfast. 8vo. 9s. 6d.

**Wolstenholme.**—**MATHEMATICAL PROBLEMS**, on Subjects included in the First and Second Divisions of the Schedule of subjects for the Cambridge Mathematical Tripos Examination. Devised and arranged by JOSEPH WOLSTENHOLME, D.Sc., late Fellow of Christ's College, sometime Fellow of St. John's College, and Professor of Mathematics in the Royal Indian Engineering College. New Edition, greatly enlarged. 8vo. 18s.

## SCIENCE.

(1) Natural Philosophy, (2) Astronomy, (3) Chemistry, (4) Biology, (5) Medicine, (6) Anthropology, (7) Physical Geography and Geology, (8) Agriculture, (9) Political Economy, (10) Mental and Moral Philosophy.

### NATURAL PHILOSOPHY.

**Airy.**—Works by Sir G. B. AIRY, K.C.B., formerly Astronomer-Royal :—

**UNDULATORY THEORY OF OPTICS.** Designed for the Use of Students in the University. New Edition. Crown 8vo. 6s. 6d.  
**ON SOUND AND ATMOSPHERIC VIBRATIONS.** With the Mathematical Elements of Music. Designed for the Use of Students in the University. Second Edition, revised and enlarged. Crown 8vo. 9s.

**A TREATISE ON MAGNETISM.** Designed for the Use of Students in the University. Crown 8vo. 9s. 6d.

**Airy (Osmond).**—**A TREATISE ON GEOMETRICAL OPTICS.** Adapted for the Use of the Higher Classes in Schools. By OSMUND AIRY, B.A., one of the Mathematical Masters in Wellington College. Extra fcap. 8vo. 3s. 6d.

**Alexander (T.).**—**ELEMENTARY APPLIED MECHANICS.** Being the simpler and more practical Cases of Stress and Strain wrought out individually from first principles by means of Elementary Mathematics. By T. ALEXANDER, C.E., Professor of Civil Engineering in the Imperial College of Engineering, Tokyo, Japan. Crown 8vo. Part I. 4s. 6d.



**Alexander — Thomson.** — **ELEMENTARY APPLIED MECHANICS.** By THOMAS ALEXANDER, C.E., Professor of Engineering in the Imperial College of Engineering, Tokei, Japan; and ARTHUR WATSON THOMSON, C.E., B.Sc., Professor of Engineering at the Royal College, Cirencester. Part II. TRANSVERSE STRESS; upwards of 150 Diagrams, and 200 Examples carefully worked out; new and complete method for finding, at every point of a beam, the amount of the greatest bending moment and shearing force during the transit of any set of loads fixed relatively to one another—*e.g.*, the wheels of a locomotive; continuous beams, &c., &c. Crown 8vo. 10s. 6d.

**Awdry.**—**EASY LESSONS ON LIGHT.** By Mrs. W. AWDRY. Illustrated. Extra fcap. 8vo. 2s. 6d.

**Ball (R. S.).**—**EXPERIMENTAL MECHANICS.** A Course of Lectures delivered at the Royal College of Science for Ireland. By R. S. BALL, M.A., Professor of Applied Mathematics and Mechanics in the Royal College of Science for Ireland. Cheaper Issue. Royal 8vo. 10s. 6d.

**Chisholm.**—**THE SCIENCE OF WEIGHING AND MEASURING, AND THE STANDARDS OF MEASURE AND WEIGHT.** By H.W. CHISHOLM, Warden of the Standards. With numerous Illustrations. Crown 8vo. 4s. 6d. (*Nature Series.*)

**Clausius.**—**MECHANICAL THEORY OF HEAT.** By R. CLAUSIUS. Translated by WALTER R. BROWNE, M.A., late Fellow of Trinity College, Cambridge. Crown 8vo. 10s. 6d.

**Cotterill.**—**A TREATISE ON APPLIED MECHANICS.** By JAMES COTTERILL, M.A., F.R.S., Professor of Applied Mechanics at the Royal Naval College, Greenwich. With Illustrations. 8vo. [*In the press.*]

**Cumming.**—**AN INTRODUCTION TO THE THEORY OF ELECTRICITY.** By LINNÆUS CUMMING, M.A., one of the Masters of Rugby School. With Illustrations. Crown 8vo. 8s. 6d.

**Daniell.**—**A TREATISE ON PHYSICS FOR MEDICAL STUDENTS.** By ALFRED DANIELL. With Illustrations. 8vo. [*In preparation.*]

**Day.**—**ELECTRIC LIGHT ARITHMETIC.** By R. E. DAY, M.A., Evening Lecturer in Experimental Physics at King's College, London. Pott 8vo. 2s.

**Everett.**—UNITS AND PHYSICAL CONSTANTS. By J. D. EVERETT, F.R.S., Professor of Natural Philosophy, Queen's College, Belfast. Extra fcap. 8vo. 4s. 6d.

**Huxley.**—INTRODUCTORY PRIMER OF SCIENCE. By T. H. HUXLEY, P.R.S., Professor of Natural History in the Royal School of Mines, &c. 18mo. 1s.

**Kempe.**—HOW TO DRAW A STRAIGHT LINE; a Lecture on Linkages. By A. B. KEMPE. With Illustrations. Crown 8vo. 1s. 6d. (*Nature Series*.)

**Kennedy.**—MECHANICS OF MACHINERY. By A. B. W. KENNEDY, M.Inst.C.E., Professor of Engineering and Mechanical Technology in University College, London. With Illustrations. Crown 8vo. [*In the press*.]

**Lang.**—EXPERIMENTAL PHYSICS. By P. R. SCOTT LANG, M.A., Professor of Mathematics in the University of St. Andrew. Crown 8vo. [*In preparation*.]

**Martineau (Miss C. A.).**—EASY LESSONS ON HEAT. By Miss C. A. MARTINEAU. Illustrated. Extra fcap. 8vo. 2s. 6d.

**Mayer.**—SOUND: a Series of Simple, Entertaining, and Inexpensive Experiments in the Phenomena of Sound, for the Use of Students of every age. By A. M. MAYER, Professor of Physics in the Stevens Institute of Technology, &c. With numerous Illustrations. Crown 8vo. 2s. 6d. (*Nature Series*.)

**Mayer and Barnard.**—LIGHT: a Series of Simple, Entertaining, and Inexpensive Experiments in the Phenomena of Light, for the Use of Students of every age. By A. M. MAYER and C. BARNARD. With numerous Illustrations. Crown 8vo. 2s. 6d. (*Nature Series*.)

**Newton.**—PRINCIPIA. Edited by Professor Sir W. THOMSON and Professor BLACKBURNE. 4to, cloth. 31s. 6d.

THE FIRST THREE SECTIONS OF NEWTON'S PRINCIPIA. With Notes and Illustrations. Also a Collection of Problems, principally intended as Examples of Newton's Methods. By PERCIVAL FROST, M.A. Third Edition. 8vo. 12s.

**Parkinson.**—A TREATISE ON OPTICS. By S. PARKINSON, D.D., F.R.S., Tutor and Prælector of St. John's College, Cambridge. New Edition, revised and enlarged. Crown 8vo. 10s. 6d.

**Perry.**—STEAM. AN ELEMENTARY TREATISE. By JOHN PERRY, C.E., Whitworth Scholar, Fellow of the Chemical Society, Lecturer in Physics at Clifton College. With numerous Woodcuts and Numerical Examples and Exercises. 18mo. 4s. 6d.

**Rayleigh.**—THE THEORY OF SOUND. By LORD RAYLEIGH, M.A., F.R.S., formerly Fellow of Trinity College, Cambridge, 8vo. Vol. I. 12s. 6d. Vol. II. 12s. 6d. [Vol. III. *in the press.*]

**Reuleaux.**—THE KINEMATICS OF MACHINERY. *Outlines of a Theory of Machines.* By Professor F. REULEAUX. Translated and Edited by Professor A. B. W. KENNEDY, C.E. With 450 Illustrations. Medium 8vo. 21s.

**Shann.**—AN ELEMENTARY TREATISE ON HEAT, IN RELATION TO STEAM AND THE STEAM-ENGINE. By G. SHANN, M.A. With Illustrations. Crown 8vo. 4s. 6d.

**Spottiswoode.**—POLARISATION OF LIGHT. By the late W. SPOTTISWOODE, P.R.S. With many Illustrations. New Edition. Crown 8vo. 3s. 6d. (*Nature Series.*)

**Stewart (Balfour).**—Works by BALFOUR STEWART, F.R.S., Professor of Natural Philosophy in the Victoria University the Owens College, Manchester.

PRIMER OF PHYSICS. With numerous Illustrations. New Edition, with Questions. 18mo. 1s. (*Science Primers.*)

LESSONS IN ELEMENTARY PHYSICS. With numerous Illustrations and Chromolitho of the Spectra of the Sun, Stars, and Nebulæ. New Edition. Fcap. 8vo. 4s. 6d.

QUESTIONS ON BALFOUR STEWART'S ELEMENTARY LESSONS IN PHYSICS. By Prof. THOMAS H. CORE, Owens College, Manchester. Fcap. 8vo. 2s.

**Stone.**—AN ELEMENTARY TREATISE ON SOUND. By W. H. STONE, M.B. With Illustrations. 18mo. 3s. 6d.

**Tait.**—AN ELEMENTARY TREATISE ON HEAT. By Professor TAIT, F.R.S.E. Illustrated. Crown 8vo. [*In the press.*]

**Thompson.**—ELEMENTARY LESSONS IN ELECTRICITY AND MAGNETISM. By SILVANUS P. THOMPSON. Professor of Experimental Physics in University College, Bristol. With Illustrations. Fcap. 8vo. 4s. 6d.

**Todhunter.**—NATURAL PHILOSOPHY FOR BEGINNERS. By I. TODHUNTER, M.A., F.R.S., D.Sc.  
Part I. The Properties of Solid and Fluid Bodies. 18mo. 3s. 6d.  
Part II. Sound, Light, and Heat. 18mo. 3s. 6d.

**Wright (Lewis).**—LIGHT; A COURSE OF EXPERIMENTAL OPTICS, CHIEFLY WITH THE LANTERN. By LEWIS WRIGHT. With nearly 200 Engravings and Coloured Plates. Crown 8vo. 7s. 6d.

### ASTRONOMY.

**Airy.**—POPULAR ASTRONOMY. With Illustrations by Sir G. B. AIRY, K.C.B., formerly Astronomer-Royal. New Edition. 18mo. 4s. 6d.

**Forbes.**—TRANSIT OF VENUS. By G. FORBES, M.A., Professor of Natural Philosophy in the Andersonian University, Glasgow. Illustrated. Crown 8vo. 3s. 6d. (*Nature Series.*)

**Godfray.**—Works by HUGH GODFRAY, M.A., Mathematical Lecturer at Pembroke College, Cambridge.

A TREATISE ON ASTRONOMY, for the Use of Colleges and Schools. New Edition. 8vo. 12s. 6d.

AN ELEMENTARY TREATISE ON THE LUNAR THEORY, with a Brief Sketch of the Problem up to the time of Newton. Second Edition, revised. Crown 8vo. 5s. 6d.

**Lockyer.**—Works by J. NORMAN LOCKYER, F.R.S.

PRIMER OF ASTRONOMY. With numerous Illustrations. 18mo. 1s. (*Science Primers.*)

ELEMENTARY LESSONS IN ASTRONOMY. With Coloured Diagram of the Spectra of the Sun, Stars, and Nebulæ, and numerous Illustrations. New Edition. Fcap. 8vo. 5s. 6d.

QUESTIONS ON LOCKYER'S ELEMENTARY LESSONS IN ASTRONOMY. For the Use of Schools. By JOHN FORBES-ROBERTSON. 18mo, cloth limp. 1s. 6d.

THE SPECTROSCOPE AND ITS APPLICATIONS. With Coloured Plate and numerous Illustrations. New Edition. Crown 8vo. 3s. 6d.

**Newcomb.**—POPULAR ASTRONOMY. By S. NEWCOMB, LL.D., Professor U.S. Naval Observatory. With 112 Illustrations and 5 Maps of the Stars. Second Edition, revised. 8vo. 18s.

"It is unlike anything else of its kind, and will be of more use in circulating a knowledge of Astronomy than nine-tenths of the books which have appeared on the subject of late years."—SATURDAY REVIEW.

## CHEMISTRY.

**Fleischer.**—A SYSTEM OF VOLUMETRIC ANALYSIS. Translated, with Notes and Additions, from the Second German Edition, by M. M. PATTISON MUIR, F.R.S.E. With Illustrations. Crown 8vo. 7s. 6d.

**Jones.**—Works by FRANCIS JONES, F.R.S.E., F.C.S., Chemical Master in the Grammar School, Manchester.

THE OWENS COLLEGE JUNIOR COURSE OF PRACTICAL CHEMISTRY. With Preface by Professor ROSCOE, and Illustrations. New Edition. 18mo. 2s. 6d.

QUESTIONS ON CHEMISTRY. A Series of Problems and Exercises in Inorganic and Organic Chemistry. Fcap. 8vo. 3s.

**Landauer.**—BLOWPIPE ANALYSIS. By J. LANDAUER. Authorised English Edition by J. TAYLOR and W. E. KAY, of Owens College, Manchester. Extra fcap. 8vo. 4s. 6d.

**Lupton.**—ELEMENTARY CHEMICAL ARITHMETIC. With 1,100 Problems. By SYDNEY LUPTON, M.A., Assistant-Master at Harrow. Extra fcap. 8vo. 5s.

**Muir.**—PRACTICAL CHEMISTRY FOR MEDICAL STUDENTS. Specially arranged for the first M.B. Course. By M. M. PATTISON MUIR, F.R.S.E. Fcap. 8vo. 1s. 6d.

**Roscoe.**—Works by H. E. ROSCOE, F.R.S. Professor of Chemistry in the Victoria University the Owens College, Manchester.

PRIMER OF CHEMISTRY. With numerous Illustrations. New Edition. With Questions. 18mo. 1s. (*Science Primers*).

LESSONS IN ELEMENTARY CHEMISTRY, INORGANIC AND ORGANIC. With numerous Illustrations and Chromolitho of the Solar Spectrum, and of the Alkalies and Alkaline Earths. New Edition. Fcap. 8vo. 4s. 6d.

A SERIES OF CHEMICAL PROBLEMS, prepared with Special Reference to the foregoing, by T. E. THORPE, Ph.D., Professor of Chemistry in the Yorkshire College of Science, Leeds, Adapted for the Preparation of Students for the Government, Science, and Society of Arts Examinations. With a Preface by Professor ROSCOE, F.R.S. New Edition, with Key. 18mo. 2s.

**Roscoe and Schorlemmer.**—INORGANIC AND ORGANIC CHEMISTRY. A Complete Treatise on Inorganic and Organic Chemistry. By Professor H. E. ROSCOE, F.R.S., and Professor C. SCHORLEMMER, F.R.S. With numerous Illustrations. Medium 8vo.

Vols. I. and II.—INORGANIC CHEMISTRY.

Vol. I.—The Non-Metallic Elements. 21s. Vol. II. Part I.—Metals. 18s. Vol. II. Part II.—Metals. 18s.

Vol. III.—ORGANIC CHEMISTRY. Part I.—THE CHEMISTRY OF THE HYDROCARBONS and their Derivatives, or ORGANIC CHEMISTRY. With numerous Illustrations. Medium 8vo. 21s. [Part II. in the press.]

**Schorlemmer.**—A MANUAL OF THE CHEMISTRY OF THE CARBON COMPOUNDS, OR ORGANIC CHEMISTRY. By C. SCHORLEMMER, F.R.S., Professor of Chemistry in the Victoria University the Owens College, Manchester. With Illustrations. 8vo. 14s.

**Thorpe.**—A SERIES OF CHEMICAL PROBLEMS, prepared with Special Reference to Professor Roscoe's Lessons in Elementary Chemistry, by T. E. THORPE, Ph.D., Professor of Chemistry in the Yorkshire College of Science, Leeds, adapted for the Preparation of Students for the Government, Science, and Society of Arts Examinations. With a Preface by Professor ROSCOE. New Edition, with Key. 18mo. 2s.

**Thorpe and Rücker.**—A TREATISE ON CHEMICAL PHYSICS. By Professor THORPE, F.R.S., and Professor RÜCKER, of the Yorkshire College of Science. Illustrated. 8vo. [In preparation.]

**Wright.**—METALS AND THEIR CHIEF INDUSTRIAL APPLICATIONS. BY C. ALDER WRIGHT, D.Sc., &c., Lecturer on Chemistry in St. Mary's Hospital Medical School. Extra f.ap. 8vo. 3s. 6d.

## BIOLOGY.

**Allen.**—ON THE COLOUR OF FLOWERS, as Illustrated in the British Flora. By GRANT ALLEN. With Illustrations. Crown 8vo. 3s. 6d. (*Nature Series.*)

**Balfour.**—A TREATISE ON COMPARATIVE EMBRYOLOGY. By F. M. BALFOUR, M.A., F.R.S., Fellow and Lecturer of Trinity College, Cambridge. With Illustrations. In 2 vols. 8vo. Vol. I. 18s. Vol. II. 21s.

**Bettany.**—FIRST LESSONS IN PRACTICAL BOTANY. By G. T. BETTANY, M.A., F.L.S., Lecturer in Botany at Guy's Hospital Medical School. 18mo. 1s.

**Darwin (Charles).**—MEMORIAL NOTICES OF CHARLES DARWIN, F.R.S., &c. By Professor HUXLEY, P.R.S., G. J. ROMANES, F.R.S., ARCHIBALD GEIKIE, F.R.S., and W. T. THISELTON DYER, F.R.S. Reprinted from *Nature*. With a Portrait, engraved by C. H. JEENS. Crown 8vo. 2s. 6d. (*Nature Series*.)

**Dyer and Vines.**—THE STRUCTURE OF PLANTS. By Professor THISELTON DYER, F.R.S., assisted by SYDNEY VINES, D.Sc., Fellow and Lecturer of Christ's College, Cambridge, and F. O. BOWER, M.A., Lecturer in the Normal School of Science. With numerous Illustrations. [*In preparation.*]

**Flower (W. H.)**—AN INTRODUCTION TO THE OSTEOLOGY OF THE MAMMALIA. Being the substance of the Course of Lectures delivered at the Royal College of Surgeons of England in 1870. By Professor W. H. FLOWER, F.R.S., F.R.C.S. With numerous Illustrations. New Edition, enlarged. Crown 8vo. 10s. 6d.

**Foster.**—Works by MICHAEL FOSTER, M.D., F.R.S., Professor of Physiology in the University of Cambridge.

PRIMER OF PHYSIOLOGY. With numerous Illustrations. New Edition. 18mo. 1s.

A TEXT-BOOK OF PHYSIOLOGY. With Illustrations. Fourth Edition, revised. 8vo. 21s.

**Foster and Balfour.**—THE ELEMENTS OF EMBRYOLOGY. By MICHAEL FOSTER, M.A., M.D., LL.D., F.R.S., Professor of Physiology in the University of Cambridge, Fellow of Trinity College, Cambridge, and the late FRANCIS M. BALFOUR, M.A., LL.D., F.R.S., Fellow of Trinity College, Cambridge, and Professor of Animal Morphology in the University. Second Edition, revised. Edited by ADAM SEDGWICK, M.A., Fellow and Assistant Lecturer of Trinity College, Cambridge, and WALTER HEAPE, Demonstrator in the Morphological Laboratory of the University of Cambridge. With Illustrations. Crown 8vo. 10s. 6d.

- Foster and Langley.**—A COURSE OF ELEMENTARY PRACTICAL PHYSIOLOGY. By Prof. MICHAEL FOSTER, M.D., F.R.S., &c., and J. N. LANGLEY, B.A. New Edition. Crown 8vo. 6s.
- Gamgee.**—A TEXT-BOOK OF THE PHYSIOLOGICAL CHEMISTRY OF THE ANIMAL BODY. Including an Account of the Chemical Changes occurring in Disease. By A. GAMGEE, M.D., F.R.S., Professor of Physiology in the Victoria University the Owens College, Manchester. 2 Vols. 8vo. With Illustrations. Vol. I. 18s. [Vol. II. in the press.]
- Gegenbaur.**—ELEMENTS OF COMPARATIVE ANATOMY. By Professor CARL GEGENBAUR. A Translation by F. JEFFREY BELL, B.A. Revised with Preface by Professor E. RAY LANKESTER, F.R.S. With numerous Illustrations. 8vo. 21s.
- Gray.**—STRUCTURAL BOTANY, OR ORGANOGRAPHY ON THE BASIS OF MORPHOLOGY. To which are added the principles of Taxonomy and Phytography, and a Glossary of Botanical Terms. By Professor ASA GRAY, LL.D. 8vo. 10s. 6d.
- Hooker.**—Works by Sir J. D. HOOKER, K.C.S.I., C.B., M.D., F.R.S., D.C.L.  
**PRIMER OF BOTANY.** With numerous Illustrations. New Edition. 18mo. 1s. (*Science Primers.*)  
**THE STUDENT'S FLORA OF THE BRITISH ISLANDS.** New Edition, revised. Globe 8vo. 10s. 6d.
- Huxley.**—Works by Professor HUXLEY, P.R.S.  
**INTRODUCTORY PRIMER OF SCIENCE.** 18mo. 1s. (*Science Primers.*)  
**LESSONS IN ELEMENTARY PHYSIOLOGY.** With numerous Illustrations. New Edition. Fcap. 8vo. 4s. 6d.  
**QUESTIONS ON HUXLEY'S PHYSIOLOGY FOR SCHOOLS.** By T. ALCOCK, M.D. 18mo. 1s. 6d.  
**PRIMER OF ZOOLOGY.** 18mo. (*Science Primers.*) [In preparation.]
- Huxley and Martin.**—A COURSE OF PRACTICAL INSTRUCTION IN ELEMENTARY BIOLOGY. By Professor HUXLEY, P.R.S., assisted by H. N. MARTIN, M.B., D.Sc. New Edition, revised. Crown 8vo. 6s.



- Lankester.**—Works by Professor E. RAY LANKESTER, F.R.S.  
 A TEXTBOOK OF ZOOLOGY. Crown 8vo. [*In preparation.*]  
 DEGENERATION: A CHAPTER IN DARWINISM. Illustrated. Crown 8vo. 2s. 6d. (*Nature Series.*)
- Lubbock.**—Works by SIR JOHN LUBBOCK, M.P., F.R.S., D.C.L.  
 THE ORIGIN AND METAMORPHOSES OF INSECTS. With numerous Illustrations. New Edition. Crown 8vo. 3s. 6d. (*Nature Series.*)  
 ON BRITISH WILD FLOWERS CONSIDERED IN RELATION TO INSECTS. With numerous Illustrations. New Edition. Crown 8vo. 4s. 6d. (*Nature Series.*)
- M'Kendrick.**—OUTLINES OF PHYSIOLOGY IN ITS RELATIONS TO MAN. By J. G. M'KENDRICK, M.D., F.R.S.E. With Illustrations. Crown 8vo. 12s. 6d.
- Martin and Moale.**—ON THE DISSECTION OF VERTEBRATE ANIMALS. By Professor H. N. MARTIN and W. A. MOALE. Crown 8vo. [*In preparation.*]  
 (See also page 41.)
- Miall.**—STUDIES IN COMPARATIVE ANATOMY.  
 No. I.—The Skull of the Crocodile: a Manual for Students. By L. C. MIALL, Professor of Biology in the Yorkshire College and Curator of the Leeds Museum. 8vo. 2s. 6d.  
 No. II.—Anatomy of the Indian Elephant. By L. C. MIALL and F. GREENWOOD. With Illustrations. 8vo. 5s.
- Mivart.**—Works by ST. GEORGE MIVART, F.R.S. Lecturer in Comparative Anatomy at St. Mary's Hospital.  
 LESSONS IN ELEMENTARY ANATOMY. With upwards of 400 Illustrations. Fcap. 8vo. 6s. 6d.  
 THE COMMON FROG. With numerous Illustrations. Crown 8vo. 3s. 6d. (*Nature Series.*)
- Müller.**—THE FERTILISATION OF FLOWERS. By Professor HERMANN MÜLLER. Translated and Edited by D'ARCY W. THOMPSON, B.A., Scholar of Trinity College, Cambridge. With a Preface by CHARLES DARWIN, F.R.S. With numerous Illustrations. Medium 8vo. 21s.
- Oliver.**—Works by DANIEL OLIVER, F.R.S., &c., Professor of Botany in University College, London, &c.  
 FIRST BOOK OF INDIAN BOTANY. With numerous Illustrations. Extra fcap. 8vo. 6s. 6d.  
 LESSONS IN ELEMENTARY BOTANY. With nearly 200 Illustrations. New Edition. Fcap. 8vo. 4s. 6d.

**Parker.**—A COURSE OF INSTRUCTION IN ZOOTOMY (VERTEBRATA). By T. JEFFREY PARKER, B.Sc. London, Professor of Biology in the University of Otago. With Illustrations. Crown 8vo. *[In the press,*

**Parker and Bettany.**—THE MORPHOLOGY OF THE SKULL. By Professor PARKER and G. T. BETTANY. Illustrated. Crown 8vo. 10s. 6d.

**Romanes.**—THE SCIENTIFIC EVIDENCES OF ORGANIC EVOLUTION. By G. J. ROMANES, M.A., LL.D., F.R.S., Zoological Secretary to the Linnean Society. Crown 8vo. 2s. 6d (*Nature Series.*)

**Smith.**—Works by JOHN SMITH, A.L.S., &c.

A DICTIONARY OF ECONOMIC PLANTS. Their History, Products, and Uses. 8vo. 14s.

DOMESTIC BOTANY: An Exposition of the Structure and Classification of Plants, and their Uses for Food, Clothing, Medicine, and Manufacturing Purposes. With Illustrations. New Issue. Crown 8vo. 12s. 6d.

## MEDICINE.

**Brunton.**—Works by T. LAUDER BRUNTON, M.D., Sc.D., F.R.C.P., F.R.S., Examiner in Materia Medica in the University of London, late Examiner in Materia Medica in the University of Edinburgh, and the Royal College of Physicians, London.

A TREATISE ON MATERIA MEDICA. 8vo. *[In the press.*

TABLES OF MATERIA MEDICA: A Companion to the Materia Medica Museum. With Illustrations. New Edition Enlarged. 8vo. 10s. 6d.

**Hamilton.**—A TEXT-BOOK OF PATHOLOGY. By D. J. HAMILTON, Professor of Pathological Anatomy (Sir Erasmus Wilson Chair), University of Aberdeen. 8vo. *[In preparation.*

**Ziegler-Macalister.**—TEXT-BOOK OF PATHOLOGICAL ANATOMY AND PATHOGENESIS. By Professor ERNST ZIEGLER of Tübingen. Translated and Edited for English Students by DONALD MACALISTER, M.A., M.B., B.Sc., Fellow of St. John's College, Cambridge. With numerous Illustrations. Medium 8vo. Part I.—GENERAL PATHOLOGICAL ANATOMY. 12s. 6d. *[PART II. in the press.*

## ANTHROPOLOGY.

**Flower.**—FASHION IN DEFORMITY, as Illustrated in the Customs of Barbarous and Civilised Races. By Professor FLOWER, F.R.S., F.R.C.S. With Illustrations. Crown 8vo 2s. 6d. (*Nature Series*).

**Tylor.**—ANTHROPOLOGY. An Introduction to the Study of Man and Civilisation. By E. B. TYLOR, D.C.L., F.R.S. With numerous Illustrations. Crown 8vo. 7s. 6d.

## PHYSICAL GEOGRAPHY & GEOLOGY.

**Blanford.**—THE RUDIMENTS OF PHYSICAL GEOGRAPHY FOR THE USE OF INDIAN SCHOOLS; with a Glossary of Technical Terms employed. By H. F. BLANFORD, F.R.S. New Edition, with Illustrations. Globe 8vo. 2s. 6d.

**Geikie.**—Works by ARCHIBALD GEIKIE, F.R.S., Director General of the Geological Surveys of the United Kingdom.

PRIMER OF PHYSICAL GEOGRAPHY. With numerous Illustrations. New Edition. With Questions. 18mo. 1s. (*Science Primers*.)

ELEMENTARY LESSONS IN PHYSICAL GEOGRAPHY. With numerous Illustrations. Fcap. 8vo. 4s. 6d.  
QUESTIONS ON THE SAME. 1s. 6d.

PRIMER OF GEOLOGY. With numerous Illustrations. New Edition. 18mo. 1s. (*Science Primers*.)

ELEMENTARY LESSONS IN GEOLOGY. With Illustrations. Fcap. 8vo. [*In preparation*].

TEXT-BOOK OF GEOLOGY. With numerous Illustrations. 8vo. 28s.

OUTLINES OF FIELD GEOLOGY. With Illustrations. New Edition. Extra fcap. 8vo. 3s. 6d.

**Huxley.**—PHYSIOGRAPHY. An Introduction to the Study of Nature. By Professor HUXLEY, P.R.S. With numerous Illustrations, and Coloured Plates. New and Cheaper Edition. Crown 8vo. 6s.

## AGRICULTURE.

**Frankland.**—AGRICULTURAL CHEMICAL ANALYSIS, A Handbook of. By PERCY FARADAY FRANKLAND, Ph.D., B.Sc., F.C.S., Associate of the Royal School of Mines, and Demonstrator of Practical and Agricultural Chemistry in the Normal School of Science and Royal School of Mines, South Kensington Museum. Founded upon *Leitfaden für die Agricultur-Chemische Analyse*, von Dr. F. KROCKER. Crown 8vo. 7s. 6d.

**Tanner.**—Works by HENRY TANNER, F.C.S., M.R.A.C., Examiner in the Principles of Agriculture under the Government Department of Science, sometime Professor of Agricultural Science, University College, Aberystwith.

ELEMENTARY LESSONS IN THE SCIENCE OF AGRICULTURAL PRACTICE. Fcap. 8vo. 3s. 6d.

FIRST PRINCIPLES OF AGRICULTURE. 18mo. 1s.

THE PRINCIPLES OF AGRICULTURE. A Series of Reading Books for use in Elementary Schools. Prepared by HENRY TANNER, F.C.S., M.R.A.C. Extra fcap. 8vo.

I. The Alphabet of the Principles of Agriculture. 6d.

II. Further Steps in the Principles of Agriculture. 1s.

III. Elementary School Readings in the Principles of Agriculture for the third stage. 1s.

## POLITICAL ECONOMY.

**Cossa.**—GUIDE TO THE STUDY OF POLITICAL ECONOMY. By Dr. LUIGI COSSA, Professor in the University of Pavia. Translated from the Second Italian Edition. With a Preface by W. STANLEY JEVONS, F.R.S. Crown 8vo. 4s. 6d.

**Fawcett.**—POLITICAL ECONOMY FOR BEGINNERS. By MILLICENT G. FAWCETT. New Edition. 18mo. 2s. 6d.

**Fawcett.**—A MANUAL OF POLITICAL ECONOMY. By Right Hon. HENRY FAWCETT, M.P., F.R.S. New Edition, revised and enlarged. Crown 8vo. 12s.

**Jevons.**—PRIMER OF POLITICAL ECONOMY. By W. STANLEY JEVONS, LL.D., M.A., F.R.S. New Edition. 18mo. 1s. (*Science Primers.*)

**Marshall.**—THE ECONOMICS OF INDUSTRY. By A. MARSHALL, M.A., late Principal of University College, Bristol, and MARY P. MARSHALL, late Lecturer at Newnham Hall, Cambridge. Extra fcap. 8vo. 2s. 6d.

**Sidgwick.**—THE PRINCIPLES OF POLITICAL ECONOMY. By HENRY SIDGWICK, M.A., Prælector in Moral and Political Philosophy in Trinity College, Cambridge, Author of "The Methods of Ethics." 8vo. 16s.

**Walker.**—POLITICAL ECONOMY. By FRANCIS A. WALKER, M.A., Ph.D., Author of "The Wages Question," "Money," "Money in its Relation to Trade," &c. 8vo. 10s. 6d.

## MENTAL & MORAL PHILOSOPHY.

**Caird.**—MORAL PHILOSOPHY, An Elementary Treatise on. By Prof. E. CAIRD, of Glasgow University. Fcap. 8vo. *[In preparation.]*

**Calderwood.**—HANDBOOK OF MORAL PHILOSOPHY. By the Rev. HENRY CALDERWOOD, LL.D., Professor of Moral Philosophy, University of Edinburgh. New Edition. Crown 8vo. 6s.

**Clifford.**—SEEING AND THINKING. By the late Professor W. K. CLIFFORD, F.R.S. With Diagrams. Crown 8vo. 3s. 6d. (*Nature Series.*)

**Jevons.**—Works by the late W. STANLEY JEVONS, LL.D., M.A., F.R.S.

PRIMER OF LOGIC. New Edition. 18mo. 1s. (*Science Primers.*)

ELEMENTARY LESSONS IN LOGIC; Deductive and Inductive, with copious Questions and Examples, and a Vocabulary of Logical Terms. New Edition. Fcap. 8vo. 3s. 6d.

THE PRINCIPLES OF SCIENCE. A Treatise on Logic and Scientific Method. New and Revised Edition. Crown 8vo. 12s. 6d.

STUDIES IN DEDUCTIVE LOGIC. Crown 8vo. 6s.

**Robertson.**—ELEMENTARY LESSONS IN PSYCHOLOGY.  
By G. CROOM ROBERTSON, Professor of Mental Philosophy, &c.,  
University College, London. *[In preparation.]*

**Sidgwick.**—THE METHODS OF ETHICS. By HENRY  
SIDGWICK, M.A., Prælector in Moral and Political Philosophy  
in Trinity College, Cambridge. Second Edition. 8vo. 14s.

---

## HISTORY AND GEOGRAPHY.

**Arnold.**—THE ROMAN SYSTEM OF PROVINCIAL ADMINISTRATION TO THE ACCESSION OF CONSTANTINE THE GREAT. By W. T. ARNOLD, B.A. Crown 8vo. 6s.

"Ought to prove a valuable handbook to the student of Roman history."—*GUARDIAN.*

**Beesly.**—STORIES FROM THE HISTORY OF ROME.  
By Mrs. BEESLY. Fcap. 8vo. 2s. 6d.

"The attempt appears to us in every way successful. The stories are interesting in themselves, and are told with perfect simplicity and good feeling."—*DAILY NEWS.*

**Brook.**—FRENCH HISTORY FOR ENGLISH CHILDREN.  
By SARAH BROOK. With Coloured Maps. Crown 8vo. 6s.

**Clarke.**—CLASS-BOOK OF GEOGRAPHY. By C. B. CLARKE, M.A., F.L.S., F.G.S., F.R.S. New Edition, with Eighteen Coloured Maps. Fcap. 8vo. 3s.

**Freeman.**—OLD-ENGLISH HISTORY. By EDWARD A. FREEMAN, D.C.L., LL.D., late Fellow of Trinity College, Oxford. With Five Coloured Maps. New Edition. Extra fcap. 8vo. 6s.

**Fyffe.**—A SCHOOL HISTORY OF GREECE. By C. A. FYFFE, M.A., Fellow of University College, Oxford. Crown 8vo. *[In preparation.]*

**Green.**—Works by JOHN RICHARD GREEN, M.A., LL.D.,  
Honorary Fellow of Jesus College, Oxford.

**SHORT HISTORY OF THE ENGLISH PEOPLE.** With  
Coloured Maps, Genealogical Tables, and Chronological Annals.  
Crown 8vo. 8s. 6d. Ninety-second Thousand.

"Stands alone as the one general history of the country, for the sake of which  
all others, if young and old are wise, will be speedily and surely set aside."—  
ACADEMY.

**ANALYSIS OF ENGLISH HISTORY**, based on Green's "Short  
History of the English People." By C. W. A. TAIT, M.A.,  
Assistant-Master, Clifton College. Crown 8vo. 3s. 6d.

**READINGS FROM ENGLISH HISTORY.** Selected and  
Edited by JOHN RICHARD GREEN. Three Parts. Globe 8vo.  
1s. 6d. each. I. Hengist to Cressy. II. Cressy to Cromwell.  
III. Cromwell to Balaklava.

**A SHORT GEOGRAPHY OF THE BRITISH ISLANDS.** By  
JOHN RICHARD GREEN and ALICE STOPFORD GREEN. With  
Maps. Fcap. 8vo. 3s. 6d.

**Grove.**—A PRIMER OF GEOGRAPHY. By Sir GEORGE  
GROVE, D.C.L., F.R.G.S. With Illustrations. 18mo. 1s.  
(*Science Primers.*)

**Guest.**—LECTURES ON THE HISTORY OF ENGLAND.  
By M. J. GUEST. With Maps. Crown 8vo. 6s.

"It is not too much to assert that this is one of the very best class books of English  
History for young students ever published."—SCOTSMAN.

**Historical Course for Schools**—Edited by EDWARD A.  
FREEMAN, D.C.L., late Fellow of Trinity College, Oxford.

I.—GENERAL SKETCH OF EUROPEAN HISTORY. By  
EDWARD A. FREEMAN, D.C.L. New Edition, revised and  
enlarged, with Chronological Table, Maps, and Index. 18mo.  
3s. 6d.

II.—HISTORY OF ENGLAND. By EDITH THOMPSON. New  
Edition, revised and enlarged, with Coloured Maps. 18mo.  
2s. 6d.

III.—HISTORY OF SCOTLAND. By MARGARET MACARTHUR.  
New Edition. 18mo. 2s.

IV.—HISTORY OF ITALY. By the Rev. W. HUNT, M.A.  
New Edition, with Coloured Maps. 18mo. 3s. 6d.

**Historical Course for Schools.**—(continued)—

V.—HISTORY OF GERMANY. By J. SIME, M.A. 18mo. 3s.

VI.—HISTORY OF AMERICA. By JOHN A. DOYLE. With Maps. 18mo. 4s. 6d.

EUROPEAN COLONIES. By E. J. PAYNE, M.A. With Maps. 18mo. 4s. 6d.

FRANCE. By CHARLOTTE M. YONGE. With Maps. 18mo. 3s. 6d.

GREECE. By EDWARD A. FREEMAN, D.C.L. [*In preparation.*]

ROME. By EDWARD A. FREEMAN, D.C.L. [*In the press.*]

**History Primers**—Edited by JOHN RICHARD GREEN, M.A., LL.D., Author of "A Short History of the English People."

ROME. By the Rev. M. CREIGHTON, M.A., late Fellow and Tutor of Merton College, Oxford. With Eleven Maps. 18mo. 1s.

"The author has been curiously successful in telling in an intelligent way the story of Rome from first to last."—SCHOOL BOARD CHRONICLE.

GREECE. By C. A. FYFFE, M.A., Fellow and late Tutor of University College, Oxford. With Five Maps. 18mo. 1s.

"We give our unqualified praise to this little manual."—SCHOOLMASTER.

EUROPEAN HISTORY. By E. A. FREEMAN, D.C.L., LL.D. With Maps. 18mo. 1s.

"The work is always clear, and forms a luminous key to European history."—SCHOOL BOARD CHRONICLE.

GREEK ANTIQUITIES. By the Rev. J. P. MAHAFFY, M.A. Illustrated. 18mo. 1s.

"All that is necessary for the scholar to know is told so compactly yet so fully, and in a style so interesting, that it is impossible for even the dullest boy to look on this little work in the same light as he regards his other school books."—SCHOOLMASTER.

CLASSICAL GEOGRAPHY. By H. F. TOZER, M.A. 18mo. 1s.

"Another valuable aid to the study of the ancient world. . . . It contains an enormous quantity of information packed into a small space, and at the same time communicated in a very readable shape."—JOHN BULL.

GEOGRAPHY. By GEORGE GROVE, D.C.L. With Maps. 18mo. 1s.

"A model of what such a work should be. . . . We know of no short treatise better suited to infuse life and spirit into the dull lists of proper names of which our ordinary class-books so often almost exclusively consist."—TIMES.



**History Primers** *Continued.*

**ROMAN ANTIQUITIES.** By Professor WALKINS. Illustrated. 18mo. 1s.

"A little book that throws a blaze of light on Roman history, and is, moreover, intensely interesting."—SCHOOL BOARD CHRONICLE.

**FRANCE.** By CHARLOTTE M. YONGE. 18mo. 1s.

"May be considered a wonderfully successful piece of work. . . . Its general merit as a vigorous and clear sketch, giving in a small space a vivid idea of the history of France, remains undeniable."—SATURDAY REVIEW.

**Hole.**—A GENEALOGICAL STEMMA OF THE KINGS OF ENGLAND AND FRANCE. By the Rev. C. HOLE. On Sheet. 1s.

**Kiepert.**—A MANUAL OF ANCIENT GEOGRAPHY. From the German of Dr. H. KIEPERT. Crown 8vo. 5s.

**Lethbridge.**—A SHORT MANUAL OF THE HISTORY OF INDIA. With an Account of INDIA AS IT IS. The Soil, Climate, and Productions; the People, their Races, Religions, Public Works, and Industries; the Civil Services, and System of Administration. By ROPER LETHBRIDGE, M.A., C.I.E., late Scholar of Exeter College, Oxford, formerly Principal of Kishnaghur College, Bengal, Fellow and sometime Examiner of the Calcutta University. With Maps. Crown 8vo. 5s.

**Michelet.**—A SUMMARY OF MODERN HISTORY. Translated from the French of M. MICHELET, and continued to the Present Time, by M. C. M. SIMPSON. Globe 8vo. 4s. 6d.

**Otté.**—SCANDINAVIAN HISTORY. By E. C. OTTÉ. With Maps. Globe 8vo. 6s.

**Ramsay.**—A SCHOOL HISTORY OF ROME. By G. G. RAMSAY, M.A., Professor of Humanity in the University of Glasgow. With Maps. Crown 8vo. [*In preparation.*]

**Tait.**—ANALYSIS OF ENGLISH HISTORY, based on Green's "Short History of the English People." By C. W. A. TAIT, M.A., Assistant-Master, Clifton College. Crown 8vo. 3s. 6d.

**Wheeler.**—A SHORT HISTORY OF INDIA AND OF THE FRONTIER STATES OF AFGHANISTAN, NEPAUL, AND BURMA. By J. TALBOYS WHEELER. With Maps. Crown 8vo. 12s.

"It is the best book of the kind we have ever seen, and we recommend it to a place in every school library."—EDUCATIONAL TIMES.

- Yonge (Charlotte M.).**—A PARALLEL HISTORY OF FRANCE AND ENGLAND: consisting of Outlines and Dates. By CHARLOTTE M. YONGE, Author of "The Heir of Redclyffe," &c., &c. Oblong 4to. 3s. 6d.
- CAMEOS FROM ENGLISH HISTORY.—FROM ROLLO TO EDWARD II.** By the Author of "The Heir of Redclyffe." Extra fcap. 8vo. New Edition. 5s.
- A SECOND SERIES OF CAMEOS FROM ENGLISH HISTORY.—THE WARS IN FRANCE.** New Edition. Extra fcap. 8vo. 5s.
- A THIRD SERIES OF CAMEOS FROM ENGLISH HISTORY.—THE WARS OF THE ROSES.** New Edition. Extra fcap. 8vo. 5s.
- CAMEOS FROM ENGLISH HISTORY—A FOURTH SERIES. REFORMATION TIMES.** Extra fcap. 8vo. 5s.
- CAMEOS FROM ENGLISH HISTORY.—A FIFTH SERIES. ENGLAND AND SPAIN.** Extra fcap. 8vo. 5s.
- EUROPEAN HISTORY.** Narrated in a Series of Historical Selections from the Best Authorities. Edited and arranged by E. M. SEWELL and C. M. YONGE. First Series, 1003—1154. New Edition. Crown 8vo. 6s. Second Series, 1088—1228. New Edition. Crown 8vo. 6s.

## MODERN LANGUAGES AND LITERATURE.

(1) English, (2) French, (3) German, (4) Modern Greek, (5) Italian.

### ENGLISH.

**Abbott.**—A SHAKESPEARIAN GRAMMAR. An attempt to illustrate some of the Differences between Elizabethan and Modern English. By the Rev. E. A. ABBOTT, D.D., Head Master of the City of London School. New Edition. Extra fcap. 8vo. 6s.

**Brooke.**—PRIMER OF ENGLISH LITERATURE. By the Rev. STOPFORD A. BROOKE, M.A. 18mo. 1s. (*Literature Primers.*)

**Butler.**—HUDIBRAS. Part I. Edited, with Introduction and Notes, by ALFRED MILNES, M.A. Crown 8vo. 3s. 6d.  
[Part II. *in the press.*]

**Cowper's TASK:** AN EPISTLE TO JOSEPH HILL, ESQ.; TIROCINIUM, or a Review of the Schools; and THE HISTORY OF JOHN GILPIN. Edited, with Notes, by WILLIAM BENHAM, B.D. Globe 8vo. 1s. (*Globe Readings from Standard Authors.*)

**Dowden.**—SHAKESPEARE. By Professor DOWDEN. 18mo. 1s. (*Literature Primers.*)

**Dryden.**—SELECT PROSE WORKS. Edited, with Introduction and Notes, by Professor C. D. YONGE. Fcap. 8vo. 2s. 6d.

**Gladstone.**—SPELLING REFORM FROM AN EDUCATIONAL POINT OF VIEW. By J. H. GLADSTONE, Ph.D., F.R.S., Member of the School Board for London. New Edition. Crown 8vo. 1s. 6d.

**Globe Readers.** For Standards I.—VI. Edited by A. F. MURISON. Sometime English Master at the Aberdeen Grammar School. With Illustrations. Globe 8vo.

Primer I. (48 pp.)	3d.	Book III. (232 pp.)	1s. 3d.
Primer II. (48 pp.)	3d.	Book IV. (328 pp.)	1s. 9d.
Book I. (96 pp.)	6d.	Book V. (416 pp.)	2s.
Book II. (136 pp.)	9d.	Book VI. (448 pp.)	2s. 6d.

"Among the numerous sets of readers before the public the present series is honourably distinguished by the marked superiority of its materials and the careful ability with which they have been adapted to the growing capacity of the pupils. The plan of the two primers is excellent for facilitating the child's first attempts to read. In the first three following books there is abundance of entertaining reading. . . . Better food for young minds could hardly be found."—THE ATHENÆUM.

#### GLOBE READINGS FROM STANDARD AUTHORS.

**Cowper's TASK:** AN EPISTLE TO JOSEPH HILL, ESQ.; TIROCINIUM, or a Review of the Schools; and THE HISTORY OF JOHN GILPIN. Edited, with Notes, by WILLIAM BENHAM, B.D. Globe 8vo. 1s.

**GLOBE READINGS FROM STANDARD AUTHORS***Continued—*

**Goldsmith's** VICAR OF WAKEFIELD. With a Memoir of Goldsmith by Professor MASSON. Globe 8vo. 1s.

**Lamb's (Charles)** TALES FROM SHAKESPEARE. Edited, with Preface, by ALFRED AINGER, M.A. Globe 8vo. 2s.

**Scott's (Sir Walter)** LAY OF THE LAST MINSTREL; and THE LADY OF THE LAKE. Edited, with Introductions and Notes, by FRANCIS TURNER PALGRAVE. Globe 8vo. 1s.

MARMION; and the LORD OF THE ISLES. By the same Editor. Globe 8vo. 1s.

**The Children's Garland from the Best Poets.—**

Selected and arranged by COVENTRY PATMORE. Globe 8vo. 2s.

**Yonge (Charlotte M.).—**A BOOK OF GOLDEN DEEDS OF ALL TIMES AND ALL COUNTRIES. Gathered and narrated anew by CHARLOTTE M. YONGE, the Author of "The Heir of Redclyffe." Globe 8vo. 2s.

**Goldsmith.**—THE TRAVELLER, or a Prospect of Society; and THE DESERTED VILLAGE. By OLIVER GOLDSMITH. With Notes, Philological and Explanatory, by J. W. HALES, M.A. Crown 8vo. 6d.

THE VICAR OF WAKEFIELD. With a Memoir of Goldsmith by Professor MASSON. Globe 8vo. 1s. (*Globe Readings from Standard Authors.*)

SELECT ESSAYS. Edited, with Introduction and Notes, by Professor C. D. YONGE. Fcap. 8vo. 2s. 6d.

**Hales.**—LONGER ENGLISH POEMS, with Notes, Philological and Explanatory, and an Introduction on the Teaching of English. Chiefly for Use in Schools. Edited by J. W. HALES, M.A., Professor of English Literature at King's College, London. New Edition. Extra fcap. 8vo. 4s. 6d.

**Johnson's** LIVES OF THE POETS. The Six Chief Lives (Milton, Dryden, Swift, Addison, Pope, Gray), with Macaulay's "Life of Johnson." Edited with Preface by MATTHEW ARNOLD. Crown 8vo. 6s.

**Lamb (Charles).**—TALES FROM SHAKESPEARE. Edited, with Preface, by ALFRED AINGER, M.A. (*Globe Readings from Standard Authors.*)

**Literature Primers**—Edited by JOHN RICHARD GREEN, M.A., LL.D., Author of "A Short History of the English People."

ENGLISH COMPOSITION. By Professor NICHOL. 18mo. 1s.

ENGLISH GRAMMAR. By the Rev. R. MORRIS, LL.D., sometime President of the Philological Society. 18mo, cloth. 1s.

ENGLISH GRAMMAR EXERCISES. By R. MORRIS, LL.D., and H. C. BOWEN, M.A. 18mo. 1s.

EXERCISES ON MORRIS'S PRIMER OF ENGLISH GRAMMAR. By JOHN WETHERELL, of the Middle School, Liverpool College. 18mo. 1s.

ENGLISH LITERATURE. By STOFFORD BROOKE, M.A. New Edition. 18mo. 1s.

SHAKSPERE. By Professor DOWDEN. 18mo. 1s.

THE CHILDREN'S TREASURY OF LYRICAL POETRY. Selected and arranged with Notes by FRANCIS TURNER PALGRAVE. In Two Parts. 18mo. 1s. each.

PHILOLOGY. By J. PEILE, M.A. 18mo. 1s.

*In preparation:—*

HISTORY OF THE ENGLISH LANGUAGE. By J. A. H. MURRAY, LL.D.

SPECIMENS OF THE ENGLISH LANGUAGE. To Illustrate the above. By the same Author.

**Macmillan's Reading Books.**—Adapted to the English and Scotch Codes. Bound in Cloth.

PRIMER. 18mo. (48 pp.) 2d.

BOOK	I. for Standard	I.	18mo.	(96 pp.)	4 <i>d.</i>
"	II.	"	II.	18mo.	(144 pp.) 5 <i>d.</i>
"	III.	"	III.	18mo.	(160 pp.) 6 <i>d.</i>
"	IV.	"	IV.	18mo.	(176 pp.) 8 <i>d.</i>
"	V.	"	V.	18mo.	(380 pp.) 1 <i>s.</i>

**Macmillan's Reading-Books** *Continued*—

BOOK VI. for Standard VI. Crown 8vo. (430 pp.) 2s.

Book VI. is fitted for higher Classes, and as an Introduction to English Literature.

"They are far above any others that have appeared both in form and substance. . . . The editor of the present series has rightly seen that reading books must aim chiefly at giving to the pupils the power of accurate, and, if possible, apt and skilful expression; at cultivating in them a good literary taste, and at arousing a desire of further reading.' This is done by taking care to select the extracts from true English classics, going up in Standard VI. course to Chaucer, Hooker, and Bacon, as well as Wordsworth, Macaulay, and Froude. . . . This is quite on the right track, and indicates justly the ideal which we ought to set before us."—*GUARDIAN*.

**Macmillan's Copy-Books**—

Published in two sizes, viz. :—

1. Large Post 4to. Price 4d. each.
2. Post Oblong. Price 2d. each.

1. INITIATORY EXERCISES AND SHORT LETTERS.
- \*2. WORDS CONSISTING OF SHORT LETTERS.
- \*3. LONG LETTERS. With words containing Long Letters—Figures.
- \*4. WORDS CONTAINING LONG LETTERS.
- 4a. PRACTISING AND REVISING COPY-BOOK. For Nos. 1 to 4.
- \*5. CAPITALS AND SHORT HALF-TEXT. Words beginning with a Capital.
- \*6. HALF-TEXT WORDS beginning with Capitals—Figures.
- \*7. SMALL-HAND AND HALF-TEXT. With Capitals and Figures.
- \*8. SMALL-HAND AND HALF-TEXT. With Capitals and Figures.
- 8a. PRACTISING AND REVISING COPY-BOOK. For Nos. 5 to 8.
- \*9. SMALL-HAND SINGLE HEADLINES—Figures.
10. SMALL-HAND SINGLE HEADLINES—Figures.
11. SMALL-HAND DOUBLE HEADLINES—Figures.
12. COMMERCIAL AND ARITHMETICAL EXAMPLES, &c.
- 12a. PRACTISING AND REVISING COPY-BOOK. For Nos. 8 to 12.

\* These numbers may be had with *Goodman's Patent Sliding Copies*. Large Post 4to. Price 6d. each.

**Martin.**—THE POET'S HOUR: Poetry selected and arranged for Children. By FRANCES MARTIN. New Edition. 18mo. 2s. 6d.

SPRING-TIME WITH THE POETS: Poetry selected by FRANCES MARTIN. New Edition. 18mo. 3s. 6d.

**Milton.**—By STOPFORD BROOKE, M.A. Fcap. 8vo. 1s. 6d. (*Classical Writers Series.*)

**Morris.**—Works by the Rev. R. MORRIS, LL.D. HISTORICAL OUTLINES OF ENGLISH ACCIDENCE, comprising Chapters on the History and Development of the Language, and on Word-formation. New Edition. Extra fcap. 8vo. 6s.

ELEMENTARY LESSONS IN HISTORICAL ENGLISH GRAMMAR, containing Accidence and Word-formation. New Edition. 18mo. 2s. 6d.

PRIMER OF ENGLISH GRAMMAR. 18mo. 1s. (See also *Literature Primers.*)

**Oliphant.**—THE OLD AND MIDDLE ENGLISH. A New Edition of "THE SOURCES OF STANDARD ENGLISH," revised and greatly enlarged. By T. L. KINGTON OLIPHANT. Extra fcap. 8vo. 9s.

**Palgrave.**—THE CHILDREN'S TREASURY OF LYRICAL POETRY. Selected and arranged, with Notes, by FRANCIS TURNER PALGRAVE. 18mo. 2s. 6d. Also in Two Parts. 18mo. 1s. each.

**Patmore.**—THE CHILDREN'S GARLAND FROM THE BEST POET'S. Selected and arranged by COVENTRY PATMORE. Globe 8vo. 2s. (*Globe Readings from Standard Authors.*)

**Plutarch.**—Being a Selection from the Lives which Illustrate Shakespeare. North's Translation. Edited, with Introductions, Notes, Index of Names, and Glossarial Index, by the Rev. W. W. SKEAT, M.A. Crown 8vo. 6s.

**Scott's (Sir Walter) LAY OF THE LAST MINSTREL; and THE LADY OF THE LAKE.** Edited, with Introduction and Notes, by FRANCIS TURNER PALGRAVE. Globe 8vo. 1s. (*Globe Readings from Standard Authors.*)

MARMION; and THE LORD OF THE ISLES. By the same Editor. Globe 8vo. 1s. (*Globe Readings from Standard Authors.*)

**Shakespeare.**—A SHAKESPEARE MANUAL, By F. G. FLEAY, M.A., late Head Master of Skipton Grammar School. Second Edition. Extra fcap. 8vo. 4s. 6d.

AN ATTEMPT TO DETERMINE THE CHRONOLOGICAL ORDER OF SHAKESPEARE'S PLAYS. By the Rev. H. PAINE STOKES, B.A. Extra fcap. 8vo. 4s. 6d.

THE TEMPEST. With Glossarial and Explanatory Notes. By the Rev. J. M. JEPHSON. New Edition. 18mo. 1s.

PRIMER OF SHAKESPEARE. By Professor DOWDEN. 18mo. 1s. (*Literature Primers.*)

**Sonnenschein and Meiklejohn.**—THE ENGLISH METHOD OF TEACHING TO READ. By A. SONNENSCHIEIN and J. M. D. MEIKLEJOHN, M.A. Fcap. 8vo.

COMPRISING :

THE NURSERY BOOK, containing all the Two-Letter Words in the Language. *id.* (Also in Large Type on Sheets for School Walls. 5s.)

THE FIRST COURSE, consisting of Short Vowels with Single Consonants. 6d.

THE SECOND COURSE, with Combinations and Bridges, consisting of Short Vowels with Double Consonants. 6d.

THE THIRD AND FOURTH COURSES, consisting of Long Vowels, and all the Double Vowels in the Language. 6d.

"These are admirable books, because they are constructed on a principle, and that the simplest principle on which it is possible to learn to read English."—SPECTATOR.

**Taylor.**—WORDS AND PLACES; or, Etymological Illustrations of History, Ethnology, and Geography. By the Rev. ISAAC TAYLOR, M.A. Third and Cheaper Edition, revised and compressed. With Maps. Globe 8vo. 6s.

**Thring.**—THE ELEMENTS OF GRAMMAR TAUGHT IN ENGLISH. By EDWARD THRING, M.A., Head Master of Uppingham. With Questions. Fourth Edition. 18mo. 2s.

**Trench (Archbishop).**—Works by R. C. TRENCH, D.D., Archbishop of Dublin.

HOUSEHOLD BOOK OF ENGLISH POETRY, Selected and Arranged, with Notes. Third Edition. Extra fcap. 8vo. 5s. 6d.

ON THE STUDY OF WORDS, Seventeenth Edition, revised. Fcap. 8vo. 5s.



**Trench (Archbishop)** Works by, *continued*—

ENGLISH, PAST AND PRESENT. Eleventh Edition, revised and improved. Fcap. 8vo. 5s.

A SELECT GLOSSARY OF ENGLISH WORDS, used formerly in Senses Different from their Present. Fifth Edition, revised and enlarged. Fcap. 8vo. 5s.

**Vaughan (C.M.)**.—WORDS FROM THE POETS. By C. M. VAUGHAN. New Edition. 18mo, cloth. 1s.

**Ward**.—THE ENGLISH POETS. Selections, with Critical Introductions by various Writers and a General Introduction by MATTHEW ARNOLD. Edited by T. H. WARD, M.A. 4 Vols. Vol. I. CHAUCER TO DONNE.—Vol. II. BEN JONSON TO DRYDEN.—Vol. III. ADDISON TO BLAKE.—Vol. IV. WORDSWORTH TO ROSSETTI. Crown 8vo. Each 7s. 6d.

**Wetherell**.—EXERCISES ON MORRIS'S PRIMER OF ENGLISH GRAMMAR. By JOHN WETHERELL, M.A. 18mo. 1s. (*Literature Primers*.)

**Wrightson**.—THE FUNCTIONAL ELEMENTS OF AN ENGLISH SENTENCE, an Examination of. Together with a New System of Analytical Marks. By the Rev. W. G. WRIGHTSON, M.A., Cantab. Crown 8vo. 5s.

**Yonge (Charlotte M.)**.—THE ABRIDGED BOOK OF GOLDEN DEEDS. A Reading Book for Schools and general readers. By the Author of "The Heir of Redclyffe." 18mo, cloth. 1s.

GLOBE READINGS EDITION. Complete Edition. Globe 8vo. 2s. (See p. 52.)

## FRENCH.

**Beaumarchais**.—LE BARBIER DE SEVILLE. Edit with Introduction and Notes, by L. P. BLOUET, Assistant Ma in St. Paul's School. Fcap. 8vo. 3s. 6d.

**Bowen**.—FIRST LESSONS IN FRENCH. By H. C THOPE BOWEN, M.A., Principal of the Finsbury Training C for Higher and Middle Schools. Extra fcap. 8vo. 1s.

**Breymann.**—Works by HERMANN BREYMAN, Ph.D., Professor of Philology in the University of Munich.

A FRENCH GRAMMAR BASED ON PHILOLOGICAL PRINCIPLES. Second Edition. Extra fcap. 8vo. 4s. 6d.

FIRST FRENCH EXERCISE BOOK. Extra fcap. 8vo. 4s. 6d.

SECOND FRENCH EXERCISE BOOK. Extra fcap. 8vo. 2s. 6d.

**Fasnacht.**—THE ORGANIC METHOD OF STUDYING LANGUAGES. By G. EUGÈNE FASNACHT, Author of "Macmillan's Progressive French Course," Editor of "Macmillan's Foreign School Classics," &c. Extra fcap. 8vo. I. French. 3s. 6d.

A SYNTHETIC FRENCH GRAMMAR FOR SCHOOLS. By the same Author. Crown 8vo. 3s. 6d.

**Macmillan's Progressive French Course.**—By G. EUGÈNE FASNACHT, sometime Senior Master of Modern Languages, Harpur Foundation Modern School, Bedford.

I.—FIRST YEAR, containing Easy Lessons on the Regular Accidence. Extra fcap. 8vo. 1s.

II.—SECOND YEAR, containing Conversational Lessons on Systematic Accidence and Elementary Syntax. With Philological Illustrations and Etymological Vocabulary. 1s. 6d.

III.—THIRD YEAR, containing a Systematic Syntax, and Lessons in Composition. Extra fcap. 8vo. 2s. 6d.

THE TEACHER'S COMPANION TO MACMILLAN'S PROGRESSIVE FRENCH COURSE. Third Year. By G. E. FASNACHT. Globe 8vo. [In the Press.]

**Macmillan's Progressive French Readers.**—By G. EUGÈNE FASNACHT.

I.—FIRST YEAR, containing Tables, Historical Extracts, Letters, Dialogues, Fables, Ballads, Nursery Songs, &c., with Two Vocabularies: (1) in the order of subjects; (2) in alphabetical order. Extra fcap. 8vo. 2s. 6d.

II.—SECOND YEAR, containing Fiction in Prose and Verse, Historical and Descriptive Extracts, Essays, Letters, Dialogues, &c. Extra fcap. 8vo. 2s. 6d.

**Macmillan's Foreign School Classics**—Edited by  
G. EUGÈNE FASNACHT, 18mo.

**FRENCH.**

CORNEILLE—LE CID. Edited by G. E. FASNACHT. 1s.

MOLIÈRE—LES FEMMES SAVANTES. By the same Editor.  
1s.

MOLIÈRE—LE MISANTHROPE. By the same Editor. 1s.

MOLIÈRE—LE MÉDECIN MALGRÉ LUI. By the same  
Editor. 1s.

MOLIÈRE—L'AVARE. Edited by L. M. MORIARTY, B.A.,  
Assistant-Master at Rossall. 1s.

MOLIÈRE—LE BOURGEOIS GENTILHOMME. By the same  
Editor. [In preparation.]

SELECTIONS FROM FRENCH HISTORIANS. Edited by  
C. COLBECK, M.A., late Fellow of Trinity College, Cambridge;  
Assistant Master at Harrow. [In preparation.]

SAND, GEORGE—LA MARE AU DIABLE. Edited by W. E.  
RUSSELL, M.A., Assistant Master in Haileybury College. 1s.

SANDEAU, JULES—MADEMOISELLE DE LA SEIGLIÈRE.  
Edited by H. C. STEEL, Assistant Master in Wellington College.  
[In the press.]

VOLTAIRE—CHARLES XII. Edited by G. E. FASNACHT.  
[In preparation.]

GRAMMAR AND GLOSSARY OF THE FRENCH LAN-  
GUAGE IN THE SEVENTEENTH CENTURY. By  
G. E. FASNACHT. [In preparation.]

\*\*\* Other volumes to follow.

(See also *German Authors*, page 61.)

**Masson (Gustave).**—A COMPENDIOUS DICTIONARY  
OF THE FRENCH LANGUAGE (French-English and English-  
French). Adapted from the Dictionaries of Professor ALFRED  
ELWALL. Followed by a List of the Principal Diverging  
Derivations, and preceded by Chronological and Historical Tables.  
By GUSTAVE MASSON, Assistant Master and Librarian, Harrow  
School. New Edition. Crown 8vo, half-bound. 6s.

**Molière.**—LE MALADE IMAGINAIRE. Edited, with Intro-  
duction and Notes, by FRANCIS TARVER, M.A., Assistant Master  
at Eton. Fcap. 8vo. 2s. 6d.

(See also *Macmillan's Foreign School Classics*.)

## GERMAN.

**Macmillan's Progressive German Course.**—By G. EUGÈNE FASNACHT.

Part I.—FIRST YEAR. Easy Lessons and Rules on the Regular Accidence. Extra fcap. 8vo. 1s. 6d.

Part II.—SECOND YEAR. Conversational Lessons in Systematic Accidence and Elementary Syntax. With Philological Illustrations and Etymological Vocabulary. Extra fcap. 8vo. 2s.

\* \* \* *Keys to the French and German Courses are in preparation.*

**Macmillan's Foreign School Classes.** Edited by G. EUGÈNE FASNACHT. 18mo.

## GERMAN.

GOETHE—GÖTZ VON BERLICHINGEN. Edited by H. A. BULL, M.A., Assistant Master at Wellington. 2s.

GOETHE—FAUST. PART I. Edited by JANE LEE, Lecturer in Modern Languages at Newnham College, Cambridge.

HEINE—SELECTIONS FROM THE PROSE WRITINGS. Edited by C. COLBECK, M.A. *[In preparation.]*

SCHILLER—DIE JUNGFRAU VON ORLEANS. Edited by JOSEPH GOSTWICK. 2s. 6d. *[In the press.]*

SCHILLER—MARIA STUART. Edited by C. SHELDON, M.A., D.Lit., Assistant Master in Clifton College. *[In the press.]*

SCHILLER—WILHELM TELL. Edited by G. E. FASNACHT. *[In preparation.]*

UHLAND—SELECT BALLADS. Adapted as a First Easy Reading Book for Beginners. Edited by G. E. FASNACHT. 1s. *[Ready.]*

SELECTIONS FROM GERMAN HISTORIANS. By the same Editor. Part I.—Ancient History. *[In preparation.]*

\* \* \* *Other Volumes to follow.*

(See also *French Authors*, page 60.)

**Pylozet.**—NEW GUIDE TO GERMAN CONVERSATION: containing an Alphabetical List of nearly 800 Familiar Words; followed by Exercises; Vocabulary of Words in frequent use; Familiar Phrases and Dialogues; a Sketch of German Literature, Idiomatic Expressions, &c. By L. PYLODET. 18mo, cloth limp. 2s. 6d.

**A SYNOPSIS OF GERMAN GRAMMAR.** From the above. 18mo. 6d.

**Whitney and Edgren.**—A COMPENDIOUS GERMAN AND ENGLISH DICTIONARY, with Notation of Correspondences and Brief Etymologies. By Professor W. D. WHITNEY, assisted by A. H. EDGREN. Crown 8vo. 7s. 6d.  
THE GERMAN-ENGLISH PART, separately, 5s.

### MODERN GREEK.

**Vincent and Dickson.**—HANDBOOK TO MODERN GREEK. By EDGAR VINCENT and T. G. DICKSON, M.A. Second Edition, revised and enlarged, with Appendix on the relation of Modern and Classical Greek by Professor JEBB. Crown 8vo. 6s.

### ITALIAN.

**Dante.**—THE PURGATORY OF DANTE. Edited, with Translation and Notes, by A. J. BUTLER, M.A., late Fellow of Trinity College, Cambridge. Crown 8vo. 12s. 6d.

### DOMESTIC ECONOMY.

**Barker.**—FIRST LESSONS IN THE PRINCIPLES OF COOKING. By LADY BARKER. New Edition. 18mo. 1s.

**Berners.**—FIRST LESSONS ON HEALTH. By J. BERNERS. New Edition. 18mo. 1s.

**Fawcett.**—TALES IN POLITICAL ECONOMY. By MILLICENT GARRETT FAWCETT. Globe 8vo. 3s.

**Frederick.**—HINTS TO HOUSEWIVES ON SEVERAL POINTS, PARTICULARLY ON THE PREPARATION OF ECONOMICAL AND TASTEFUL DISHES. By Mrs. FREDERICK. Crown 8vo. 1s.

"This unpretending and useful little volume distinctly supplies a desideratum. . . . The author steadily keeps in view the simple aim of 'making every-day meals at home, particularly the dinner, attractive,' without adding to the ordinary household expenses."—SATURDAY REVIEW.

**Grand'homme.**—CUTTING-OUT AND DRESSMAKING. From the French of Mdlle. E. GRAND'HOMME. With Diagrams. 18mo. 1s.

**Tegetmeier.**—HOUSEHOLD MANAGEMENT AND COOKERY. With an Appendix of Recipes used by the Teachers of the National School of Cookery. By W. B. TEGETMEIER. Compiled at the request of the School Board for London. 18mo. 1s.

**Thornton.**—FIRST LESSONS IN BOOK-KEEPING. By J. THORNTON. New Edition. Crown 8vo. 2s. 6d.

The object of this volume is to make the theory of Book-keeping sufficiently plain for even children to understand it.

**Wright.**—THE SCHOOL COOKERY-BOOK. Compiled and Edited by C. E. GUTHRIE WRIGHT, Hon Sec. to the Edinburgh School of Cookery. 18mo. 1s.

---

## ART AND KINDRED SUBJECTS.

**Anderson.**—LINEAR PERSPECTIVE, AND MODEL DRAWING. A School and Art Class Manual, with Questions and Exercises for Examination, and Examples of Examination Papers. By LAURENCE ANDERSON. With Illustrations. Royal 8vo. 2s.

**Collier.**—A PRIMER OF ART. With Illustrations. By JOHN COLLIER. 18mo. 1s.

**Delamotte.**—A BEGINNER'S DRAWING BOOK. By P. H. DELAMOTTE, F.S.A. Progressively arranged. New Edition improved. Crown 8vo. 3s. 6d.

**Ellis.**—SKETCHES FROM NATURE. A Handbook for Students and Amateurs. By TRISTRAM J. ELLIS. With a Frontispiece and Ten Illustrations, by H. STACY MARKS, R.A., and Twenty-seven Sketches by the Author. Crown 8vo. 2s. 6d. (*Art at Home Series.*)

**Hunt.**—TALKS ABOUT ART. By WILLIAM HUNT. With a Letter from J. E. MILLAIS, R.A. Crown 8vo. 3s. 6d.

**Taylor.**—A PRIMER OF PIANOFORTE PLAYING. By FRANKLIN TAYLOR. Edited by GEORGE GROVE. 18mo. 1s.

## WORKS ON TEACHING.

**Blakiston**—THE TEACHER. Hints on School Management. A Handbook for Managers, Teachers' Assistants, and Pupil Teachers. By J. R. BLAKISTON, M.A. Crown 8vo. 2s. 6d. (Recommended by the London, Birmingham, and Leicester School Boards.)

"Into a comparatively small book he has crowded a great deal of exceedingly useful and sound advice. It is a plain, common-sense book, full of hints to the teacher on the management of his school and his children."—SCHOOL BOARD CHRONICLE.

**Calderwood**—ON TEACHING. By Professor HENRY CALDERWOOD. New Edition. Extra fcap. 8vo. 2s. 6d.

**Fearon**—SCHOOL INSPECTION. By D. R. FEARON, M.A., Assistant Commissioner of Endowed Schools. New Edition. Crown 8vo. 2s. 6d.

**Gladstone**—OBJECT TEACHING. A Lecture delivered at the Pupil-Teacher Centre, William Street Board School, Hammersmith. By J. H. GLADSTONE, Ph.D., F.R.S., Member of the London School Board. With an Appendix. Crown 8vo. 3d.

"It is a short but interesting and instructive publication, and our younger teachers will do well to read it carefully and thoroughly. There is much in these few pages which they can learn and profit by."—THE SCHOOL GUARDIAN.

---

## DIVINITY.

\*.\* For other Works by these Authors, see THEOLOGICAL CATALOGUE.

**Abbott (Rev. E. A.)**—BIBLE LESSONS. By the Rev. E. A. ABBOTT, D.D., Head Master of the City of London School. New Edition. Crown 8vo. 4s. 6d.

"Wise, suggestive, and really profound initiation into religious thought."—GUARDIAN.

**Arnold.**—A BIBLE-READING FOR SCHOOLS.—THE GREAT PROPHECY OF ISRAEL'S RESTORATION (Isaiah, Chapters xl.—lxvi.). Arranged and Edited for Young Learners. By MATTHEW ARNOLD, D.C.L., formerly Professor of Poetry in the University of Oxford, and Fellow of Oriel. New Edition. 18mo, cloth. 1s.

ISAIAH XL.—LXVI. With the Shorter Prophecies allied to it. Arranged and Edited, with Notes, by MATTHEW ARNOLD. Crown 8vo. 5s.

**Cheetham.**—A CHURCH HISTORY OF THE FIRST SIX CENTURIES. By the Ven. ARCHDEACON CHEETHAM. Crown 8vo. [In the press.]

**Curteis.**—MANUAL OF THE THIRTY-NINE ARTICLES. By G. H. CURTEIS, M.A., Principal of the Lichfield Theological College. [In preparation.]

**Gaskoin.**—THE CHILDREN'S TREASURY OF BIBLE STORIES. By Mrs. HERMAN GASKOIN. Edited with Preface by the Rev. G. F. MACLEAR, D.D. PART I.—OLD TESTAMENT HISTORY. 18mo. 1s. PART II.—NEW TESTAMENT. 18mo. 1s. PART III.—THE APOSTLES: ST. JAMES THE GREAT, ST. PAUL, AND ST. JOHN THE DIVINE. 18mo. 1s.

**Golden Treasury Psalter.**—Students' Edition. Being an Edition of "The Psalms Chronologically arranged, by Four Friends," with briefer Notes. 18mo. 3s. 6d.

**Greek Testament.**—Edited, with Introduction and Appendices, by CANON WESTCOTT and Dr. F. J. A. HORT. Two Vols. Crown 8vo. 10s. 6d. each.

Vol. I. The Text.

Vol. II. Introduction and Appendix.

**Greek Testament.**—Edited by Canon WESTCOTT and Dr. HORT. School Edition of Text. Globe 8vo. [In the press.]

**Hardwick.**—Works by Archdeacon HARDWICK:—

A HISTORY OF THE CHRISTIAN CHURCH. Middle Age. From Gregory the Great to the Excommunication of Luther. Edited by WILLIAM STUBBS, M.A., Regius Professor of Modern History in the University of Oxford. With Four Maps. Fourth Edition. Crown 8vo. 10s. 6d.



**Hardwick.**—Works by Archdeacon HARDWICK (*continued*)—

A HISTORY OF THE CHRISTIAN CHURCH DURING THE REFORMATION. Fourth Edition. Edited by Professor STUBBS. Crown 8vo. 10s. 6d.

**Jennings and Lowe.**—THE PSALMS, WITH INTRODUCTIONS AND CRITICAL NOTES. By A. C. JENNINGS, B.A.; assisted in parts by W. H. LOWE. In 2 vols. Crown 8vo. 10s. 6d. each.

**Lightfoot.**—Works by Right Rev. J. B. LIGHTFOOT, D.D., Bishop of Durham:—

ST. PAUL'S EPISTLE TO THE GALATIANS. A Revised Text, with Introduction, Notes, and Dissertations. Seventh Edition, revised. 8vo. 12s.

ST. PAUL'S EPISTLE TO THE PHILIPPIANS. A Revised Text, with Introduction, Notes, and Dissertations. Seventh Edition, revised. 8vo. 12s.

ST. CLEMENT OF ROME—THE TWO EPISTLES TO THE CORINTHIANS. A Revised Text, with Introduction and Notes. 8vo. 8s. 6d.

ST. PAUL'S EPISTLES TO THE COLOSSIANS AND TO PHILEMON. A Revised Text, with Introductions, Notes, and Dissertations. Sixth Edition, revised. 8vo. 12s.

THE IGNATIAN EPISTLES. 8vo. [In the press.]

**Maclear.**—Works by the Rev. G. F. MACLEAR, D.D., Warden of St. Augustine's College, Canterbury, and late Head-Master of King's College School, London:—

A CLASS-BOOK OF OLD TESTAMENT HISTORY. New Edition, with Four Maps. 18mo. 4s. 6d.

A CLASS-BOOK OF NEW TESTAMENT HISTORY, including the Connection of the Old and New Testaments. With Four Maps. New Edition. 18mo. 5s. 6d.

A SHILLING BOOK OF OLD TESTAMENT HISTORY, for National and Elementary Schools. With Map. 18mo, cloth. New Edition.

A SHILLING BOOK OF NEW TESTAMENT HISTORY, for National and Elementary Schools. With Map. 18mo, cloth. New Edition.

**Maclear.**—Works by the Rev. G. F. MACLEAR, D.D., *continued*—

These works have been carefully abridged from the author's large manuals.

CLASS-BOOK OF THE CATECHISM OF THE CHURCH OF ENGLAND. New Edition. 18mo, cloth. 1s. 6d.

A FIRST CLASS-BOOK OF THE CATECHISM OF THE CHURCH OF ENGLAND. With Scripture Proofs, for Junior Classes and Schools. New Edition. 18mo. 6d.

A MANUAL OF INSTRUCTION FOR CONFIRMATION AND FIRST COMMUNION. WITH PRAYERS AND DEVOTIONS. 32mo, cloth extra, red edges. 2s.

**Maurice.**—THE LORD'S PRAYER, THE CREED, AND THE COMMANDMENTS. A Manual for Parents and Schoolmasters. To which is added the Order of the Scriptures. By the Rev. F. DENISON MAURICE, M.A. 18mo, cloth, limp. 1s.

**Procter.**—A HISTORY OF THE BOOK OF COMMON PRAYER, with a Rationale of its Offices. By Rev. F. PROCTER, M.A. Sixteenth Edition, revised and enlarged. Crown 8vo. 10s. 6d.

**Procter and Maclear.**—AN ELEMENTARY INTRODUCTION TO THE BOOK OF COMMON PRAYER. Rearranged and supplemented by an Explanation of the Morning and Evening Prayer and the Litany. By the Rev. F. PROCTER and the Rev. Dr. MACLEAR. New and Enlarged Edition, containing the Communion Service and the Confirmation and Baptismal Offices. 18mo. 2s. 6d.

**The Psalms, with Introductions and Critical Notes.**—By A. C. JENNINGS, B.A., Jesus College, Cambridge, Tyrwhitt Scholar, Crosse Scholar, Hebrew University, Prizeman, and Fry Scholar of St. John's College; assisted in Parts by W. H. LOWE, M.A., Hebrew Lecturer and late Scholar of Christ's College, Cambridge, and Tyrwhitt Scholar. In 2 vols. Crown 8vo. 10s. 6d. each.

**Ramsay.**—THE CATECHISER'S MANUAL; or, the Church Catechism Illustrated and Explained, for the Use of Clergymen, Schoolmasters, and Teachers. By the Rev. ARTHUR RAMSAY, M.A. New Edition. 18mo. 1s. 6d.

**Simpson.**—AN EPITOME OF THE HISTORY OF THE CHRISTIAN CHURCH. By WILLIAM SIMPSON, M.A. New Edition. Fcap. 8vo. 3s. 6d.

**St. John's Epistles.**—The Greek Text with Notes and Essays, by BROOKE FOSS WESTCOTT, D.D., Regius Professor of Divinity and Fellow of King's College, Cambridge, Canon of Peterborough, &c. 8vo. 12s. 6d.

**St. Paul's Epistles.**—Greek Text, with Introduction and Notes.

THE EPISTLE TO THE GALATIANS. Edited by the Right Rev. J. B. LIGHTFOOT, D.D., Bishop of Durham. Seventh Edition. 8vo. 12s.

THE EPISTLE TO THE PHILIPPIANS. By the same Editor. Seventh Edition. 8vo. 12s.

THE EPISTLE TO THE COLOSSIANS. By the same Editor. Sixth Edition. 8vo. 12s.

THE EPISTLE TO THE ROMANS. Edited by the Very Rev. C. J. VAUGHAN, D.D., Dean of Llandaff, and Master of the Temple. Fifth Edition. Crown 8vo. 7s. 6d.

THE EPISTLE TO THE THESSALONIANS, COMMENTARY ON THE GREEK TEXT. By JOHN EADIE, D.D., LL.D. Edited by the Rev. W. YOUNG, M.A., with Preface by Professor CAIRNS. 8vo. 12s.

**The Epistle to the Hebrews.** In Greek and English. With Critical and Explanatory Notes. Edited by Rev. FREDERIC RENDALL, M.A., formerly Fellow of Trinity College, Cambridge, and Assistant-Master at Harrow School. Crown 8vo. 6s.

**Trench.**—Works by R. C. TRENCH, D.D., Archbishop of Dublin.

NOTES ON THE PARABLES OF OUR LORD. Fourteenth Edition, revised. 8vo. 12s.

NOTES ON THE MIRACLES OF OUR LORD. Eleventh Edition, revised. 8vo. 12s.

COMMENTARY ON THE EPISTLES TO THE SEVEN CHURCHES IN ASIA. Third Edition, revised. 8vo. 8s. 6d.

LECTURES ON MEDIEVAL CHURCH HISTORY. Being the substance of Lectures delivered at Queen's College London. Second Edition, revised. 8vo. 12s.

SYNONYMS OF THE NEW TESTAMENT. Ninth Edition, revised. 8vo. 12s.

**Westcott.**—Works by BROOKE FOSS WESTCOTT, D.D., Canon of Peterborough, Regius Professor of Divinity, and Fellow of King's College, Cambridge.

A GENERAL SURVEY OF THE HISTORY OF THE CANON OF THE NEW TESTAMENT DURING THE FIRST FOUR CENTURIES. Fifth Edition. With Preface on "Supernatural Religion." Crown 8vo. 10s. 6d.

INTRODUCTION TO THE STUDY OF THE FOUR GOSPELS. Fifth Edition. Crown 8vo. 10s. 6d.

THE BIBLE IN THE CHURCH. A Popular Account of the Collection and Reception of the Holy Scriptures in the Christian Churches. New Edition. 18mo, cloth. 4s. 6d.

THE EPISTLES OF ST. JOHN. The Greek Text, with Notes and Essays. 8vo. 12s. 6d.

THE EPISTLE TO THE HEBREWS. The Greek Text Revised, with Notes and Essays. 8vo. *[In preparation.]*

**Westcott and Hort.**—THE NEW TESTAMENT IN THE ORIGINAL GREEK. The Text Revised by B. F. WESTCOTT, D.D., Regius Professor of Divinity, Canon of Peterborough, and F. J. A. HORT, D.D., Hulsean Professor of Divinity; Fellow of Emmanuel College, Cambridge: late Fellows of Trinity College, Cambridge. 2 vols. Crown 8vo. 10s. 6d. each.

Vol. I. Text.

Vol. II. Introduction and Appendix.

**Wilson.**—THE BIBLE STUDENT'S GUIDE to the more Correct Understanding of the English Translation of the Old Testament, by reference to the original Hebrew. By WILLIAM WILSON, D.D., Canon of Winchester, late Fellow of Queen's College, Oxford. Second Edition, carefully revised. 4to. cloth. 25s.

**Yonge (Charlotte M.).**—SCRIPTURE READINGS FOR SCHOOLS AND FAMILIES. By CHARLOTTE M. YONGE. Author of "The Heir of Redclyffe." In Five Vols.

FIRST SERIES: GENESIS TO DEUTERONOMY. Extra fcap. 8vo. 1s. 6d. With Comments, 3s. 6d.

SECOND SERIES. From JOSHUA to SOLOMON. Extra fcap. 8vo. 1s. 6d. With Comments, 3s. 6d.

THIRD SERIES. The KINGS and the PROPHETS. Extra fcap.  
8vo. 1s. 6d. With Comments, 3s. 6d.

FOURTH SERIES. The GOSPEL TIMES. 1s. 6d. With Comments,  
extra fcap. 8vo, 3s. 6d.

FIFTH SERIES. APOSTOLIC TIMES Extra fcap. 8vo. 1s. 6d.  
With Comments, 3s. 6d.

**Zechariah—Lowe.**—THE HEBREW STUDENT'S COMMENTARY ON ZECHARIAH, HEBREW AND LXX. With Excursus on Syllable-dividing, Metheg, Initial Dagesh, and Siman Rapheh. By W. H. LOWE, M.A., Hebrew Lecturer at Christ's College, Cambridge. Demy 8vo. 10s. 6d.

## MACMILLAN'S GLOBE LIBRARY.

*In Cloth Binding, Globe 8vo, 3s. 6d. each.*

"The 'Globe' Editions are admirable for their scholarly editing, their typographical excellence, their compendious form, and their cheapness."—*Saturday Review*.

"Not only truly cheap, but excellent in every way."—*Literary World*.

"A wonderfully cheap and scholarly series."—*Daily Telegraph*.

**SHAKESPEARE'S COMPLETE WORKS.** Edited by W. G. CLARK, M.A., and W. ALDIS WRIGHT, M.A. With Glossary.

**MORTE D'ARTHUR.** The Book of King Arthur and of his Noble Knights of the Round Table. The Original Edition of Caxton revised for Modern Use, with Introduction, Notes, and Glossary. By Sir E. STRACHEY.

**ROBINSON CRUSOE.** Edited after the Original Editions. With a Biographical Introduction by HENRY KINGSLEY, F.R.G.S.

**SIR WALTER SCOTT'S POETICAL WORKS.** Edited with Biographical and Critical Memoir, by F. T. PALGRAVE. With Introduction and Notes.

**VIRGIL.** Rendered into English Prose, with Introductions, Notes, Analysis, and Index by J. LONSDALE, M.A., and S. LEE, M.A.

**HORACE.** Rendered into English Prose, with Introductions, Running Analysis, Notes, and Index, by J. LONSDALE, M.A., and S. LEE, M.A.

**DRYDEN'S POETICAL WORKS.** Edited, with a Memoir, Revised Text, and Notes, by W. D. CHRISTIE, M.A.

**COWPER'S POETICAL WORKS.** Edited, with Biographical Introduction and Notes, by W. BENHAM, B.D.

**BURNS'S COMPLETE WORKS.** Edited from the best Printed and MS. Authorities, with Memoir and Glossarial Index, by ALEXANDER SMITH.

**GOLDSMITH'S MISCELLANEOUS WORKS.** With Biographical Introduction by Professor MASSON.

**POPE'S POETICAL WORKS.** Edited, with Notes and Introductory Memoir, by Professor WARD, of Owen's College, Manchester.

**SPENSER'S COMPLETE WORKS.** Edited from the Original Editions and Manuscripts, with Glossary, by R. MORRIS, and a Memoir by J. W. HALES, M.A.

**MILTON'S POETICAL WORKS.** Edited, with Introductions, by Professor MASSON.

MACMILLAN AND CO., LONDON.

Now Publishing, in Crown 8vo, price 2s. 6d. each.  
Also in stiff boards, uncut edges, price 2s. 6d. each.

## ENGLISH MEN OF LETTERS.

EDITED BY JOHN MORLEY.

---

"These excellent biographies should be made class-books for Schools."—*Westminster Review*.

"This admirable series"—*British Quarterly Review*.

"Enjoyable and excellent little books."—*Academy*.

---

**JOHNSON.** By LESLIE STEPHEN.  
**SCOTT.** By R. H. HUTTON.  
**GIBBON.** By J. C. MORISON.  
**SHELLEY.** By J. A. SYMONDS.  
**HUME.** By Professor HUXLEY.  
P.R.S.  
**GOLDSMITH.** By WILLIAM BLACK.  
**DEFOE.** By W. MINTO.  
**BURNS.** By Principal SHAIRP.  
**SPENSER.** By the Very Rev. the  
DEAN OF ST. PAUL'S.  
**THACKERAY.** By ANTHONY  
TROLLOPE.  
**BURKE.** By JOHN MORLEY.  
**BUNYAN.** By J. A. FROUDE.  
**POPE.** By LESLIE STEPHEN.  
**BYRON.** By Professor NICHOL.  
**COWPER.** By GOLDWIN SMITH.  
**LOCKE.** By Professor FOWLER.  
**WORDSWORTH.** By F. W. H.  
MYERS.

**DRYDEN.** By G. SAINTSBURY.  
**LANDOR.** By Professor SIDNEY  
COLVIN.  
**CHARLES LAMB.** By Rev. A.  
AINGER.  
**BENTLEY.** By Professor R. C.  
JEBB.  
**DICKENS.** By Prof. A. W. WARD.  
**MACAULAY.** By J. C. MORISON.  
**DE QUINCEY.** By Prof. MASSON.  
**MILTON.** By MARK PATTISON.  
**HAWTHORNE.** By HENRY JAMES.  
**SOUTHEY.** By Professor DOWDEN.  
**CHAUCE.** By Prof. A. W. WARD.  
**GRAY.** By E. W. GOSSE.  
**SWIFT.** By LESLIE STEPHEN.  
**STERNE.** By H. D. TRAILL.  
**FIELDING.** By AUSTIN DORSON.  
**SHERIDAN.** By Mrs. OLIPHANT.

\* \* \* Other Volumes to follow.

---

*Macmillan & Co.'s New Illustrated Catalogue of Books suitable for  
PRESENTATION and SCHOOL PRIZES, at prices ranging  
from SIXPENCE upwards, may now be had, post free.*

---

MACMILLAN AND CO., LONDON.





